

## eCommons@AKU

Department of Pathology and Laboratory Medicine

Medical College, Pakistan

October 2001

# Morphological pattern of salivary gland tumours

M. S. Gill Aga Khan University

S. Muzaffar Aga Khan University

I. N. Soomro Aga Khan University

Naila Kayani Aga Khan University

A. S. Hussainy *Aga Khan University* 

See next page for additional authors

Follow this and additional works at: http://ecommons.aku.edu/ pakistan\_fhs\_mc\_pathol\_microbiol

Part of the <u>Digestive System Diseases Commons</u>, <u>Oncology Commons</u>, and the <u>Pathology</u> <u>Commons</u>

#### **Recommended** Citation

Gill, M. S., Muzaffar, S., Soomro, I. N., Kayani, N., Hussainy, A. S., Pervez, S., Hasan, S. H. (2001). Morphological pattern of salivary gland tumours. *Journal of Pakistan Medical Association*, *51*(10), 343-346. **Available at:** http://ecommons.aku.edu/pakistan\_fhs\_mc\_pathol\_microbiol/286

#### Authors

M. S. Gill, S. Muzaffar, I. N. Soomro, Naila Kayani, A. S. Hussainy, Shahid Pervez, and S. H. Hasan

# **Morphological Pattern of Salivary Gland Tumours**

M. S. Gill, S. Muzaffar, I. N. Soomro, N. Kayani, A. S. Hussainy, S. Pervez, S. H. Hasan (Department of Pathology, The Aga Khan University Hospital, Karachi.)

#### Abstract

**Objective:** To delineate the spectrum of salivary gland tumors in our setup. **Setting:** The Aga Khan University Medical Centre, Karachi.

**Method:** Tumors were analysed considering histological type, age and sex of the patients and anatomic location. The diagnosis of individual tumours was based on the 1991 World Health Organisation Classification.

**Results:** During the span of eight years (1991-1998), 379 cases of salivary gland tumours were diagnosed. Of these, 205 (65.7%) were male and 174 (34.3%) were female. The median age at the time of diagnosis was 35 years. The median age for patients with malignant lesions (44 years) was 12 years older than those with benign tumours (34 years). Overall, malignant tumours were seen more frequently in males, however benign tumours were distributed equally between the two sexes. The most common site was parotid gland (82.85%). Only five cases of minor salivary gland tumours were seen. The most frequently diagnosed benign salivary gland neoplasm was pleomorphic adenoma (84.5%), followed by Warthin's tumours (6.18%), Mucoepidermoid carcinoma was the most commonly encountered malignant lesion (56.9%), followed by adenoid cystic carcinoma (19.6%). **Conclusion:** Plemorphic adenoma was the most frequent malignant neoplasm. Parotid gland was the most common site of origin in both benign and malignant tumours. The overall relative frequency of salivary gland tumours in this series correlates with that reported in the international literature (JPMA 51:343,2001).

#### Introduction

Tumours of the salivary glands are relatively infrequent and account for less than 2% of all human tumours<sup>1,2</sup>. These comprise a wide variety of benign and malignant neoplasms, which exhibit difference not only in biological behavior but in prognosis as well<sup>3-6</sup>. Overall, benign tumours of the salivary glands tend to present somewhat earlier than malignant ones<sup>6</sup>.

Many studies have been carried out in the West<sup>3-6</sup>, however published data in Pakistan is scanty. The risk factor and demographics of our population are different which may lead to a different age of the population at risk. Therefore, this series describes the prevalence of salivary tumours, in a hospital-based sample and characterizes them according to age, sex and anatomic location.

#### **Materials and Methods**

Data was collected from the surgical pathology files of the histopathology section of pathology department of the Aga Khan University Medical Centre. The study included all cases of salivary tumours diagnosed during the eight-year period, 1991 to 1998. Cases were then characterized according to their histological subtypes, age, sex and anatomic location of the tumours. The histological evaluation of these lesion was performed on Hematoxylin and Eosin stained Sections of paraffin embedded tissue. Special stains like Periodic Acid Schiff (PAS) with and without diastase, PAS with Alcian Blue Reticulin were done wherever indicated. The immun oh istochem ical evaluation,

if required, was performed using Peroxidase Anti-Peroxidase (PAP) technique. The monoclonal antibodies used include Cytokeratins, Vimentin, Leukocyte Common Antigen (LCA), Pan-B (L-26, CD 20), Pan T (UCHL 1), S-100 protein and Alpha Smooth Muscle Antigen (ASMA) depending upon the nature of lesion. The diagnosis of individual tumours was based on the 1991 World Health Organization Classification. Patients with more than one specimen from the same lesion were counted once.

#### Results

During the eight-year period, 1991-1998, 379 cases of salivary tumours were diagnosed at the histopathology section of pathology department. Of these, 205 (54.09%) were male and 174 (45.9 1%) were female, giving a male to female ratio of 1.2:1. There were more men (65.7%) than women (34.3%) affected by malignant tumours, but benign tumours were equally distributed between the two sexes.

Histological subtype	No. of cases	%
Benign	277	73.09
Pleomorphic Adenoma	234	61.74
Warthin's Tumour	17	4.49
Cavernous Hemangioma	5	1.32
Basal Cell Adenoma	4	1.06
Lipoma	4	1.06
Schwannoma	4	1.06
Capillary Hemangioma	3	0.79
Monomorphic Adenoma	2	0.53
Trabecular Adenoma	1	0.26 0.26
Angiolipoma	1	
Myoepithelioma	1	0.26
Hamartoma	1	0.26
Malignant	102	26.91
Mucoepidermoid Carcinoma	58	15.30
Adenoid Cystic Carcinoma	20	5.28
Malignant Mixed Tumour	5	1.32
Undifferentiated Carcinoma	5	1.32
Acinic Cell Carcinoma	4	1.06
Non-Hodgkin's Lymphoma	. 4	1.06
Salivary Duct Carcinoma	2	0.53
Adenocarcinoma	1	0.26
Papillary Adenocarcinoma	1	0.26
Metastatic Leukemia	1	0.26
Hodgkin's Disease	1	0.26

Table 1. Distribution of histological entities in 379 cases.

Table 1 describes the relative distribution of various histological subtypes of salivary tumours. Of these, 73 .09% were classified as benign and 26.9 1% as malignant.

Pleomorphic adenoma was the most frequently diagnosed benign salivary neoplasm (6 1.74% of all cases and 84.5% of benign tumours), followed by Warthin's tumours (4.49% of all tumours and 6.14%

of benign lesions). Mucoepidermoid carcinoma was the most commonly encountered malignant tumour, accounting for 15.3% of all. tumours and 56.9% of malignant lesions. Adenoid cystic carcinoma was the second most common type, making up 5.28% of all tumours and 19.6% of all malignant tumours. Many other categories were also represented in relatively smaller numbers.

Age (Years)	Benign		Malignant		Total	
	No.	%	No.	%	No.	%
0-9	6	2.17	1	0. 98	7	1.85
10-19	35	12.64	10	9.80	45	11.87
20-29	73	26.35	15	14.70	88	23.22
30-39	63	22.74	12	11.77	75	19.79
40-49	53	19.13	21	20. 59	74	19. 52
50-59	29	10.47	20	19.61	49	12.93
60-69	11	3.97	12	11.77	23	6.07
70+	3	1.08	8	7.84	11	2.90
Unknown	4	1.45	3	2.94	7	1.85

Table 2. Distribution of salivary tumours by Age.

Table 2 summarises the distribution of salivary tumours in various age groups. Majority (54.9%) of the salivary tumours were diagnosed during 2nd to 4th decades of life. The median age at diagnosis was 35 years, with a range of 1-79 years. The median age for patients with malignant lesions (44 years) was 12 years older than for patients with benign tumours (34 years).

Regarding anatomic distribution of salivary gland tumours, three hundred and fourteen (82.85%) originated in the parotid glands, forty-seven (12.4%) in submandibular glands, three (0.8%) in minor salivary glands and two (0.52%) in sublingual salivary glands. Site of origin was not kown in thirteen

(3.43%) cases. Overall, there was a slight predominanace of left-sided tumours.

The four most common sub-categories were then analysed on the basis of age, sex and location. There were more women than men affected by pleomorphic adenoma and adenoid cystic carcinoma. However, a predominance of males was seen in mucoepidemoid carcinoma and Warthin's tumours. In each morphological subtype, parotid gland was the most common site of the tumour. The median age varied greatly, with the youngest (32 years) for pleomorphic adenoma and the eldest (50 years) for Warthin's tumour.

## Discussion

Tumours of the salivary glands are relatively uncommon, however, their multifaceted clinical presentation, varied morphologic configuration and relatively unpredictable prognosis continue to attract significant medical interest<sup>7</sup>. Environmental and hereditary factors have a considerable impact upon the geographical distribution of cancer<sup>8</sup>.

In the present study, the tumours of the salivary glands were more common in males especially the malignant lesions. High male to female ratio has been reported<sup>2,9-11</sup>. This finding contrasts with the overall female predominance reported in Western Literature<sup>1</sup>. In our series, benign tumours, on the average, presented a decade earlier than malignant tumours, is in accordance with the published literature<sup>3-6</sup>.

More than 90% of the tumours in our study were of epithelial type and parotid was the most commonly involved gland, similar to the published literature  $^{1,6,12,13}$ .

In this study, Pleomorphic adenoma was the most frequent histological type, mostly in the 3rd decade of life. It compromised 61% of tumours correlating well with other series<sup>3,6</sup>. Warthin's tumour was the second common benign neoplasm.

Amongst malignant tumours, mucoepidermoid carcinoma was the most common lesion in our series, predominantly affecting the parotid gland (85.7%). Most of these were seen in 4th, 5th and 6th decades. This was followed by adenoid cystic carcinoma which is the most common malignant tumour of minor salivary glands<sup>14</sup>, however in our series, 80% of these were seen in parotid. There were only five cases of minor salivary tumours in our study, accounting for this discordant finding. Hemangiomas are occasionally seen in the parotid<sup>15</sup>. In our study, these vascular neoplasms comprised 2% all parotid tumours.

Six cases of lipoma (including one case of angiolipoma) were seen in our study. Literature review reveals that these are rare neoplasms of parotid, however occasional cases have been reported in the submandibular gland<sup>16,17</sup>. Schwannoma presenting as primary salivary gland neoplasm is a recognized entity<sup>18</sup>. These are thought to arise from the radicals of facial nerve. We observed four cases of Schwannoma, all affecting parotid gland.

Malignant lymphomas of the salivary glands may either be a primary process or a secondary involvement by extraglandular diseases<sup>14</sup>. The incidence of primary lymphoma of the salivary glands ranges from 0.6 to  $2.4\%^{19,20}$ . In our study, there were four cases of non-Hodgkin's lymphoma and one case of Hodgkin's disease. As this was a retrospective analysis, the primary or secondary nature of these neoplasms could not be determined.

Plemorphic adenoma was the most common benign salivary gland tumour and mucoepidermoid carcinoma was the most frequent malignant neoplasm. Parotid gland was the most common site of origin in both benign and malignant tumours. The overall relative frequency of salivary gland tumours in this series correlates with that reported in the international literature.

## References

1.Cotran RS, Kwnar R, Collins T. Pathologic basis of disease. 6th ed. Philadelphia, W. B. Saunders Co., 1999,p.769.

2.Bhurgri Y, Bhurgn A, Rahim, et at. The pattern of malignancies in Karachi (1995 - 1996). 3. Pak. Med. Assoc., 1999;49:157-61.

3.Ellis GL, Auclair PL. Tumors of the Salivary Glands. Atlas of Tumor Pathology, Third Series, Fascicle 17. Washington, DC, Armed Forces Inst. of Pathology., 1996.

4.Simpson RH. Classification of salivary gland tumors: a brief pathological review. Histol. Histopathol., 1995;10:737.

5.Shrestha P. Primary epithelial tumors of salivay glands- histogenesis, histomorphological and immunohistochemical implications, diagnosis and clinical management. Crit. Rev. Oncol. Hematol., 1996;23:239.

6.Nagler RM, Laufer D. Tumors of the major and minor salivary glands: review of 25 years of experience. Anticancer Res., 1997;17:701.

7.Huvos AG. Salivary glands In: Sternberg SS(ed). Diagnostic surgical pathology. 2nd ed. Philadelphia;Lippincott-Raven, 1994,p.813.

8.Magrath I, Litrak J. Cancer in developing countries: opportunity and challenge. J. Nat. Cancer Inst., 1993;85:862-74.

9.Ahmad J, Hashmi MA, Naveed IA, et al. Spectrwn of malignanacies, in Faisalabad., 1986-1990. Pak. J. Pathol., 1992,3:103-10.

10.Jafarey NA. Frequency of malignant tumors in seven centres of Pakistan. Pakistan Medical Research Council Cancer Study Group. J. Pak. Med. Assoc., 1977;27:335-39.

11, Khan SM, Gillani J, Nasreen 5, et al. Cancer in North-West Pakistan. Pak. J. Med. Res., 1996;35:167-69.

12.Eveson JW, Cawson RA. Salivary gland tumors: a review of 2410 cases with particular reference to histological types, site, age, and sex distribution. J. Pathol., 1985;146:51-8.

13.Spiro RH. Salivary neoplasms: overview of a 35-year experience with 2807 patients. Head Neck Surg., 1986;8:177-84.

14.Rosai J. Major and minor salivary glands in: Rosai J(ed). Ackennan's surgical pathology. 8th ed. Missouri; Mosby., 1996;pp.837-56.

15.Mantravadi J, Roth LM, Kafrawy AH. Vascular neoplasms of the partoid gland: parotid vascular tumors. Oral Surg. Oral Med. Oral Pathol., 1 993;75:70-5.

16.Muzaffar S, Kayani N, Hasan SH. Partoid gland lipoma: a rare entity. J. Pak. Med. Assoc., 1996;46:262-63.

17.Maaita JK, Al-Qaisi N. Submandibular salivary gland masses: a clinicopathological study. JCPSP. 1999;9:268-70.

18.Katz AD, Passy V, Kaplan L. Neurogenous neoplasms of major nerves of face and neck. Arch. Surg., 1971;103:51-56.

19.Nichols RD. Rebuck JW, Sullivan JC. Lymphoma and the partoid gland. Laryngoscope., 198;92:365-69.

20.Wolvulus EB, Vander valk P, Vander Wal JE, et al. Primary non-Hodgkin's lymphoma of salivary gland: an analysis of 22 cases. J. Oral Path, Med., 1996;25:177-81.