



THE AGA KHAN UNIVERSITY

eCommons@AKU

Anaesthesiology, East Africa

Medical College, East Africa

January 2008

Knowledge, attitude and use of labour pain relief methods among women attending antenatal clinic in Nairobi

Vitalis Mung'ayi

Aga Khan University, vitalis.mung'ayi@aku.edu

D. Netyton

Aga Khan University

Robinson Karuga

Aga Khan University

Follow this and additional works at: http://ecommons.aku.edu/eastafrica_fhs_mc_anaesth



Part of the [Anesthesiology Commons](#)

Recommended Citation

Mung'ayi, V., Netyton, D., Karuga, R. (2008). Knowledge, attitude and use of labour pain relief methods among women attending antenatal clinic in Nairobi. *East African Medical Journal*, 85(9), 438-441.

Available at: http://ecommons.aku.edu/eastafrica_fhs_mc_anaesth/11

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/26302310>

Knowledge, Attitude and Use of Labour Pain Relief Methods among Women Attending Antenatal Clinic in Nairobi

Article in *East African medical journal* · September 2008

DOI: 10.4314/eamj.v85i9.117084 · Source: PubMed

CITATIONS

16

READS

409

3 authors, including:



Vitalis Mung'ayi

Aga Khan University Hospital, Nairobi

15 PUBLICATIONS 46 CITATIONS

[SEE PROFILE](#)



Robinson Njoroge Karuga

LVCT Health

11 PUBLICATIONS 36 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Effects of intravenous diclofenac on postoperative sore throat in patients undergoing laparoscopic surgery at Aga Khan University Hospital, Nairobi: A prospective, randomized, double blind controlled trial [View project](#)



Introducing Pre-Exposure Prophylaxis in Combination Prevention (IPCP) [View project](#)

East African Medical Journal Vol. 85 No. 9 September 2008

KNOWLEDGE, ATTITUDE AND USE OF LABOUR PAIN RELIEF METHODS AMONG WOMEN ATTENDING ANTENATAL CLINIC IN NAIROBI

V. Mung'ayi, MBChB, MMed (Anaesthesia), Fell. CCM. Senior Instructor, D. Nekyon, MBChB, MMed (Anaesthesia), Senior Instructor, Department of Anaesthesia and R. Karuga BSc, MPH, Research Support Unit, Aga Khan University Hospital, P. O. Box 30270-00100, Nairobi, Kenya

Request for reprints to: Dr. V. Mung'ayi, Department of Anaesthesia, Aga Khan University Hospital, P. O. Box 30270-00100, Nairobi, Kenya

KNOWLEDGE, ATTITUDE AND USE OF LABOUR PAIN RELIEF METHODS AMONG WOMEN ATTENDING ANTENATAL CLINIC IN NAIROBI

V. MUNG'AYI, D. NEKYON and R. KARUGA

ABSTRACT

Background: Labour analgesia has rapidly gained popularity in obstetric practice. Low usage in the developing world has raised concern.

Objectives: To assess the knowledge, attitude and use of labour pain relief methods in women attending antenatal clinic in Nairobi.

Design: A prospective study.

Setting: Aga Khan University Hospital, a teaching and referral hospital in Nairobi, Kenya.

Subjects: Two hundred and two consecutive expectant mothers attending antenatal clinic at the Aga Khan University Hospital, Nairobi.

Results: Fifty six per cent of the participants had knowledge about labour pain relief methods. Friends, the antenatal clinic and books/leaflets were the major source on information. Ninety per cent indicated they would intend to have some form of labour pain relief at their next delivery. Eighteen percent had been offered some form of pain relief at their last delivery with 82% of those offered having effective pain relief ($P < 0.001$).

Conclusions: While most of our participants were well educated, level of knowledge of labour analgesia is still low. Use of labour analgesia is also still quite low in comparison to the western World. It is recommended that obstetricians and anaesthetists participate in knowledge dissemination and setup of dedicated labour analgesia services in this region.

INTRODUCTION

Natural child birth is associated with labour pain that exceeds expectations. The scientific definition of pain was introduced by the International Association for the Study of Pain. Pain is an unpleasant sensory and emotional experience associated with actual or potential tissue damage, or described in terms of such damage (1).

Bonica (2) has defined acute pain as a complex constellation of unpleasant sensory, perceptual, and emotional experiences and certain associated autonomic, psychological, emotional, and behavioural responses.

Lowe (3) has described the nature of labour pain which begins with an understanding of the nociceptive stimuli that may be centrally perceived by the parturient and called pain. During the dilatation

phase of labour (first stage), it is the visceral pain that predominates with pain (nociceptive) stimuli arising from the mechanical distension of the lower uterine segment and cervical dilatation. High threshold myometrial mechanoreceptors may also generate nociceptive stimuli in response to uterine contractions especially in long, protracted labours. The increasing intensity of pain commonly observed with the progression of the dilatation may be partially attributable to a lowered activation threshold in the mechanoreceptors, and the chemoreceptor stimulation produced by repeated stimulation of uterine contractions. As the pelvic or descent phase of labour progresses (late first stage and second stage), somatic pain predominates from distension and traction on pelvic structures surrounding the vaginal vault and from distension of the pelvic floor and perineum.

The negative physiologic effects of labour pain are believed to originate mainly in alterations in the maternal respiratory pattern and the catecholamine mediated stress response.

Brownridge (4) has reviewed the potential physiological effects of severe labour pain which may include increased oxygen consumption and hyperventilation with hypocarbia and respiratory alkalosis, and autonomic stimulation and catecholamine release with gastric inhibition and increased gastric acidity, lipolysis, increased peripheral vascular resistance, cardiac output, and blood pressure, decreased placental perfusion and incoordinate uterine activity. At the extreme end of the spectrum, these responses are hypothesised to produce maternal metabolic acidaemia, fetal distress and dysfunctional labour. However Brownridge states that "such effects may be largely innocuous during the course of an uncomplicated labour".

Several surveys have been conducted that show a large proportion of women go through labour using various methods of pain relief (5 - 7).

In contrast to women in Western countries, the majority of women in the developing world including Africa have no access to pain relief methods in labour (8, 9). In order to assess the magnitude of the problem, we conducted a questionnaire survey of antenatal women regarding their knowledge, attitude and use of labour pain relief methods.

MATERIALS AND METHODS

Approval from the Aga Khan University Hospital Research and Ethics Committee for our survey was sought and obtained prior to commencement of the survey.

The survey was conducted in May, 2007. We surveyed 202 women attending antenatal clinic at the Aga Khan University Hospital, Nairobi over a period of 20 days using a questionnaire. The questionnaire captured their demographics, knowledge and attitude of labour pain relief methods.

An investigator was available to explain any technical terms in the questionnaire that the participants did not comprehend. Both primigravida and parous women were included in the survey. All descriptive and analytical statistics were performed using Intercooled Stata (StataCorp 200. Stata Statistical Software: Release 9. College Station, TX: Stata Corporation.). A p-value of <0.05 was considered significant for all statistical tests.

RESULTS

In May, 2007, 202 consecutive women attending the antenatal clinic at our hospital were enrolled

in our survey. All of them completed the provided questionnaires. Characteristics of the women interviewed are shown in Table 1.

The majority of the women were young (mean age 27 years), had a college level of education (52%) and were parous (65%). Fifty six per cent of the participants had knowledge of labour pain relief methods. Among participants who had knowledge of labour analgesia, friends, the antenatal clinic and books / leaflets were the major source on information.

Table 1

Baseline variables of interviewed women (n= 202)

Variable	No.	(%)
Mean age	27	-
Primary	5	2
Secondary	24	12
College	105	52
University	68	34
Primigravida	71	35
Multiparous	131	65
Knowledge of labour pain relief method	111	56
Doctor	14	13
Antenatal classes	28	25
Friends	26	23
Book / leaflets	35	32
Internet	8	7

There was no significant relationship between level of education and knowledge of pain relief methods, $P= 0.494$. Ninety four per cent of the parous women had experienced pain at their last delivery, with 54% of them having felt severe pain, 32% moderate pain, eight per-cent mild pain and six percent had not experienced any pain. A minority of the parous women (18%) had been offered any form of pain relief at their last delivery. Of those who had been offered pain relief, a majority of 82% had effective pain relief.

Table 2 shows the pain experience and management. There was a significant relation between pain relief administration and effectiveness, $P<0.001$. The majority of the participants in our survey, 90%, indicated they would intend to have some form of labour pain relief at their next delivery so long as they were assured it would be safe for them and their babies. Eight per cent declined and two per cent were unsure whether they would like pain relief at their next delivery.

Table 2
Pain experience and management (n=131)

Variable	No.	(%)
No pain	8	6
Mild pain	10	8
Moderate pain	42	32
Severe pain	71	54
Offered pain relief	22	18
Not offered pain relief	109	82
Offered pain relief (n= 22)		
Pain relief effective	18	82
Pain relief not effective	4	18

DISCUSSION

Our study shows that just over half of women attending antenatal clinic at the university hospital had knowledge about labour pain relief methods. This is comparable to findings of a similar study conducted in South Africa by Mugambe *et al* (10). Minhas *et al* (9) found higher figures of knowledge about labour pain relief methods at a university hospital in Pakistan. Other studies in Africa by Okeke *et al* (11) and Akpan *et al* (12) have shown much lower levels of knowledge of 38.9 and 4.5% respectively.

A striking finding in our setting was that while most of our survey participants were of college or university education, this was not associated with a higher level of knowledge on labour pain relief methods overall. This could be due to lack of emphasis on family based subjects in college and university curricula and shows the substantial unmet need for relevant health related information even among educated women.

Friends, booklets/leaflets and the antenatal clinic were the main sources of information in our survey. Stewart *et al* (13) found comparable results in their study. Although more than half of the women in our survey knew about labour analgesia, only a small percentage availed this service. This compares with a study by Onah *et al* (14) which showed that only 22.1% of parturients received some pain relieving drug during labour.

This low availability in our study could be due to extra expenses that may be involved in availing this service. Cultural and religious practices, lack of adequate information on safety and use of labour analgesia during antenatal clinics or via leaflets could also explain the low usage. There are no hospitals offering a well established labour analgesia service in our region. This could be another explanation for low availability of labour analgesia in our setup.

Although only a small percentage of our participants had received labour analgesia at their last delivery, it was reported as having been effective.

The level of effectiveness was comparable to findings elsewhere.

Findley *et al* (5) has shown that with pharmacological labour analgesia methods, entonox was helpful to about 85% of the women who used it, opiates provided 85% of women who used them with a good experience of pain relief while over 90% of the women who used epidural analgesia found it good or very good and 85% would choose to use it again. In our study the various pharmacological labour analgesia methods had a comparable level of effectiveness. A majority of participants in our survey expressed willingness to have labour analgesia at their next delivery if safety was assured for them and their babies.

What do the present findings add to knowledge about the relationship between social and educational context, knowledge, demand and use of pain relief methods in labour? Our patients might be considered to be socio-culturally located within a transition from traditional social norms for coping with labour pain, to a cultural milieu heavily influenced by Western values and expectations. Our findings suggest that the expected linkages between education in general, specific knowledge of pain relief methods, demand for and use of services are highly fragmented. However, it is clear that when offered relevant information on pain relief options our patients are certainly interested in these options. The implication for maternity care providers in urban settings of developing countries such as ours is that specific measures are needed to overcome the substantial information, access and service delivery gaps that currently exist.

Our survey has some limitations. The results of our study cannot be generalised as it was conducted in only one centre and most of the study participants were educated and employed. This would have had an impact on their level of knowledge, attitude and use of labour analgesia as compared to a similar survey conducted elsewhere. Also, the survey method was not able to provide the detailed insight into the meaning of labour pain for women that would be obtained through a qualitative study.

In conclusion, our study has shown that despite a big percentage of women attending our antenatal clinic being well educated, knowledge on labour analgesia methods is still below the Western World levels. Friends, books/leaflets and the antenatal classes are the most common source of information. We recommend reinforcement of these sources and urge the obstetricians and anaesthetists to become more actively involved in labour analgesia knowledge dissemination. A majority of the participants in our study have shown willingness to have a labour analgesia at their next delivery. Setting up dedicated labour analgesia services in this region would therefore go a long way in alleviating labour pain and its associated suffering.

REFERENCES

1. Merskey, H. Pain terms: a list with definitions and a note on usage. Recommended by the International Association for the Study of pain (IASP) subcommittee on taxonomy. *Pain*. 1979; **6**: 249-252
2. Bonica, J.J. Definitions and taxonomy of pain. In: Bonica J.J, editor. The management of pain. 2nd edition. Philadelphia: Lea & Febieger; 1990; p18-27.
3. Lowe, N.K. The nature of labour pain. *Am. J. Obstet. Gynecol.* 2002; **186**: S16- S24.
4. Brownridge, P. The nature and consequences of childbirth pain. *Eur. J. Obstet. Gynecol. and Reprod. Biol.* 1995; **59**: S9-S15.
5. Findley, I. and Chamberlain, G. ABC of labour care: Relief of pain. *Brit. Med. J.* 1999; **318**: 927-930.
6. Oystan, I. Obstetrical anaesthesia in Ontario. *Can. J. Anaesth.* 1995; **42**: 1117-1125.
7. Stamer, U.M., Messersschmidt, A., Wulf, H. and Hoeft, A. Practice of epidural analgesia for labour pain: a German survey. *Eur. J. Anaesthesiol.* 1999; **16**: 308-314.
8. Chan, Y.K. and Ng, K.P. Survey of regional analgesia and anaesthesia for obstetrics in selected countries in the Far East. *Int. J. Obstet. Anesth.* 2000; **9**: 225-232.
9. Minhas, M.R., Kamal, R., Afshan, G. and Raheel, H. Knowledge, attitude and practice of parturients regarding epidural analgesia for labour in a university hospital in Karachi. *J. Pak. Med. Assoc.* 2005; **55**: 63-66.
10. Mugambe, J.M., Nel, M., Hiemstra, L.A. and Steinburg, W.J. Knowledge of and attitude towards pain relief during labour of women attending Cecilia Makiwane Hospital's antenatal clinic, South Africa. *South Afr. Fam. Pract.* 2007; **49**: 16.
11. Okeke, C.I., Merah, N.A., Cole, S.U., *et al.* Knowledge and perception of obstetric analgesia among prospective parturients at the Lagos university Teaching Hospital. *Niger Postgrad. Med. J.* 2005; **12**: 258-261.
12. Akpan, S.G., Eshlet, A.I., Ilori, L.U., *et al.* Attitude of Nigerian mothers to labour pain and its relief. *Mary Slessor J. Med.* 2003; **3**: 12-13.
13. Stewart, A., Sodhi, V., Harper, N., *et al.* Assessment of the effect upon maternal knowledge of an information leaflet about pain relief in labour. *Anaesthesia.* 2003; **58**: 1015-1019.
14. Onah, H.E., Obi, S.N., Oguanuo, T.C., *et al.* Pain perception among parturients in Enugu, South Eastern Nigeria. *J. Obstet. Gynaecol.* 2007; **27**: 585-588.