



THE AGA KHAN UNIVERSITY

eCommons@AKU

Section of Dental-Oral Maxillofacial Surgery

Department of Surgery

8-2019

Evaluation of the root and canal morphology of mandibular first permanent molars in a sample of Pakistani population by cone-beam computed tomography

Muhammad Rizwan Nazeer
Bahria University Medical and Dental College

Farhan Raza Khan
Aga Khan University, farhan.raza@aku.edu

Follow this and additional works at: https://ecommons.aku.edu/pakistan_fhs_mc_surg_dent_oral_maxillofac

Recommended Citation

Nazeer, M. R., Khan, F. R. (2019). Evaluation of the root and canal morphology of mandibular first permanent molars in a sample of Pakistani population by cone-beam computed tomography. *Journal of Pakistan Medical Association*, 69(8), 1084-1089.

Available at: https://ecommons.aku.edu/pakistan_fhs_mc_surg_dent_oral_maxillofac/116

Evaluation of the root and canal morphology of mandibular first permanent molars in a sample of Pakistani population by cone-beam computed tomography

Muhammad Rizwan Nazeer,¹ Farhan Raza Khan²

Abstract

Objective: To evaluate the root canal morphology of permanent mandibular first molars using Cone Beam Computed Tomography.

Methods: The retrospective study was done at Aga Khan University Hospital, Karachi, and comprised medical records of patients who visited the dental clinics from December 2016 to March 2017. Mandibular first permanent molars were evaluated on Cone Beam Computed Tomography images. Frequency distribution of Vertucci's classification was determined, and so was the association between Vertucci's classifications and gender. SPSS 20 was used for data analysis.

Results: A total of 142 mandibular first permanent molars were evaluated on 78 Cone Beam Computed Tomography images. The most common Vertucci's classification was Type IV for mesial root 86(60.56%) and Type I for distal root 63(44%). There was no difference in the two genders for root canal morphology ($p>0.05$).

Conclusions: Type IV Vertucci's were prevalent in the mesial root and Type I were common in the distal root of permanent mandibular first molars.

Keywords: Root canal, Endodontics, CBCT, Vertucci's classification. (JPMA 69: 1084; 2019)

Introduction

The successful outcome of an endodontic therapy depends on certain factors in which the most important is root canal preparation.¹ One of the most common causes of root canal failure is inadequate debridement of root canal space, leaving pulp tissue remnants, microorganisms or their byproducts in the canal space. The remnants act as foci of infection, resulting in post-treatment disease.² For better treatment outcomes, it is essential for a clinician to have a detailed knowledge of the basic root and canal morphology or its morphological variants.^{3,4} There are many factors responsible for root canal variations such as ethnicity,⁵ gender⁶ or the assessment method⁷ used for its evaluation.

In the oral cavity, mandibular molars are the first permanent teeth to erupt and are the most common teeth that require endodontic treatment because of early caries.⁵ Variability in the root canal morphology exists among different races.⁶ Root canal morphology of mandibular first molar has been extensively studied among various population groups and their results demonstrate that it exhibits diverse morphologic variations.⁸⁻¹³ Gu et al.⁸ studied root canal anatomy in Chinese population and reported that the most common

root canal configuration in the root was Type IV (64.4%) followed by Type II (11.1%). However, Zaatari et al.⁹ reported in Kuwaiti population that the most common was Type II (56.5%) followed by Type IV (42.9%). Wasti et al.¹⁰ found that Type IV (66.7%) was the most frequent canal morphology found in the mesial root of Pakistani population, followed by Type II (23.3%), and so did Faraz et al.¹¹ who observed Type IV (70.7%) and Type II (26.8%), respectively.

Regarding distal root canal morphology, Rwenyonyi et al.¹² reported in Ugandan population the frequency of Type I (84.8%), Type IV (9.4%) and Type II (1.3%). However, Ahmed et al.,¹³ in Sudanese population reported the frequency of Type I (38%) followed by Type II (28%) and Type IV (22%). Wasti et al.¹⁰ reported in Pakistani population frequency of Type I (30%), Type II (26.7%) and Type IV (20%). Faraz et al.¹¹ reported higher frequency of Type I (65.8%), Type IV (19.5%) and Type II (14.6%). Hence, variations in canal morphology exist among various population groups.

Various methods have been reported in literature for the evaluation of root canal anatomy. These include canal staining and clearing technique,^{14,15} conventional and digital radiography,¹⁶ clinical operating microscopy¹⁷ and radiographic assessment enhanced with contrast medium.¹⁸ Conventional or digital periapical radiographs are most commonly used for evaluation of canal morphology before and during the treatment.

¹Operative Dentistry, Bahria University Medical and Dental College,

²Department of Surgery, Aga Khan University Hospital, Karachi.

Correspondence: Farhan Raza Khan. Email: farhan.raza@aku.edu

Nevertheless, due to inherent limitations, like overlapping of anatomical structures and distortions like elongation or shortening etc.,¹⁹ these radiographs have limited diagnostic value. Cone Beam Computed Tomography (CBCT) was introduced in dentistry a few years ago and is considered an effective diagnostic modality as it provides three-dimensional (3D) information and is accepted as the most reliable in vivo method for visualising canal morphology in endodontics.^{19,20} CBCT provides clinicians an opportunity to visualise root canal anatomy not only in cross-sectional, sagittal and coronal planes, but also in oblique planes. Provision of images in three different planes eliminates the overlap of anatomical structure, hence increasing the diagnostic accuracy. It helps clinicians in visualising the exact number of root canals and their convergence and divergence with precision.¹⁹⁻²² These advantages enable clinicians to perform effective endodontic treatment with reduced chance of failure.^{20,22}

Plenty of studies have been published on permanent mandibular first molar root canal anatomy among various population groups using CBCT.^{21,22} In local literature, so far only two studies have been published on root canal morphology of these teeth.^{10,11} However, none of the studies used CBCT as an assessment tool. The current study was planned to evaluate the root canal morphology of permanent mandibular first molars using CBCT.

Material and Methods

The retrospective study was conducted at the Aga Khan University Hospital (AKUH), Karachi, and comprised medical records of patients who visited the dental clinics between December 2016 and March 2017. After

exemption was obtained from the institutional ethics review board, permanent mandibular first molars were evaluated on CBCT scans Using non-probability consecutive sampling technique. World Health Organisation (WHO) sample size calculator²³ was used. With prior knowledge,²⁴ it was assumed that 38% of the distal roots of mandibular first molars have Vertucci's Class I canal morphology. Taking this population proportion with 8% absolute precision at 95% confidence level, the sample size needed for the study was calculated.

CBCT scans having permanent mandibular first molars both males and females of Pakistani origin aged 16-60 years and mature roots with no root canal fillings were included. Scans with periapical pathology, indirect restorations and blurred images were excluded.

CBCT scans were obtained using Sirona Dental system (D-64625 Bensheim, Germany). The voltage was kept at 85 KVp and current was 7mAs. Three-dimensional scans in each plane were studied using GALAXIS version 1.9 (SICAT GmbH & Co. KG, Bonn, Germany) on a 10-inch monitor.

SPSS 20 was used for data analysis. Vertucci's classification²⁵ was used for categorising different canal types. Chi-square test or Fisher's exact test, as needed, were used to determine any association between gender and canal type. P<0.05 was taken as statistically significant.

Results

A total of 142 mandibular first permanent molars were evaluated on 78 CBCT images. Of them, 80(56.3%) belonged to males and 62(43.7%) belonged to females. In

Table-1: Frequency distribution of Vertucci's classification and number of apical foramina in the mesial and distal roots of mandibular first molar.

Canal Morphology	Mesial root			p-value	Distal root			p-value
	Gender		Total n (%)		Gender		Total n (%)	
	Male n (%)	Female n (%)			Male n (%)	Female n (%)		
Type I (1)				0.091	36 (25.2)	27 (18.9)	63 (44.1)	0.095
Type II (2-1)	24 (16.9)	20 (14.1)	44 (31)		17 (11.9)	15 (10.5)	32 (22.4)	
Type III (1-2-1)					4 (2.8)	10 (7)	14 (9.8)	
Type IV (2)	50 (35.2)	36 (25.4)	86 (60.6)		11 (7.7)	7 (5)	18 (12.6)	
Type V (1-2)					8 (5.6)	2 (1.4)	10 (7)	
Type VI (2-1-2)		4 (2.8)	4 (2.8)		3 (2.1)		3 (2.1)	
Type VII (1-2-1-2)					2 (1.4)		2 (1.4)	
Others	6 (4.2)	2 (1.4)	8 (5.6)			1 (0.7)	1 (0.7)	
Total	80 (56.4)	62 (43.6)	142		81 (56.6)	62 (43.4)	143	
		Number of apical foramina				Number of apical foramina		
One			44 (31)				110(76.8)	
Two			96 (67.6)				33 (23.2)	
Three			2 (1.4)					
Total			142				143	

* Chi- square test (Fisher's exact test) was applied at 5% level of significance.

terms of Vertucci's classification, Type IV 86(60.56%) and Type II 44(30.99%) were most frequent in mesial root. The two genders did not differ significantly for mesial root anatomy ($p = 0.09$). Since one of the specimens had two distal roots, we had a total of 143 distal roots. The most frequent Vertucci's types in distal roots were Type I 63(44%) followed by Type II 32(22.3%). The anatomy of the distal root did not differ between the genders ($p = 0.09$).

Table-1 also shows frequency of the apical foramen in the mesial and distal roots of mandibular first molars. Two apical foramina were common in the mesial root 96(67.6%) followed by single apical foramen 44(31%) and three apical foramina 2(1.4%), respectively. In the distal root, the frequency of one and two apical foramina were 110(76.7%) and 33(23.2%) respectively (Table-1).

Discussion

The outcome of root canal treatment becomes predictable only if all the root canals are identified, adequately cleaned and filled-in 3D. Inadequate removal of pulpal tissue remnants, microorganism or their by-products can act as the foci of infection and cause eventual failure.² Hence, it is imperative that a clinician should be well aware of the canal morphology of a particular tooth, its variations along with typical features of different ethnicities, as this knowledge will aid in achieving favourable treatment outcomes. The chance of having a procedural error is higher in teeth

with atypical root canal morphology. These errors includes perforation, ledge formation, canal perforation and apical zipping etc.^{2,26}

A CBCT scan allows a clinician to visualise tooth anatomy in three anatomical planes i.e. sagittal, axial and coronal. All these planes can be viewed in thin slices, increasing its accuracy for the detection of small changes in the canal anatomy and the surrounding peri-apical tissues which were previously impossible.²⁷ In a CBCT software, several tools are available which makes it possible to provide images in an oblique plane, hence increasing its precision. Moreover, options such as contrast and brightness alteration are available that help to enhance the image quality. Similarly, tools are available that can help to decipher the nerves at the tooth apex with high precision. Therefore, CBCT can be considered one of the most valuable tools in dentistry.^{27,28}

Quite a few studies have reported the number of apical foramina in the mesial and distal roots of lower first molars (Table-2). In our sample, two apical foramina were common in the mesial root whereas distal root had predominately one apical foramina. These results were comparable to those reported by Chen et al.²⁹ (68.3%) in Chinese population. Our results were also comparable to those reported in Pakistani population by Wasti et al.¹⁰ (73.3%) and Faraz et al.¹¹ (73.1%). However, the high frequency of two apical foramina is reported by Wang et al.²¹ i.e. 95.8% and the lowest frequency is reported by

Table-2: Frequency of number of apical foramina (%) in mesial and distal roots of mandibular first molars as reported in the literature.

Author/year/ sample size	Method	Mesial root				Distal root			
		One (%)	Two (%)	Three (%)	Other (%)	One (%)	Two (%)	Three (%)	Other (%)
Skidmore & Bjorndahl /1971 /45 ³	In vitro (Plastic cast)	40	60			88.9	9.1		
Vertucci et al. /1984 / 100 ³⁹	Clearing method)	40	59	1		85	15		
Caliskanet al./ 1995 /100 ⁴⁰	Clearing method	39	57.6	3.4					
Zaatar et al. / 1997 / 49 ⁹	In vivo (PA)	57.1	42.9			83.7	16.3		
Al-Nazhan et al. /1999 / 251 ³⁴	In vivo (PA)	52.6	47.4			77.3	22.7		
Wasti et al.2001 /30 ¹⁰	Clearing method	23.3	73.3			56.7	43.3		
Gulavibala et al. /2002/ 118 ³²	Clearing method	35.6	60.2	3.4		77	17.8	5.2	
Chen et al. / 2009 / 183 ²⁹	Clearing method	31.7	68.3			67.2	32.8		
Rwonyonyi et al. /2009 / 224 ¹²	Clearing method	46.9	56.1			87.1	12.9		
Al-Qudah & Awawdeh/2009/ 330 ³³	Clearing method	39.1	58.5	2.4		77	22.1	0.9	
Wang et al./ 2010 / 410 ²¹	CBCT	4.2	95.8			98.6	1.4		
Chourasia et al. 2012 / 150 ³⁰	Clearing method	62.6	37.4			87.4	12.6		
Mukhaimer et al. /2014/320 ⁴¹	CBCT	42.5	57.5			90.6	9.4		
Faraz et al./ 2015/ 123 ¹¹	Clearing method	26.9	73.1			80.5	19.5		
Estrela et al./ 2015 / 100 ³¹	CBCT		36	49	15	5	54	40	1
Akhlaghi et al. / 2017/ 150 ³⁷	Clearing method	44.6	55.4			89.9	10.1		
Present study/ 2017/ 142	CBCT	67.3	31	1.4		76.8	23.2		

CBCT: Cone Beam Computed Tomography.

Table-3: Root canal configuration in mesial root of mandibular first molars according to various population groups.

Author/year/ sample size	Population	Method	Type II (%)	Type IV (%)	Type VIII (%)	Others (%)
Skidmore & Bjorndahl /1971 /45 ³	Caucasian	Clearing method	33.3	60		6.7
Vertucci et al. /1984 / 100 ³⁹	N/A	In vivo (PA)	28	43	1	28
Caliskan et al./ 1995 /100 ⁴⁰	Turkish	Clearing method	37.3	44.1	3.4	15.2
Zaatar et al. / 1997 / 49 ⁹	Kuwait	In vivo (PA))	56.5	42.9		0.9
Al-Nazhan et al. /1999 / 251 ³⁴	Saudian	In vivo (PA)	52.6	47.4		
Wasti et al. 2001 /30 ¹⁰	Pakistani	Clearing method	23.3	66.7		
Sperber and Moreau/ 1998 / 480 ³⁵	Senelgese	Invitro (sections)	16	84		
Gulavibala et al. /2001/ 139 ³⁶	Burmese	Clearing method	23.3	66.7	3.3	7.7
Gulavibala et al. /2002/ 118 ³²	Thai	Clearing method	28.8	38.1	0.7	32.4
Ahmed et al. / 2007/ 100 ¹³	Sudanese	Clearing method	14	73	2	11
Chen et al. / 2009 / 183 ²⁹	Taiwenese	Clearing method	29.5	55.2	5.5	9.8
Rwenyonyi et al. /2009 / 224 ¹²	Ugandian	Clearing method	13.8	44.6	0	41.6
Al-Qudah & Awawdeh/2009/ 330 ³³	Jordanian	Clearing method	36	52.7	0.3	11
Gu et al. / 2010 / 45 ⁸	Chinese	Invitro (mCT)	11.1	64.4	2.2	28.3
Wang et al. / 2010 / 410 ²¹	Chinese	CBCT	1.7	93.9	0.2	4.2
Chourasia et al. / 2012 / 150 ³⁰	Indian	Clearing method	36.6	54		9.4
Mukhaimer et al. /2014/320 ⁴¹	Palestinian	CBCT	38.8	53.8		7.4
Nuret al./ 2014/ 966 ³⁸	Turkish	CBCT	1	89	0.2	9.8
Faraz et al.2015/ 123 ¹¹	Pakistani	Clearing method	26.8	70.7		2.5
Akhlaghi et al. / 2017/ 150 ³⁷	Irani	Clearing method	41.3	55.3		
Present study/ 2017/ 142	CBCT	CBCT	60.56	30.99		8.45

CBCT: Cone Beam Computed Tomography

N/A: Not Applicable

PA: Periapical Radiograph

mCT: Micro Computed Tomography

Chourasia et al.³⁰ i.e. 37.4%. Estrela et al.³¹ reported that the occurrence of three apical foraminas (49%) was highest in the mesial root followed by two (36%) and four (15%), but none of their subjects had single apical foramina.

In our sample, the frequency of single apical foramen was highest in the distal root (76.76%) followed by two foramina (23.24%). Regarding the frequency of the single apical foramen, our results were in agreement with those reported by Gulavibala et al.³² (77%), Al-Qudah & Awwadeh³³ (77%) and Al Nazhan et al.³⁴ (77.3%). Our results were also comparable with those reported by Faraz et al.¹¹ (80.5%). However, Estrela et al.³¹ reported that the prevalence of two apical foramina (54%) was highest in the distal root followed by three (40%) and one (5%).

Literature has quite a few studies evaluating root canal configuration in mesial root of mandibular first molars according to various population groups (Table-3). The most common types reported for mesial roots are Type IV and Type II.^{3,10-12,29,30,32} In the current study, the highest frequency was of Type IV (60.5%) followed by Type II (30.9%) and Type V (2.8%). Our results were comparable to those reported by Skidmore and Bjorndahl³ (Type IV:

60%) and by Gu et al.⁸ (Type IVL 64.6%). These results were also comparable to those reported in Pakistani sub-population by Wasti et al.¹⁰ (Type IV: 66.7%). Exception to these findings was reported by Al-Nazhan et al.³⁴ and Zaatar et al.,⁹ with Type II being the most frequent finding (52.6% and 56.5%, respectively). The high frequencies of Type IV (93.9% and 84%) were reported by Wang et al.²¹ and Sperber and Moreau.³⁵ Moreover, we found canal types other than the Vertucci's eight types.²⁴ The frequency of such additional canal types reported in our sample was 5.63%. Out of 142, 5 mesial roots had 3-2 configuration which is Type XII, two roots had 2-3 configuration classified as Type XIII by Gulavibala et al.³⁶ and, 1 mesial root had 2-3-2 configuration categorised as Type XXI by Al-Qudah and Awwadeh.³³ It is often easy for a clinician to debride and obturate Type II and Type IV. However, presence of additional canal types requires substantial effort and manual dexterity. Failure to disinfect such a root canal system poses a challenge and may affect the outcome of the treatment.

Root canal configuration in distal root of mandibular first molars has also been studied frequently (Table-4). Of the 142 mandibular first molars in the current study, 1 tooth had two distal roots which means a total of 143 distal roots. The most common canal configuration reported for

Table-4: Root canal configuration in distal root of mandibular first molars according to various population groups.

Author/year/ sample size	Population	Method	Type I (%)	Type II (%)	Type IV (%)	Others (%)
Skidmore & Bjorndahl /1971 /45 ³	Caucasian	Clearing method	71.1	17.8	8.9	2.2
Vertucci et al. /1984 / 100 ³⁹	N/A	In vivo (PA)	70	15	5	10
Zaatar et al. / 1997 / 49 ⁹	Kuwait	In vivo (PA))	70.7	12.9	16.4	
Al-Nazhan et al. /1999 / 251 ³⁴	Saudian	In vivo (PA)	42.2	35.1	22.7	
Wasti et al. 2001 /30 ¹⁰	Pakistani	Clearing method	30	26.7	20	23.3
Gulavibala et al. /2001/ 139 ³⁶	Burmese	Clearing method	66.2	15.8	10.1	7.9
Gulavibala et al. /2002/ 118 ³²	Thai	Clearing method	61	4.2	15.3	19.5
Ahmed et al. / 2007/ 100 ¹³	Sudanese	Clearing method	38	28	22	12
Chen et al. / 2009 / 183 ²⁹	Taiweneese	Clearing method	54.1	12.6	25.1	8.2
Rwenyonyi et al. /2009 / 224 ¹²	Ugandian	Clearing method	84.8	1.3	9.4	4.5
Al-Qudah & Awawdeh/2009/ 330 ³³	Jordanian	Clearing method	54.2	17	9.4	19.4
Gu et al. / 2010 / 45 ⁸	Chinese	Invitro (mCT)	82.2	2.2	2.2	13.4
Wang et al. / 2010 / 410 ²¹	Chinese	CBCT	62.9	9.7	25.1	2.3
Chourasia et al. / 2012 / 150 ³⁰	Indian	Clearing method	65.3	20.6	9.3	4.8
Mukhaimer et al. /2014/320 ⁴¹	Palestinian	CBCT	57.5	22.5	8.1	11.9
Nur et al./ 2014/ 966 ³⁸	Turkish	CBCT	60	14	20	6
Faraz et al.2015/ 123 ¹¹	Pakistani	Clearing method	66	14.5	19.5	
Akhlaghi et al. / 2017/ 150 ³⁷	Irani	Clearing method	61.3	26.6	9.4	
Present study/ 2017/ 142	CBCT	CBCT	44.06	22.38	12.59	

CBCT: Cone Beam Computed Tomography.

distal root was Type I (44.0%) followed by Type II (22.3%), Type IV (12.5%), Type III (9.7%), Type V (6.9%), Type VI (2.1%) and Type VII (1.4%). One distal root had additional canal configuration i.e. 2-3-2, categorised as type XXI.³³ The predominance of Vertucci's Type I observed in our sample is similar to the findings reported by other studies ranging between 54% and 82%.^{3,8,11,12,21,29,30,32-34,36-38} Our frequency was similar to that reported by Al-Nazhan et al.³⁴ (Type I: 42.2%) and Ahmed et al.¹³ (Type I: 38%). In distal root, the frequency of Type I reported by Wasti et al.¹⁰ in sub-Pakistan population was 30%, which shows that prevalence of Type I is variable.

In the light of our findings, CBCT is a valuable tool for evaluating canal morphology. However, it does not replace the peri-apical radiograph for diagnosis and management in daily endodontic practice. The use of CBCT should be reserved in cases where its benefits outweigh the harm from additional radiation exposure. Scenarios such as abnormal or complex canal morphology warrants use of CBCT for precise determination of the pulp anatomy, canal orifices location and canal configuration. It can help in the detection of additional canal types, hence improving the outcome of endodontic treatment.

To our knowledge, the current study is the first on canal morphology done in a Pakistani sub-population using CBCT. It identified canal morphology of permanent

mandibular first molar and its variants which can be applied to endodontic practice to improve the quality of care and reduce the failures. The primary limitation of the present study is that the images were assessed by a single at a single centre. We recommend a larger sample size and in other populations which are poorly studied.

Conclusion

In a sample of Pakistani subjects visiting university dental clinics for the treatment of permanent mandibular first molars, Type IV Vertucci's was the most frequent canal configuration in the mesial root whereas Type I was the most prevalent in the distal root. There was no difference in terms of gender for root canal morphology.

Disclaimer: None.

Conflict of Interest: None.

Source of Funding: None.

References

1. Peters OA. Current challenges and concepts in the preparation of root canal systems: a review. *J Endod.* 2004; 30:559-67.
2. Lin LM, Skribner JE, Gaengler P. Factors associated with endodontic treatment failures. *J Endod.* 1992; 18: 625-7.
3. Skidmore AE, Bjorndal AM. Root canal morphology of the human mandibular first molar. *Oral Surg Oral Med Oral Pathol.* 1971; 32:778-84.
4. Reyhani MF, Rahimi S, Shahi S. Root canal therapy of a mandibular first molar with five root canals: a case report. *Iran Endod J.* 2007;

- 2:110-2.
5. De Pablo ÓV, Estevez R, Sánchez MP, Heilborn C, Cohenca N. Root anatomy and canal configuration of the permanent mandibular first molar: a systematic review. *J Endod.* 2010; 36:1919-31.
 6. Sert S, Bayirli GS. Evaluation of the root canal configurations of the mandibular and maxillary permanent teeth by gender in the Turkish population. *J Endod.* 2004; 30:391-8.
 7. Assadian H, Dabbaghi A, Gooran M, Eftekhari B, Sharifi S, Shams N, et al. Accuracy of CBCT, Digital radiography and cross-sectioning for the evaluation of mandibular Incisor root canals. *Iran Endod J.* 2016; 11:106-10.
 8. Gu Y, Lu Q, Wang H, Ding Y, Wang P, Ni L. Root canal morphology of permanent three-rooted mandibular first molars—part I: pulp floor and root canal system. *J Endod.* 2010; 36:990-4.
 9. Zaatar El, Al-Kandari AM, Alhomaidah S, Al-Yasin IM. Frequency of endodontic treatment in Kuwait: radiographic evaluation of 846 endodontically treated teeth. *J Endod.* 1997; 23:453-6.
 10. Wasti F. Root canal systems of the mandibular and maxillary first permanent molar teeth of South Asian Pakistanis. *Int Endod J.* 2001; 34:263-6.
 11. Faraz SA, Tariq A, Jameel A. Root canal morphology of mandibular first permanent molars-karachi sample. *Pak Oral Dent J.* 2015; 35:294-8.
 12. Rwenyonyi CM, Kutesa A, Muwazi LM, Buwembo W. Root and canal morphology of mandibular first and second permanent molar teeth in a Ugandan population. *Odontology.* 2009; 97:92-6.
 13. Ahmed HA, Abu-bakr NH, Yahia NA, Ibrahim YE. Root and canal morphology of permanent mandibular molars in a Sudanese population. *Int Endod J.* 2007; 40:766-71.
 14. Vertucci FJ. Root canal morphology and its relationship to endodontic procedures. *Endodontic Topics.* 2005; 10:3-29.
 15. Weng XL, Yu SB, Zhao SL, Wang HG, Mu T, Tang RY, et al. Root canal morphology of permanent maxillary teeth in the Han nationality in Chinese Guanzhong area: a new modified root canal staining technique. *J Endod.* 2009; 35:651-6.
 16. Weine FS, Hayami S, Hata G, Toda T. Canal configuration of the mesiobuccal root of the maxillary first molar of a Japanese sub-population. *Int Endod J.* 1999; 32:79-87.
 17. Baldassari-Cruz LA, Lilly JP, Rivera EM. The influence of dental operating microscopes in locating the mesiolingual canal orifices. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2002; 93:190-4.
 18. Naoum HJ, Love RM, Chandler NP, Herbison P. Effect of X-ray beam angulation and intraradicular contrast medium on radiographic interpretation of lower first molar root canal anatomy. *Int Endod J.* 2003; 36:12-9.
 19. Cotton TP, Geisler TM, Holden DT, Schwartz SA, Schindler WG. Endodontic applications of cone-beam volumetric tomography. *J Endod.* 2007; 33:1121-32.
 20. Patel S, Dawood A, Ford TP, Whaites E. The potential applications of cone beam computed tomography in the management of endodontic problems. *Int Endod J.* 2007; 40:818-30.
 21. Wang Y, Zheng QH, Zhou XD, Tang L, Wang Q, Zheng GN, et al. Evaluation of the root and canal morphology of mandibular first permanent molars in a western Chinese population by cone-beam computed tomography. *J Endod.* 2010; 36:1786-9.
 22. Silva EJNL, Nejaim Y, Silva AV, Haiter-Neto F, Cohenca N. Evaluation of root canal configuration of mandibular molars in a Brazilian population by using cone-beam computed tomography: an in vivo study. *J Endod.* 2013; 39:849-52.
 23. Lwanga SK, Lemeshow S. Sample size determination in health studies: A practical manual. World Health Organization. Geneva. 1991:1-3.
 24. Razmi H, Shokouhinejad N, Hooshyar M. An In vitro study of the number of distal roots and canals in mandibular first molars in Iranian population. *Iran Endod J.* 2008; 2:126-30.
 25. Vertucci F, Seelig A, Gillis R. Root canal morphology of the human maxillary second premolar. *Oral Surg Oral Med Oral Pathol.* 1974; 38:456-64.
 26. Gutmann JL, Fan B. Tooth morphology, isolation, and access. In: Cohen S, Hargraves KM, Eds. *Pathways of the Pulp.* 11th ed. Mosby Elsevier, St Louis; 2016:193-9.
 27. Baratto-Filho F, Zaitter S, Haragushiku GA, De Campos EA, Abuabara A, Correr GM. Analysis of the internal anatomy of maxillary first molars by using different methods. *J Endod.* 2009; 35:337-42.
 28. D'addazio P, Carvalho A, Campos C, Devito K, Özcan M. Cone beam computed tomography in endodontics. *Int Endod J.* 2016; 49:311-2.
 29. Chen G, Yao H, Tong C. Investigation of the root canal configuration of mandibular first molars in a Taiwan Chinese population. *Int Endod J.* 2009; 42:1044-9.
 30. Chourasia HR, Meshram GK, Warhadpande M, Dakshindas D. Root canal morphology of mandibular first permanent molars in an Indian population. *Int J Dent.* 2012; 745152: 1-6
 31. Estrela C, Bueno MR, Couto GS, Rabelo LEG, Alencar AHG, Silva RG, et al. Study of root canal anatomy in human permanent teeth in a subpopulation of Brazil's center region using Cone beam computed tomography-Part 1. *Braz Dent J.* 2015; 26:530-6.
 32. Gulabivala K, Opasanon A, Ng YL, Alavi A. Root and canal morphology of Thai mandibular molars. *IntEndod J.* 2002; 35:56-62.
 33. Al-Qudah AA, Awawdeh LA. Root and canal morphology of mandibular first and second molar teeth in a Jordanian population. *Int Endod J.* 2009; 42:775-84.
 34. Al Nazhan S. Incidence of four canals in root-canal-treated mandibular first molars in a Saudi Arabian sub-population. *Int Endod J.* 1999; 32:49-52.
 35. Sperber GH, Moreau JL. Study of the number of roots and canals in Senegalese first permanent mandibular molars. *Int Endod J.* 1998; 31:117-22.
 36. Gulabivala K, Aung TH, Alavi A, Ng YL. Root and canal morphology of Burmese mandibular molars. *Int Endod J.* 2001; 34:359-70.
 37. Akhlaghi NM, Khalilak Z, Vatanpour M, Mohammadi S, Pirmoradi S, Fazlyab M, et al. Root canal anatomy and morphology of mandibular first molars in a selected Iranian population: an in-vitro study. *Iran Endod J.* 2017; 12:87 -91.
 38. Nur BG, Ok E, Altunsoy M, Aglarci OS, Colak M, Gungor E. Evaluation of the root and canal morphology of mandibular permanent molars in a south-eastern Turkish population using cone-beam computed tomography. *Eur J Dent.* 2014; 8:154-9.
 39. Vertucci FJ. Root canal anatomy of the human permanent teeth. *Oral Surg Oral Med Oral Pathol.* 1984; 58:589-99.
 40. Caliskan MK, Pehlivan Y, Sepetcioglu F, Turkun M, Tuncer SS. Root canal morphology of human permanent teeth in a Turkish population. *J Endod.* 1995; 21:200-4.
 41. Mukhaimer RH. Evaluation of Root canal configuration of mandibular first molars in a Palestinian population by using Cone-beam computed tomography: an ex vivo study. *Int Sch Res Notices.* 2014; 2014:583621.