



THE AGA KHAN UNIVERSITY

eCommons@AKU

Section of Orthopaedic Surgery

Department of Surgery

2-2019

Inflammatory arthritis and mycobacterium tuberculosis infection: A diagnostic and management challenge for knee arthroplasty in endemic areas

Obada Hasan

Aga Khan University, obada.husseinali@aku.edu

Riaz Hussain Lakhdawala

Aga Khan University, riaz.lakhdawala@aku.edu

Vickash Kumar

South City Hospital Karachi

Syed Faisal Mahmood

Aga Khan University, faisal.mahmood@aku.edu

Shahryar Noordin

Aga Khan University, shahryar.noordin@aku.edu

Follow this and additional works at: https://ecommons.aku.edu/pakistan_fhs_mc_surg_orthop



Part of the [Infectious Disease Commons](#), [Orthopedics Commons](#), and the [Surgery Commons](#)

Recommended Citation

Hasan, O., Lakhdawala, R. H., Kumar, V., Mahmood, S. F., Noordin, S. (2019). Inflammatory arthritis and mycobacterium tuberculosis infection: A diagnostic and management challenge for knee arthroplasty in endemic areas. *Journal of the Pakistan Medical Association*, 69(1 (Supplement-1)), S-101-S-107.

Available at: https://ecommons.aku.edu/pakistan_fhs_mc_surg_orthop/118



THE AGA KHAN UNIVERSITY

eCommons@AKU

Department of Surgery

Department of Surgery

February 2019

Inflammatory arthritis and Mycobacterium Tuberculosis infection: a diagnostic and management challenge for Knee arthroplasty in endemic areas

Obada Hussein Hasan

Aga Khan University, obada.husseinali@aku.edu

Riaz Hussain Lakdawala

Aga Khan University, riaz.lakdawala@aku.edu

Vickash Kumar

Syed Faisal Mahmood

Aga Khan University, faisal.mahmood@aku.edu

Shahryar Noordin

Aga Khan University, shahryar.noordin@aku.edu

Follow this and additional works at: https://ecommons.aku.edu/pakistan_fhs_mc_surg_surg

Recommended Citation

Hasan, O. H., Lakdawala, R., Kumar, V., Mahmood, S., Noordin, S. (2019). Inflammatory arthritis and Mycobacterium Tuberculosis infection: a diagnostic and management challenge for Knee arthroplasty in endemic areas. *The Journal of the Pakistan Medical Association*, 69(Supl. 1), 101-107.

Available at: https://ecommons.aku.edu/pakistan_fhs_mc_surg_surg/765

Inflammatory arthritis and Mycobacterium Tuberculosis infection: a diagnostic and management challenge for Knee arthroplasty in endemic areas

Obada Hasan,¹ Riaz Lakdawala,² Vickash Kumar,³ Faisal Mahmood,⁴ Shahryar Noordin⁵

Abstract

Tuberculosis continues to be one of the most challenging health problems more prevalent in developing countries. Pakistan ranks 5th in tuberculosis prevalence among the high-burden countries. Prosthetic joint infection of the knee by acid fast bacilli is a rare and distressing complication, occurring in nearly 1% of primary joint arthroplasties requiring prolonged medical treatment and multiple surgeries. A recent publication extensively reviewed English literature from 1952 to 2016, and reported only 64 prosthetic joint infection with tuberculosis, of which 27 cases involved the knee. Tuberculosis is a global health problem adding to the challenges that arthroplasty surgeons face in our resource-constrained setting. Furthermore, it presents as other inflammatory arthritis with almost same laboratory and radiological findings. The current paper was planned to highlight the preoperative and postoperative challenges that the arthroplasty surgeon may have in diagnosis and management of this rare infection. We included studies from 1996 to date which reported knee tuberculosis prosthetic joint infection that were managed by medication alone or with surgical intervention in patients who had undergone arthroplasty.

Keywords: Knee arthroplasty, TB prosthetic joint infection, Endemic, Inflammatory arthritis.

Introduction

Tuberculosis (TB) continues to be one of the most challenging health problems and is more prevalent in developing countries. Pakistan ranks fifth in TB prevalence amongst the high-burden countries.¹ In this day and age joint replacement surgeries have become popular in developing countries where TB also has high prevalence. Prosthetic joint infection (PJI) of the knee by acid fast bacilli (AFB) is a rare and distressing complication occurring in nearly 1% of primary joint arthroplasties^{2,3}

requiring prolonged medical treatment and multiple surgeries. Patients can present with a palpable mass (cold abscess), draining sinus/fistula, and painful erythema. Therefore high clinical suspicion is mandatory for diagnosis of PJI by mycobacterium. Adding to this challenge is inflammatory arthritis affecting the knee joint present with almost identical features on clinical examination, laboratory tests and radiographic findings. Laboratory tests include tuberculin skin test (TST), erythrocytes sedimentation rate (ESR) and cultures of synovial fluid, cytology and histology with different rates of accuracy. In endemic areas with mycobacterium TB (MTB), the low threshold of suspicious is indeed helpful for early diagnosis and management of this catastrophic infection.

Factors contributing to delayed diagnosis include its low incidence, varied clinical manifestations, co-infection with pyogenic bacteria accounting for 37% cases, and low index of suspicion.⁴

First case of peri-PJI with MTB was reported in 1977(5). A recently 2018 publication extensively reviewed English-language literature from 1952 to 2016, and reported only 64 PJI with TB, of which 27 cases involved the knee.⁶

Discussion

Pathology: Three pathogenic mechanisms have been reported: active TB arthritis present at the time of surgery but not known to clinician; TB spread by haematogenous route from foci elsewhere; and surgical trauma to old granulomas resulting in recurrence of TB arthritis.⁵⁻⁸ Recent studies have highlighted human immunodeficiency virus (HIV) as an important risk factor for re-activation of TB in previously infected joints.^{9,10} Estimated risk of re-activation has been reported between 0% and 31%, with total knee arthroplasty (TKA) more at risk than total hip arthroplasty (THA) (27% and 6%).¹¹ Staphylococcus is the most common organism for prosthetic infections followed by gram-negative and streptococcus species, while atypical infections are rare.^{18,12}

^{1,2,5}Department of Surgery, Section of Orthopaedics, The Aga Khan University, Karachi Pakistan. ³South City Hospital Karachi. ⁴Infectious Disease, Department of medicine, The Aga Khan University, Karachi Pakistan

Correspondence: Obada Hasan. e-mail: Obada.husseinali@aku.edu

MTB has limited biofilm-formation capacity which, if formed, is very thin and has a lesser tendency to adhere to implants compared to the biofilm-forming staphylococci which have ample biofilm-forming capacity and stronger ability to multiply and adhere on surface of all types of implants.¹³⁻¹⁵ Because of these factors, MTB is more susceptible to anti-TB agents and, if implant is stable, removal of hardware is not needed.^{13,16} Nevertheless, emergence of multidrug-resistant (MDR) and extensively drug-resistant (XDR) TB has complicated its management.¹⁶

Clinical presentation: PJI of knee with MTB has a varied clinical presentation. Constitutional symptoms, including fever, weight loss and night sweats, are not seen in all cases and a previous history of pulmonary Koch's is often absent.^{17,18} Patients can present with a palpable mass (cold abscess), draining sinus/fistula, and painful erythema. Therefore high clinical suspicion is mandatory for diagnosis of PJI by mycobacterium. Onset of symptoms within 2 month after arthroplasty is considered "early presentation" whereas onset of symptoms after 2 months is labelled as "delayed presentation".^{7,19}

Role of lab investigation: The ESR level may be elevated, but has a low specificity. ESR levels remain elevated for about 90 days or more than a year after arthroplasty, making it a less valuable test at least during this timeframe.^{14,20} TST is sensitive, but has a reduced positive predictive value (PPV) in populations with a low infection rate. It may report false negative results in patients who are immune-compromised, malnourished or HIV-positive.³ It is also not able to differentiate between active and latent TB. Clinical utility of different tests in diagnosing extra-pulmonary TB also need to be considered (Table 1).

Role of radiological investigation: Plain radiographs either of knee or chest are not specific. Radiological findings may show subchondral cysts, subchondral erosions or implant loosening, but these findings may also be present with pyogenic infection, aseptic loosening and will be absent in early presentation where the implant is stable.^{21,22}

Magnetic resonance imaging (MRI) has low sensitivity due to artefacts caused by the implant. MRI findings

Table-1: Clinical utility of different tests in diagnosing extra-pulmonary TB(27, 28).

	Tuberculin Skin Test (TST)	Smear for AFB	Xpert(39)	Histology
Sensitivity, (95% CI)	94% (87-98)	25% (12-42)	70% (50-85)	100% (85.7-100)
Specificity, (95% CI)	88% (74-96)	100% (75-100)	100% (75-100)	100% (95.5-100)
PPV, (95% CI)	95% (88-98)	100% (66-100)	100% (86-100)	100% (85.7-100)
NPV, (95% CI)	86% (72-94)	32.5% (19-49)	72.8% (31-97)	100% (95.5-100)

reveal large intra-articular effusions, peri-articular osteoporosis, and gross thickening of the remaining synovial membrane which are not clinically relevant in isolation after arthroplasty.⁸

Nuclear medicine is the most valuable diagnostic procedure to detect the prosthesis involvement and loosening,²³ but literature review has reported their use to be limited. Bone scans findings with technecium-99m or indium-111 are non-specific because findings mimic pyogenic infection, metastatic disease and non-specific inflammation.⁵

Grams staining / histological diagnosis: Aspiration of synovial fluid is another modality, but yield is low, with reported sensitivity of 80%. Literature reports synovial biopsy as gold standard in diagnosing PJI of knees.²⁴⁻²⁶ It has sensitivity of about 90%, but should always be added with cultures to get information about antibiotic resistance.^{27,28}

Culture specimens taken from draining sinuses are usually contaminated.^{3,6} Staining and visualisation of mycobacterium with Ziehl-nelson is time-saving and cost-effective, but yield of positive test is less than culture.¹⁷

Polymerase chain reaction (PCR), though one of the recent diagnostic modalities, has significant specificity, but is less sensitive at 60%. There are limited studies mentioning its use in the diagnosis of PJI.²⁹⁻³²

Treatment: Since PJI of knee with mycobacterium is rare, with varied presentation and delayed diagnoses; there are no specific guidelines for its management. Several different treatment plans have been advised for PJI of knee by AFB. Management varies from case to case. Literature reports treatment options that range from chemotherapy alone to arthrodesis or revision in addition to chemotherapy.

Conservative management: Early onset PJI of knee has been managed by retention of the implant with prolonged chemotherapy. There is great controversy regarding treatment duration of chemotherapy and combination of drugs in the literature. Cases of PJI with

Table-2: The cases of tuberculosis-prosthetic joint infection (TB-PJI) managed with medications alone.

Author/year (ref.)	Age/sex	Time elapsed from arthroplasty to joint infection	Time elapsed from joint infection to diagnosis	Concomitant infections	Medical therapy (duration in months)*	Follow-up from end of therapy (months)	Complications
Marschall et al. 2008 ⁴⁰	48/M	6 months	3 months	None	I, P, E, M ⁶ → I, P, E, R ⁶	Not reported	Death due to disseminated TB
Neogi et al. 2009 ¹⁸	73/F	14 years	2 months	None	I, P, E, R ⁴ → I, P, R ³ → I, R ¹¹	36	Not reported
Kadokia AP et al. 2007 ¹⁶	85/F	1 month	3 months	Staphylococcus	I, P, E, R ⁶	Not reported	Not reported
Veloci1et al(2018) ⁶	34/F	8 months	4 years	None	I, R ¹⁸ , P, E ²	24	None
Egües Dubuc C et al. 2014 ⁴¹	77/F	Not reported	1 year	None	I, P, R (on therapy)	48	None

I: Isoniazid P: Pyrazinamide R: Rifampicin E: Ethambutol M: Moxifloxacin

MTB managed with medications alone are worth taking a look at (Table 2).

Surgical management: Surgical treatment options include debridement alone, single-stage or two-stage implant exchange or removal of prosthesis and arthrodesis. Chemotherapy alone or with surgical debridement has been used in early onset PJI. In late onset PJI, implant usually gets loose, and removal of implant is often required.³³ Wolfgang shared his experience of late onset PJI in knee, managed with removal of implant, extensive debridement and two-stage revisions with adjuvant chemotherapy and good results at 1-year follow-up.³² A successful case of staged procedure in TB arthritis is also known. Figure 1 and 2 summarize a case of 55 years old lady presented with right knee pain, stiffness and difficulty walking not responding to conservative measures. Right knee was swollen, warm and tender with moderate effusion. Active range of motion 0-110 degrees. No varus/valgus or AP instability was seen on clinical examination. Blood tests were normal except for ESR and CRP, 75mm/hr and 2.14mg/dL, respectively. These features suggested erosive arthritis, which could be inflammatory. However TB should be considered high up in the differentials particularly in endemic areas with MTB. Patient



Figure-1: Preoperative knee x-ray showing juxta-articular osteopenia, peripheral osseous erosions and narrowing of joint space.

underwent debridement and open biopsy at first stage, which confirmed the diagnosis of TB.

Multidisciplinary approach including the surgeon, infectious disease team and microbiologist was followed. Following ATT for 10 months, patient had a successful TKA with an excellent outcome.

This strengthens the concept that not every erosive arthritis is a systemic inflammatory rheumatoid type, particularly so in our endemic areas of TB. Patient had no varus/valgus or AnteroPosterior (AP) instability on clinical examination. Blood tests were normal except for raised ESR and CRP. Pre-operative knee X-ray showed juxta-articular osteopenia, peripheral osseous erosions and narrowing of joint space. These features suggested erosive arthritis, which could be inflammatory. However, TB should be considered high up in the differentials, particularly in endemic areas with MTB. Patient underwent debridement and open biopsy at first stage, which confirmed the diagnosis of TB.

In endemic areas, one, therefore, has to make sure that TB is considered in the differential diagnosis of inflammatory arthritis as the radiological features alone



Figure-2: Postoperative x-rays show proper sagittal and coronal alignment. She is pain free at her 2nd year follow-up and is walking without any support.

Table-3: The cases of tuberculosis-prosthetic joint infection (TB-PJI) managed surgically.

Author/year (ref.)	Age/sex	Time elapsed from arthroplasty to joint infection	Time elapsed from joint infection to diagnosis	Concomitant infections	Medical therapy (duration in months)*	Surgical procedure	Surgery Time elapsed from start of medical therapy to surgery (months)	Follow-up from end of therapy (months)	Complications
Veloci et al. 2008 ⁶	62/M	Postoperatively	3 years	None	I, R (18), P ²	Debridement and implant retained	Not reported	1 month	None
Su et al. 1996 ⁴²	16 knees, in 2 groups (A & B)	A: 8 knees before TKA B: 8 knees diagnosed within 12 months post TKA	Variable	None	A: 20m before TKA and 12m post TKA B: 12m post TKA only	Only Anti Tuberculosis Therapy (ATT) for 1 case Debridement + ATT for 3 cases Excisional arthroplasty for 1 case of recurrent infection in group A	Within 12m	36	5 cases of reactivation (1 in group A and 4 in group B)
Tokumoto et al. 1995 ³¹	71/F	2 months	Not reported	S. aureus	I, E ¹⁸	Debridement, explant and later arthrodesis	9	96	None
Tokumoto et al. 1995 ³¹	70/F	38 years	3 months	None	I, R ¹²	Debridement and explant	Surgery first	24	Not reported
Lusk et al. 1995 ⁴³	75/F	16 years	Not reported	Not reported	I, R, P (0.7) → I, P, E (5.3)	Debridement and removal of arthroplasty	Surgery first	6	Death (other reasons)
Spinner et al. 1996 ⁴⁴	70/F	5 years	2 months	Coagulase negative Staphylococcus	R, E ¹²	Debridement	Surgery first	30	None
Al-Shaikh & Goodman 2003 ⁴⁵	73/F	8 months	5 months	S. aureus	I, r, P, E (2.1) → I, R, P ¹²	Debridement and later arthrodesis	5 months	12	None
Marmor et al. 2004 ¹⁴	66/M	2 months	4 months	None	I, R, P ¹⁶	Debridement and staged exchange	(procedures 14 months apart)	60	None
Marmor et al. 2004 ¹⁴	65/F	3 months NR	6 months	None	I, E, P ⁸	Debridement and staged exchange	Removal first, reimplantation 14 months	84	None
Wang et al. 2007 ⁴⁶	72/M	1 year	3 months	None	I, R, P, E (0.7)	Debridement	Surgery first	1	Death due to MODS
Khater et al. 2007 ⁸	75/F	3 months	1 month	S. aureus	I, R, P, E (unclear duration) → I, E ¹⁸	Debridement and explant	NR	18	None
de Haan et al. 2008 ²⁰	75/F	3 months	1 month	None	I, R, P, E ⁹	Debridement	9	36	None
Lee et al. 2009 ⁴⁷	79/F	2 months	2 months	Not reported	R, E ¹²	Debridement	Surgery first	13	None
Uppal S et al. 2010 ³⁶	27/F	1 month	3 years	Not reported	I, R, P, E ¹⁸	Debridement	9	Not Reported	None
Kadakia AP et al. 2007 ¹⁶	85/F	1 month	3 months	Staphylococcus	I, R, P, E ⁶	None	Variable	Not reported	Not reported
Harwin SF et al. [11], 2013 ⁴⁸	60/F	7 months	2 years	None	I, R ²¹ , P, E ¹²	Staged exchange	13	24	None
Klein GR et al. 2012 ³⁸	36/F	11 months	1 year	None	I, R, E, P, M, A ¹⁹	Staged exchange 36 months	Removal first, reimplantation	7	None
Carrega G et al. 2012 ³⁷	72/F	7 years	Not reported	None	I, R ¹² , E ¹²	Staged exchange	Surgery first	12	None
Carrega G et al. [19], 2012 ³⁷	80/F	Not reported	Not reported	None	I, R ¹⁴ , E ¹²	Staged exchange	Surgery first	7	Not reported

I: Isoniazid P: Pyrazinamide R: Rifampicin E: Ethambutol M: Moxifloxacin A: Amikacin




Figure-3: Right knee swelling and healed healthy scar with discharging sinus and non-healing wound over the lateral aspect of popliteal fossa (arrow).




Figure-4: X-rays showing the osteolysis around the unstable implant with significant bone loss and erosions with knee effusion.

are not enough to differentiate between these conditions. Indeed, these two different pathologies of arthritis present with same laboratory and radiological findings include elevated ESR and CRP, as well as erosive arthritis without formation of osteophytes and with mono-articular involvement.

Cases of TB-PJI managed surgically by debridement and retention of implant, debridement and explant, staged procedure (debridement followed by TKA) or arthrodesis have been known (Table 3).³⁴

Surgical treatment depends on the status of implant fixation. Implant may be retained if it is stable and only debridement followed by chemotherapy may be required. This strategy has been reported successfully in multiple cases and studies.^{19,29,35,36}

In case of implant loosening or co-infection with pyogenic organism, removal of loose implant and staged revision have been reported.^{37,38} In recurrent infection cases in severely sick patients, arthrodesis, if bone stock is available, or above-knee amputation, in cases with

significant bone loss and destruction, can be potential options to improve patients' quality of life. Figure 3 and 4 showing a 68 year old lady with multiple comorbid. Right TKA done for reported advanced OA outside our institution. Presented with pain and swelling of right knee within 11 months postoperatively. Patient underwent debridement and implant removal. Peroperative cultures grew MTB. Planned for revision TKA once free from disease or arthrodesis. Patient was kept on ATT but was not compliant and had multiple hospital admissions due to heart failure, asthma, electrolyte imbalance and drowsiness. This unfortunate sick lady had recurrent TB infection with significant bone loss and ultimately underwent above knee amputation.

Conclusion

Obtaining synovial specimens and specifically requesting TB culture and histology are the most pertinent investigations. Early diagnosis and treatment may prevent prosthetic loosening and avoid revision surgery with significant benefit to the patient and optimising outcomes and resources. Adding to this challenge is inflammatory arthritis affecting the knee joint present with almost identical features on clinical examination, laboratory tests and radiographical findings. TB should always be amongst differential diagnosis in cases of erosive inflammatory arthritis in endemic areas. Needless to say that this issue should be dealt with in multi-disciplinary setting including experienced surgeon, infection disease control consultant, histopathologist and public health workers to improve the outcome of these patients in a developing country.

Disclaimer: We declare that this paper was not previously presented or published anywhere. This was not part of any other projects.

Conflict of Interest: No conflicts of interest to disclose.

Sources of Funding: No funding from any source for this paper.

References

1. World Health Organization. Global tuberculosis control: epidemiology, planning, financing; WHO report 2009. [Internet] Geneva: World Health Organization; 2009 [cited 2018 December 30] Available from: <http://www.who.int/iris/handle/10665/44035>
2. Barnes S, Salemi C, Fithian D, Akiyama L, Barron D, Eck E, et al. An enhanced benchmark for prosthetic joint replacement infection rates. *Am J Infect Control* 2006;34:669-72.
3. Miandad M, Burke F, Nawaz-ul-Huda S, Azam M. Tuberculosis incidence in Karachi: a spatio-temporal analysis. *Geografia Malays J Soc Space* 2017;10:1-8.

4. Salemi C, Anderson D, Flores D. American Society of Anesthesiology scoring discrepancies affecting the National Nosocomial Infection Surveillance System: surgical-site-infection risk index rates. *Infect Control Hosp Epidemiol* 1997;18:246-7.
5. McCullough C. Tuberculosis as a late complication of total hip replacement. *Acta Orthop Scand* 1977;48:508-10.
6. Veloci S, Mencarini J, Lagi F, Beltrami G, Campanacci DA, Bartoloni A, et al. Tubercular prosthetic joint infection: two case reports and literature review. *Infection* 2018;46:55-68.
7. Horsburgh Jr CR, Rubin EJ. Latent tuberculosis infection in the United States. *N Engl J Med* 2011;364:1441-8.
8. Khater FJ, Samnani IQ, Mehta JB, Moorman JP, Myers JW. Prosthetic joint infection by *Mycobacterium tuberculosis*: an unusual case report with literature review. *South Med J* 2007;100:66-9.
9. Berbari EF, Hanssen AD, Duffy MC, Steckelberg JM, Osmon DR. Prosthetic joint infection due to *Mycobacterium tuberculosis*: a case series and review of the literature. *Am J Orthop (Belle Mead NJ)* 1998;27:219-27.
10. Johnson R, Barnes K, Owen R. Reactivation of tuberculosis after total hip replacement. *J Bone Joint Surg Br* 1979;61-B:148-50.
11. Burger JD, de Jongh H. Total knee replacement infected with *Mycobacterium tuberculosis*: a case study and review of the literature. *SA Orthop J* 2013;12:64-8.
12. Poultsides LA, Karydakos G, Karachalios T, Kaitelidou D, Papakonstantinou V, Liarpoulos L, et al. The impact of infection after total knee arthroplasty on hospital and surgeon resource utilization. A micro costing analysis. *Orthop Proc* 2009;91-B(Suppl II):314.
13. Ha KY, Chung YG, Ryoo SJ. Adherence and biofilm formation of *Staphylococcus epidermidis* and *Mycobacterium tuberculosis* on various spinal implants. *Spine (Phila Pa 1976)* 2005;30:38-43.
14. Marmor M, Parnes N, Dekel S. Tuberculosis infection complicating total knee arthroplasty: report of 3 cases and review of the literature. *J Arthroplasty* 2004;19:397-400.
15. Koruk ST, Sipahio?lu S, Cali?ir C. Periprosthetic tuberculosis of the knee joint treated with antituberculous drugs: a case report. *Acta Orthop Traumatol Turc* 2013;47:440-3.
16. Kadakia AP, Williams R, Langkamer VG. Tuberculous infection in a total knee replacement performed for medial tibial plateau fracture: a case report. *Acta Orthop Belg* 2007;73:661-4.
17. Barr DA, Whittington AM, White B, Patterson B, Davidson RN. Extrapulmonary tuberculosis developing at sites of previous trauma. *J Infect* 2013;66:313-9.
18. Neogi DS, Kumar A, Yadav CS, Singh S. Delayed periprosthetic tuberculosis after total knee replacement: is conservative treatment possible? *Acta Orthop Belg* 2009;75:136-40.
19. Griffith DE, Aksamit T, Brown-Elliott BA, Catanzaro A, Daley C, Gordin F, et al. An official ATS/IDSA statement: diagnosis, treatment, and prevention of nontuberculous mycobacterial diseases. *Am J Respir Crit Care Med* 2007;175:367-416.
20. de Haan J, Vreeling A, van Hellemond GG. Reactivation of ancient joint tuberculosis of the knee following total knee arthroplasty after 61 years: a case report. *Knee* 2008;15:336-8.
21. Sundfeldt M, Carlsson LV, Johansson CB, Thomsen P, Gretzer C. Aseptic loosening, not only a question of wear: a review of different theories. *Acta Orthop* 2006;77:177-97.
22. Trampuz A, Zimmerli W. Prosthetic joint infections: update in diagnosis and treatment. *Swiss Med Wkly* 2005;135:243-51.
23. Krappel FA, Harland U. Failure of osteosynthesis and prosthetic joint infection due to *Mycobacterium tuberculosis* following a subtrochanteric fracture: a case report and review of the literature. *Arch Orthop Trauma Surg* 2000;120:470-2.
24. Ahmad SS, Shaker A, Saffarini M, Chen AF, Hirschmann MT, Kohl S. Accuracy of diagnostic tests for prosthetic joint infection: a systematic review. *Knee Surg Sports Traumatol Arthrosc* 2016;24:3064-74.
25. Della Valle C, Parvizi J, Bauer TW, Dicesare PE, Evans RP, Segreti J, et al. Diagnosis of periprosthetic joint infections of the hip and knee. *J Am Acad Orthop Surg* 2010;18:760-70.
26. Suren C, Harrasser N, Pohlif F, Banke IJ, Lenze U, Lenze F, et al. Prospective analysis of a sterile, semi-automated tissue biopsy homogenization method in the diagnosis of prosthetic joint infections. *In Vivo* 2017;31:937-42.
27. Lee JE, Kim HJ, Lee SW. The clinical utility of tuberculin skin test and interferon-? release assay in the diagnosis of active tuberculosis among young adults: a prospective observational study. *BMC Infect Dis* 2011;11:96. doi: 10.1186/1471-2334-11-96.
28. Li Y, Jia W, Lei G, Zhao D, Wang G, Qin S. Diagnostic efficiency of Xpert MTB/RIF assay for osteoarticular tuberculosis in patients with inflammatory arthritis in China. *PLoS One* 2018;13:e0198600. doi: 10.1371/journal.pone.0198600.
29. Nocera RM, Sayle B, Rogers C, Wilkey D. Tc-99m MDP and indium-111 chloride scintigraphy in skeletal tuberculosis. *Clin Nucl Med* 1983;8:418-20.
30. Palestro CJ. Nuclear medicine and the failed joint replacement: past, present, and future. *World J Radiol* 2014;6:446-58.
31. Tokumoto JI, Follansbee SE, Jacobs RA. Prosthetic joint infection due to *Mycobacterium tuberculosis*: report of three cases. *Clin Infect Dis* 1995;21:134-6.
32. Wolfgang GL. Tuberculosis joint infection following total knee arthroplasty. *Clin Orthop Relat Res* 1985;201:162-6.
33. Chen WH, Jiang LS, Dai LY. Influence of bacteria on spinal implant-centered infection: an in vitro and in vivo experimental comparison between *Staphylococcus aureus* and *Mycobacterium tuberculosis*. *Spine (Phila Pa 1976)* 2011;36:103-8.
34. Kim SJ, Kim JH. Late onset *Mycobacterium tuberculosis* infection after total knee arthroplasty: a systematic review and pooled analysis. *Scand J Infect Dis* 2013;45:907-14.
35. Titov AG, Vyshnevskaya EB, Mazurenko SI, Santavirta S, Konttinen Y. Use of polymerase chain reaction to diagnose tuberculous arthritis from joint tissues and synovial fluid. *Arch Pathol Lab Med* 2004;128:205-9.
36. Uppal S, Garg R. Tubercular infection presenting as sinus over ankle joint after knee replacement surgery. *J Glob Infect Dis* 2010;2:71-2.
37. Carrega G, Bartolacci V, Burastero G, Finocchio GC, Ronca A, Riccio G. Prosthetic joint infections due to *Mycobacterium tuberculosis*: a report of 5 cases. *Int J Surg Case Rep* 2013;4:178-81.
38. Klein GR, Jacqueline GM. Prosthetic knee infection in the young immigrant patient - do not forget tuberculosis! *J Arthroplasty* 2012;27:1414.e1-4. doi: 10.1016/j.arth.2011.09.020.
39. Held M, Laubscher M, Mears S, Dix-Peek S, Workman L, Zar H, et al. Diagnostic accuracy of the Xpert MTB/RIF assay for extrapulmonary tuberculosis in children with musculoskeletal infections. *Pediatr Infect Dis J* 2016;35:1165-8.
40. Marschall J, Evison JM, Droz S, Studer U, Zimmerli S. Disseminated tuberculosis following total knee arthroplasty in an HIV patient. *Infection* 2008;36:274-8.
41. Egües Dubuc C, Uriarte Ecenarro M, Errazquin Aguirre N, Belzunegui Otano J. Prosthesis infection by *Mycobacterium tuberculosis* in a patient with rheumatoid arthritis: a case report and literature review. *Reumatol Clin* 2014;10:347-9.
42. Su JY, Huang TL, Lin SY. Total knee arthroplasty in tuberculous arthritis. *Clin Orthop Relat Res* 1996;323:181-7.
43. Lusk RH, Wienke EC, Milligan TW, Albus TE. Tuberculous and foreign?body granulomatous reactions involving a total knee prosthesis. *Arthritis Rheum* 1995;38:1325-7.
44. Spinner RJ, Sexton DJ, Goldner RD, Levin LS. Periprosthetic infections due to *Mycobacterium tuberculosis* in patients with no prior history of tuberculosis. *J Arthroplasty* 1996;11:217-22.

45. Al-Shaikh R, Goodman SB. Delayed-onset Mycobacterium tuberculosis infection with staphylococcal superinfection after total knee replacement. *Am J Orthop (Belle Mead NJ)* 2003;32:302-5.
 46. Wang PH, Shih KS, Tsai CC, Wang HC. Pulmonary tuberculosis with delayed tuberculosis infection of total knee arthroplasty. *J Formos Med Assoc* 2007;106:82-5.
 47. Lee CL, Wei YS, Ho YJ, Lee CH. Postoperative Mycobacterium tuberculosis infection after total knee arthroplasty. *Knee* 2009;16:87-9.
 48. Harwin SF, Banerjee S, Issa K, Kapadia BH, Pivec R, Khanuja HS, et al. Tubercular prosthetic knee joint infection. *Orthopedics* 2013;36:e1464-9. doi: 10.3928/01477447-20131021-35.
-