



THE AGA KHAN UNIVERSITY

eCommons@AKU

Department of Anaesthesia

Medical College, Pakistan

January 2018

Is homocystinuria a real challenge for an anesthetist? Are we making a difference?

Muhammad Umar Slote

Aga Khan University, muhammad.slote@aku.edu

Karima Karam Khan,

Aga Khan University, karima.karam@aku.edu

Ausaf Ahmed Khan

Aga Khan University

Mohsin Nazir Butt

Aga Khan University, mohsin.nazir@aku.edu

Follow this and additional works at: https://ecommons.aku.edu/pakistan_fhs_mc_anaesth



Part of the [Anesthesiology Commons](#)

Recommended Citation

Slote, M. U., Khan, K. K., Khan, A. A., Butt, M. N. (2018). Is homocystinuria a real challenge for anesthetist? Are we making a difference?. *Saudi Journal of Anesthesia*, 12(1), 172-173.

Available at: https://ecommons.aku.edu/pakistan_fhs_mc_anaesth/135

Is homocystinuria a real challenge for anesthesiologist? Are we making a difference?

Sir,

Homocystinuria is characterized by impaired synthesis of cystathionine, decreased cysteine, and increased methionine and homocysteine due to cystathionine b-synthase deficiency. Weakened collagen in connective tissue due to decrease cysteine explains lax ligaments and ectopia lentis as being the common presentation in children. Vascular injury through oxidative mechanism and platelet aggregation leading to hypercoagulable state is another manifestation but usually in adults. Accumulated methionine causes increase insulin release resulting in hypoglycemia. These patients usually seek medical attention due to wide spectrum of the disease and are therefore at increased risk for venous thrombosis^[1] and hypoglycemia.^[2]

A 10-year-old school going girl presented with cyanosis of fingertips for a month. Further history and examination was unremarkable, suspecting homocystinuria its levels were advised, and results were much higher (350 $\mu\text{mol/L}$) than normal (5–12 $\mu\text{mol/L}$) confirming the diagnosis. She was started on protein-restricted diet along with aspirin, pyridoxine, folic acid, trimethylglycine, and Vitamin C supplements. Fortunately, she responded and homocysteine levels dropped down (99.8 $\mu\text{mol/L}$) with improvement of symptoms.

Within a month of diagnosis, she started complaining of decreased vision in both eyes. On examination, bilateral ectopia lentis was diagnosed, she was scheduled for lensectomy under general anesthesia. Although lensectomy is a day care procedure, after explaining perioperative risks, patient was electively admitted. Preoperative assessment was done, homocysteine levels were 83.76 $\mu\text{mol/L}$, and written informed consent was taken. Compression stockings were provided and subcutaneous enoxaparin 0.5 mg/kg/dose every 12 hourly was started. Intravenous line was maintained and $\frac{1}{2}$ strength dextrose started and continued perioperatively. Blood glucose level was measured every 4 hourly and routine medications were continued.

In operating room, monitoring for electrocardiogram, noninvasive blood pressure, and oxygen saturation were applied. Intravenous induction with propofol 2 mg/kg, fentanyl 0.2 $\mu\text{g/kg}$, and atracurium 0.6 mg/kg was done. Anesthesia was maintained with a mixture of isoflurane in

air:oxygen in 50:50 ratio to maintain a mixture of air-oxygen of 0.9. Nitrous oxide was avoided, it increases homocysteine levels by inhibiting methionine synthase and is associated with postoperative morbidity and mortality.^[3] Intermittent pneumatic compression stockings were applied over both the legs with pressure of 40–50 mmHg. After completion of 21 min surgery, the patient was extubated and shifted to recovery room.

Patient was discharged on the 2nd postoperative day from hospital. She was doing well with improvement in vision and started going back to school after a week of surgery. Monthly follow-up with homocysteine level was advised. On her last follow-up in February 2017, levels were 4.76 $\mu\text{mol/L}$ which were almost normal.

A perioperative multidisciplinary approach including pediatrician, anesthesiologist, surgeon, and parents is required for these patients to prevent complication and improves outcome. Preoperatively lower levels of homocystinuria <50 $\mu\text{mol/L}$ should be targeted. Maintaining adequate hydration and providing standard antithrombotic measures perioperatively decreases the risk of thrombosis. Rapid recovery after anesthesia using short-acting agents, minimizing the surgical time, and encouraging for early mobilization postoperatively further reduces the complications.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

**MUHAMMAD UMAR SLOTE, KARIMA KARAM KHAN,
AUSAF AHMED KHAN, MOHSIN NAZIR BUTT**

Department of Anaesthesiology, The Aga Khan University Hospital, Karachi, Pakistan

Address for correspondence:

Dr. Karima Karam Khan,
Department of Anaesthesiology, The Aga Khan University Hospital,
P. O. Box 3500, Stadium Road, Karachi, Pakistan.
E-mail: karima.karam@aku.edu


References

1. Morris AA, Kožich V, Santra S, Andria G, Ben-Omran TI, Chakrapani AB, *et al.* Guidelines for the diagnosis and management of cystathionine

Letters to Editor

- beta-synthase deficiency. *J Inherit Metab Dis* 2017;40:49-74.
2. Lowe S, Johnson DA, Tobias JD. Anesthetic implications of the child with homocystinuria. *J Clin Anesth* 1994;6:142-4.
 3. Selzer RR, Rosenblatt DS, Laxova R, Hogan K. Adverse effect of nitrous oxide in a child with 5,10-methylenetetrahydrofolate reductase deficiency. *N Engl J Med* 2003;349:45-50.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

Access this article online	
Website: www.saudija.org	Quick Response Code 
DOI: 10.4103/sja.SJA_274_17	

How to cite this article: Slote MU, Khan KK, Khan AA, Butt MN. Is homocystinuria a real challenge for anesthetist? Are we making a difference?. *Saudi J Anaesth* 2018;12:172-3.

© 2018 Saudi Journal of Anesthesia | Published by Wolters Kluwer - Medknow