



THE AGA KHAN UNIVERSITY

eCommons@AKU

Section of Neurology

Department of Medicine

May 2006

MS in Asian countries

M Wasay

Aga Khan University, mohammad.wasay@aku.edu

I A. Khatri

B Khealani

Aga Khan University, bhojo.khealani@aku.edu

M Sheerani

Follow this and additional works at: https://ecommons.aku.edu/pakistan_fhs_mc_med_neurol



Part of the [Neurology Commons](#)

Recommended Citation

Wasay, M., Khatri, I. A., Khealani, B., Sheerani, M. (2006). MS in Asian countries. *International MS Journal*, 13(2), 58-65.

Available at: https://ecommons.aku.edu/pakistan_fhs_mc_med_neurol/162

MS in Asian Countries

M Wasay¹, IA Khatri², B Khealani¹, M Sheerani¹

¹The Aga Khan University, Karachi, Pakistan; ²University of Medicine and Dentistry of New Jersey, New Jersey, USA

Summary

This paper reviews all available English language literature on MS from Asian countries published between 1970 and 2005. Although limited data are available, the review reveals that western Asia – including the Middle East – has the highest prevalence of MS across the continent, and that MS in Asia largely resembles

conventional MS in western countries. Opticospinal MS (a distinct clinical entity from conventional MS) is more common in eastern Asian regions. Larger epidemiological and genetic studies, with more complete ascertainment in various Asian populations, are needed so that we can understand the diversity of Asian MS.

KEY WORDS:

MULTIPLE SCLEROSIS; OPTICOSPINAL MULTIPLE SCLEROSIS; ASIA; MIDDLE EAST; LITERATURE REVIEW

Introduction

The diversity of MS in Asia has been a topic of debate for years. Although the prevalence of MS varies in different parts of this region (with Israel, the Middle East and west-Asian countries having the highest reported MS prevalences), nationwide MS surveys have rarely been undertaken, and differences in the availability of advanced medical care for MS patients may confound cross-border comparisons.

We believe that there are wide variations in the clinical presentation, male to female ratio and human leucocyte antigen (HLA) typing among MS patients throughout Asia. In addition, it is a matter of debate whether clinical and immunological features of Asian MS patients as a group are distinct from European or North American MS patients. For example, although opticospinal MS (OSMS) is reported to be common in Japanese literature and is sometimes referred to as Asian-type MS, this term may be misleading. Neuromyelitis optica – which is similar to OSMS in many respects – is found globally. It may be that historical variations in diagnostic criteria may be a major factor in the perceived excess of opticospinal-type MS cases in Asia. Moreover, in some Asian countries, OSMS is reported to be rare.

An extensive and systematic review of how MS is reported in Asia has not previously been undertaken. In the present study, therefore, we reviewed published MS-related literature (in the English language) from 22 Asian countries.

Methods

A MEDLINE® search using the key words ‘multiple sclerosis’ and names of countries was performed, covering the period from 1970 to 2005. In total, 432 articles were identified, and papers published in English were included. Only epidemiological studies or case series larger than 15 patients were included. As a result, we reviewed 105 articles. Asian countries were divided into four geographical regions – eastern, southern, western, and central – to identify regional or racial differences in MS across the continent. We compared data from countries within each region in tabulated form. Data were input onto a pre-specified questionnaire that focused on prevalence, male to female ratio, clinical presentation (particularly opticospinal cord involvement) and HLA typing.

Results

Eastern Asia

There is relatively limited literature in the English language on MS from this region. We reviewed papers from Japan,^{1–7} China,^{8,9} Hong Kong,^{10–12} Taiwan,¹³ Singapore,¹⁴ Thailand,^{15,16} Korea¹⁷ and Malaysia.¹⁸ No papers were available from Vietnam, Cambodia or Indonesia.

In this region, the MS prevalence is estimated to be between 0.8 and 2 cases per 100 000 (Table 1),

although not all countries have prevalence data. A recent paper from northern Japan,⁶ however, reported a higher prevalence of 8.57 cases per 100 000 population. An increase in the incidence, change in the diagnostic criteria, and the role of magnetic resonance imaging (MRI) in the diagnosis of MS may explain this apparent rise.

Female preponderance is reported by all series except Kurtzke *et al.*¹⁷ The male to female ratio ranges from 1:1.3 to approximately 1.5. An unusually lower male to female ratio of 1:9.6 was reported from Hong Kong¹⁰ in a 2002 article. The mean age of MS onset across Asia ranges from 25 to 33 years.

Chiemchanya and Vishudhiphan¹⁶ found MS in 17 paediatric patients from Thailand, six of whom developed the disease before 5 years of age. For most patients, disease onset was acute (<1 week) or subacute (1 week to 1 month), and optical symptoms and/or spinal cord involvement at onset and during the disease course were common (Table 1).

The prominence of opticospinal involvement in Asian populations has long been recognized. Many authors have divided MS into opticospinal and conventional forms, based on their differential clinical phenotypes. Kira *et al.*⁵ consider these to be different clinical entities and have also shown immunogenetic differences between the types: they found an association between conventional MS and the DRB1*1501 and DRB5*0101 alleles in MS patients, but not those with opticospinal involvement. A subsequent paper⁴ demonstrated a change in the clinical phenotype of MS in Japan over a 50-year period, identifying an increasing incidence of conventional MS and a concomitant decrease in the incidence of OSMS. Others have reported similar trends.¹⁹

High mortality rates (6–36%) have been described, with the mean duration of illness being between 20 months and 8 years before death in some countries.^{9,11,13,15,16,18} The pathological findings from autopsies (which represent very few cases) show extensive opticospinal involvement. In addition to classic demyelinating lesions, destructive, necrotic lesions with hypercellularity have been found.^{11,13}

Table 1: Characteristics of MS in eastern Asian countries

Lead author/ Reference No.	Year of publication	Population	No. of patients	Prevalence per 100 000	Males:females with MS	Mean age of onset (years)	Opticospinal presentation (%)
Kuroiwa ¹	1975	Japanese	1084	-	1:1.30	33±13	43
Houzen ⁶	2003	Japanese	31	8.57	1:2.90	29.1±14.2	16
Fukazawa ⁷	1992	Japanese	62	-	1:1.95	31	27
Zhao ⁹	1981	Chinese	70	-	1:1.80	30.6	30
Lau ¹⁰	2002	Hong Kong Chinese	53	0.77	1:9.60	29.4	45
Yu ¹¹	1989	Hong Kong Chinese	47	0.88	1:1.80	29	33
Hung ¹³	1976	Taiwanese	25	0.80	1:3.20	32.1	56
Das ¹⁴	1998	Singaporean	21	-	1:3.20	-	43
Jitpimolmard ¹⁵	1994	Thai	50	-	1:4.00	30.4	68
Kurtzke ¹⁷	1968	Korean	22	2	2.1:1.0	25.8	64
Tan ¹⁸	1988	Malaysian	30	2	1:5	29.7	33

-, not reported

Southern Asia

Southern Asia (India, Pakistan, Bangladesh, Sri Lanka, Nepal, the Maldives and Bhutan), Iran, Myanmar and Afghanistan are considered to be areas of low MS prevalence (<5 per 100 000 population).^{20–23} There are no published reports in English from Bangladesh, Nepal, the Maldives, Bhutan, Afghanistan and Myanmar. Literature on the subject is scarce from Pakistan,²⁴ Sri Lanka²⁵ and Iran,^{20,26} although several reports are available from India,^{21,27–34} the majority being hospital-based case series.

There are no nationwide prevalence reports from these countries, although two community-based prevalence studies from India, conducted in the late 1980s, revealed a prevalence of 21–58 per 100 000 population among the Parsi community.^{31,32} MS diagnosis in both of these surveys was based on Schumacher's criteria. In a hospital-based study, Gangopadhyay *et al.*²¹ reported an incidence of 0.62 per 100 000 population based on patients who visit neurology outpatient clinics. Although a population-based survey in Indian Kashmir, conducted in 1986, failed to detect any MS cases, another hospital-based study showed that, over time, the total proportion of MS-related neurology department admissions increased from 1.58% to 2.54%.²⁸ This apparent rise is likely to be multifactorial and has been linked to increased awareness (possibly because of a growing number of neurologists) and the availability of sophisticated diagnostic tools, especially MRI.

In general, research undertaken since the 1990s indicates that MS affects more women than men,³⁵ although, an earlier Indian paper indicated a male preponderance.²⁷ The mean age of MS onset has been reported between 25 and 30 years (Table 2) (ranging from 4.5 years²⁷ to 55 years²¹). Kalanie *et al.*²⁰ found a higher age of onset for patients with progressive disease, especially primary-progressive MS, which is also recognized in Western countries.³⁶

Visual symptoms were the most common presenting features (followed by motor symptoms) in Indian patients,²⁷ while in Iran, motor manifestations were the most common, followed by sensory and visual symptoms in that order.²⁰ Cerebellar involvement was both less common at onset and was relatively less frequent during the course of illness compared with expectations with western-type MS.

Key Points

- The prevalence of MS is high in western Asia and the Middle East compared with eastern or south-eastern Asia
- Opticospinal MS is more common in eastern Asia; patients from western or south-eastern Asia predominantly suffer from western-type MS
- The genetic make-up of patients with MS in western Asia is probably closer to Caucasians than those from eastern Asia
- Large multicentre studies, involving all Asian countries, are needed to understand the epidemiology and course across the continent

The reported frequency of OSMS is 23–33% in the majority of Indian papers,^{21,27, 29} but some studies report as high as 47–58%^{27,28}, whereas OSMS is relatively less common in Iran.²⁰ Common presenting features in Sri Lankan patients were visual and pyramidal, which is consistent with reports of western-type MS.²⁵ No papers describing the clinical manifestations of MS in Pakistan have been published.

Few studies from India have identified an immunological or genetic basis for MS, although one case-control study found that measles infection and exposure to dogs were significantly more common in MS patients than in controls.³⁷ An Iranian study found that 5% of patients had a family history of MS in first-degree relatives,²⁰ but an Indian study failed to detect a single case with a positive MS history in a first-degree relative.³⁰ Various HLA antigens have been noted to be more common in MS patients compared with the general population, although the frequency of HLA markers from people born in this region differs from that reported in western literature. An Iranian study²⁶ found a higher occurrence of HLA DR2, DR15 and A24 in patients with definite MS, with a relative risk of 1.9 for each marker. Indian investigators^{33,34} identified an association between HLA B12 and MS: this antigen was detected in about 80% of cases, compared with about 15% of controls.

There is little information on disability and prognosis in people with MS from this region, although the clinical course of MS tends to be more severe compared with patients from western

Table 2: Characteristics of MS in India and Iran

Lead author/ Reference No.	Jain ²⁷	Syal ²⁸	Mani ²⁹	Gangopadhyay ²¹	Bansil ³⁰	Kalanie ²⁰
Years covered by study	1957–1980	1986–1998	1991–1996	1989–1999	1990s	1996–2001
Population	Indian	Indian	Indian	Indian	Indian	Iranian
No. of patients	354	100	31	45	81	200
Mean age of onset (years)	27	28.5	25.3	30.5	27.5	27.0
Males:females with MS	1:1.32	1:1.32	1:2.10	1:1.50	1:2.25	1:2.50
Optic-spinal presentation (%)	22–58	47	23	33	–	20

–, not reported

countries, with higher percentages of southern Asian patients being severely disabled or bedridden. Some papers have included Expanded Disability Status Scale (EDSS) scores of <5 in most patients,^{20,21} and annual attack rates of 0.4 from Iran²⁰ and 0.63 from India,²⁸ in patients with relapsing–remitting MS. The median EDSS score for Indian patients was 3.7 (mean of 6 years from disease onset),²¹ whereas it was 2.1 in Iran at 5.5 years post onset.²⁰ The Iranian investigators²⁰ also noted that 12% of their patients had EDSS scores <2 at 5 years and 14% had EDSS scores <3 at 10 years.

Western Asia and Middle East

This region can be considered as a medium-risk zone for MS, where the disease prevalence is 5–29 per 100 000 population, as opposed to a high-risk zone (prevalence rates >30 per 100 000 population), or low-risk zone (prevalence <5 per 100 000 population).³⁸ Surveys from the Middle East have demonstrated MS to be less common in this region compared with western Europe and North America, but the prevalence of MS in this region is higher than in any other part of Asia. A limited number of English language papers on MS from this region have been published. We reviewed papers from Kuwait,^{39,40} Turkey,⁴¹ Jordan,⁴² Iraq,^{43,44} Israel,^{45–47} and Saudi Arabia.^{48–50} No English language papers were available from Qatar, United Arab Emirates, Syria, Palestine, Yemen, Bahrain, Oman and Lebanon.

Excluding Israel and Turkey, most west-Asian countries have predominantly Arab populations. The

MS prevalence is approximately 10–20 per 100 000 population (Table 3), compared with 30–80 per 100 000 population in North America. One study, however, reported a higher extrapolated prevalence of 20 per 100 000 Jordanians and 42 per 100 000 Palestinians.⁴² The prevalence of MS in Palestinians is reported to be the highest among Arab countries.^{39,40,42} Palestinians living in Kuwait also have a higher incidence of MS than native Kuwaitis; reasons for this are unclear, but may be related to the admixture of northern European peoples who have settled in this area during migration or invasion. A similar explanation may apply for other non-Arab nations – including the Kurds who live in northern Iraq – who also have higher MS prevalence rates than others native to these regions.^{43,44}

The MS prevalence was found to be 0.4/100 000 in Saudi Arabia.⁴⁸ Israel, however, is considered to be a high-risk zone for MS with a prevalence of more than 30 per 100 000 population,⁴⁵ with higher rates in native Jews of Asian/African origin (52/100 000) than immigrant Jews of Asian/African origin (22/100 000).⁴⁷ In Turkish people, the male to female ratio of MS ranged from 1:1.1 to 1:1.9.⁴¹

The mean age of MS disease onset in this Asian region ranges from 12 to 34 years, although local studies have rarely assessed the signs, symptoms and course of MS. The few that have done so have usually studied Arab populations, in whom the most common symptoms are visual loss and sensory symptoms. Daif *et al.*⁴⁹ reviewed 89 MS cases and Yaqub and Daif⁵⁰ reported 16 cases; both studies

Table 3: Characteristics of MS in the Middle East

Lead author/ Reference no.	Al-Din ³⁹	Al-Din ⁴⁰	Kantarci ⁴¹	Karni ⁴⁷	Al-Din ⁴²
Year of publication	1990	1986	1998	2003	1995
Population	Kuwaitis and Palestinians	Kuwaitis	Turkish	Israel	Jordanians
No. of patients	201	89	1259	272	139
Prevalence per 100 000	10.2	8.33	–	46.2	20.0
Males:females with MS	1:1.7	1.11:1	1:1.19	1.64:1	1:1.9
Mean age of onset (years)	34.7	31.2	27.6±8.8	–	29.6±8.1
Opticospinal presentation (%)	–	27	25	–	28

–, not reported

were undertaken in Saudi Arabia and found the presentation and course of MS in these patients to be no different to western MS. A review of 300 Iraqi patients included clinical and demographic data similar to Caucasian populations.⁴³

A significant association between MS and HLA-DR2 and HLA-DQW1 was found in Palestinians,³⁹ and a population-based cohort of clinically definite MS patients showed a significant association between HLA class II alleles (DRB1, DQA1, DQB1) and MS, in Ashkenazi and non-Ashkenazi Jewish populations.⁴⁵ These data also suggested that DRB1*1501 is the susceptibility allele for MS in the Jewish population.⁴⁵ An Israeli MS study showed that Ashkenazi Jews originating from Northern Europe had the highest $\Delta 32\text{CCR5}$ allele frequency, and that the $\Delta 32\text{CCR5}$ mutation, correlated with slower MS disease progression.⁵¹

Central Asia

Literature from this region published in the English language is limited. We reviewed four papers from Uzbekistan and Tajikistan.^{52–55} The Tajikistani study reviewed 182 patients with conventional MS and suggested climatic factors for pathogenesis.⁵² Three studies by Alaev *et al.*^{53–55} reviewed over 100 MS cases from Uzbekistan, concluding that MS was more common in people of Russian descent compared with those of Uzbeki origin. OSMS was not specifically mentioned in these studies.

Discussion

The present study presents a systematic and comprehensive review of the prevalence and clinical presentation of MS throughout Asia. Our findings are greatly limited by the weakness of available data and the extreme variability of ascertainment bias, which could influence prevalence, gender ratio, disease subtypes, clinical course and genetic associations. The data are also limited by the variable quality of the different studies, small sample sizes, diagnostic difficulties, and incomplete and variable reporting methods. Most studies were hospital-based case series rather than population-based studies, which limited the scope of our review. Although reports or case series from many Asian countries have not been published, we believe that MS exists throughout the Asian region. The MS diagnosis is closely related to the availability of neurologists and diagnostic tools, therefore the prevalence of MS in developing countries may increase when access to these services is more widespread.

The overall prevalence of MS in Asia is lower than in Europe or North America, which may in part be related to genetic factors. In support of this conclusion, the prevalence of MS in Asians born in western countries is lower than that of the native population.^{56,57} Large epidemiological studies are needed to study MS incidence and prevalence, and its natural disease course, in developing Asian countries.

Although OSMS is perceived to be more common in Asia than elsewhere, such findings mostly emerge from research undertaken in eastern Asia rather than across the continent.⁵⁸ Reports from India, Iran, the Middle East and Iraq show OSMS to be rarer in these countries than in eastern Asia. OSMS is uncommon – although it may be more correct to say under-recognized – in North America, because the disease has traditionally been diagnosed only in those with classic features of Devic's disease. This form of MS is more common in African-Americans compared with Caucasian Americans.⁵⁹ Various reports suggest that OSMS is clinically, radiologically and immunologically distinct from conventional MS,^{60–70} although another study suggests that OSMS and neuromyelitis optica may be the same condition.⁷¹ Our literature review is unable to differentiate the true prevalence of neuromyelitis optica or OSMS from cases of western MS with optic and spinal involvement, primarily because the criteria that allow these entities to be differentiated have only recently emerged.

It is generally felt that neuromyelitis optica or OSMS do not respond to interferon therapy. Recent research, however, indicates a trend in favour of high-dose beta interferon therapy for reducing relapses in people with opticospinal MS, although the results were not statistically significant.⁷² Further studies are needed to establish whether patients with opticospinal MS do respond to beta interferon therapy.

Finally, one should note that it is extremely difficult to compare genetic characteristics of western and Asian MS: the Asian subcontinent comprises racially and geographically diverse populations, and few studies from Asian countries have reported HLA typing in MS patients. Available data, however, suggest that the genetic make-up of MS patients from western Asia is probably closer to that of Caucasians than of those from eastern Asia.⁷³

Acknowledgement

The authors are thankful to Dr Sten Fredrikson, Karolinska Institute, Stockholm, Sweden for manuscript review and valuable suggestions.

Conflicts of Interest

No conflicts of interest were declared in relation to this article.

Address for Correspondence

M Wasay, Department of Neurology, The Aga Khan University, Stadium Road, Karachi 74800, Pakistan
E-mail: mohammad.wasay@aku.edu;
mohammadwasay@hotmail.com

Original received: 14 March 2005

Revised received: 26 September 2005

Revised accepted: 9 December 2005

References

- Kuroiwa Y, Igata A, Itahara K, Koshijima S, Tsubaki T, Toyokura Y *et al.* Nationwide survey of multiple sclerosis in Japan. Clinical analysis of 1,084 cases. *Neurology* 1975; **25**: 845–851.
- Kuroiwa Y, Hung TP, Landsborough D, Park CS, Singhal BS, Soemargo S *et al.* Multiple sclerosis in Asia. *Neurology* 1977; **27**: 188–192.
- Kuroiwa Y. Clinical and epidemiological aspects of multiple sclerosis in Japan. *Jpn J Med* 1982; **21**: 135–140.
- Kira J, Kanai T, Nishimura Y, Yamasaki K, Matsushita S, Kawano Y *et al.* Western versus Asian types of multiple sclerosis: immunogenetically and clinically distinct disorders. *Ann Neurol* 1996; **40**: 569–574.
- Kira J, Yamasaki K, Horiuchi I, Ohyagi Y, Taniwaki T, Kawano Y. Changes in the clinical phenotypes of multiple sclerosis during the past 50 years in Japan. *J Neurol Sci* 1999; **166**: 53–57.
- Houzen H, Niino M, Kikuchi S, Toshiyuki F, Nogoshi S, Matsumoto H *et al.* The prevalence and clinical characteristics of MS in northern Japan. *J Neurol Sci* 2003; **211**: 49–53.
- Fukazawa T, Tashiro K, Hamada T, Moriwaka F, Matsumoto A, Shima K *et al.* Multiple sclerosis in Hokkaido, the northernmost island of Japan: prospective analyses of clinical features. *Intern Med* 1992; **31**: 349–352.
- Hou JB, Zhang ZX. Prevalence of multiple sclerosis: A door-to-door survey in Lan Cang La Hu Zu Autonomous County, Yunnan Province of China. *Neuroepidemiology* 1992; **11**: 52.
- Zhao BX, Liu XQ, Guo YP, Yang YC, Huang HF. Multiple Sclerosis. A clinical study of 70 cases. *Eur Neurol* 1981; **20**: 394–400.
- Lau KK, Wong LKS, Li LS, Chan YW, Li HL, Wong Y. Epidemiological study of multiple sclerosis in Hong Kong Chinese: questionnaire survey. *Hong Kong Med J* 2002; **8**: 77–80.
- Yu YL, Woo E, Hawkins BR, Ho HC, Huang CY. Multiple sclerosis amongst Chinese in Hong Kong. *Brain* 1989; **112**: 1445–1467.
- Lau KK, Lee PO, Chan KY, Chan YW, Chin KF. Interferon treatment for multiple sclerosis patients in Hong Kong. *Hong Kong Med J* 2000; **6**: 221–223.
- Hung TP, Landsborough D, Hsi MS. Multiple sclerosis amongst Chinese in Taiwan. *J Neurol Sci* 1976; **27**: 459–484.
- Das A, Puvanendran K. A retrospective review of patients with clinically definite multiple sclerosis. *Ann Acad Med Singapore* 1998; **27**: 204–209.
- Jitpimolmard S, Vejjajiva A. Clinical features and clinical course of multiple sclerosis in Thai patients: a report of 50 cases. *J Med Assoc Thai* 1994; **77**: 239–243.
- Chiemchanya S, Visudhiphan P. Multiple sclerosis in children: a report of 17 Thai pediatric patients. *J Med Assoc Thai* 1993; **76**: 28–33.
- Kurtzke JF, Park CS, Oh SJ. Multiple sclerosis in Korea. Clinical features and prevalence. *J Neurol Sci* 1968; **6**: 463–481.
- Tan CT. Multiple sclerosis in Malaysia. *Arch Neurol* 1988; **45**: 624–627.
- Onodera H, Nakashima I, Fujihara K, Nagata T, Itoyama Y. Elevated plasma level of plasminogen activator inhibitor-1 (PAI-1) in patients with relapsing-remitting multiple sclerosis. *Tohoku J Exp Med* 1999; **189**: 259–265.
- Kalanie H, Gharagozli K, Kalanie AR. Multiple sclerosis: report on 200 cases from Iran. *Mult Scler* 2003; **9**: 36–38.
- Gangopadhyay G, Das SK, Sarda P, Saha SP, Gangopadhyay PK, Roy TN *et al.* Clinical profile of multiple sclerosis in Bengal. *Neurol India* 1999; **47**: 18–21.
- Kurtzke JF, Wallin MT. *Neuroepidemiology. In: Neurology in Clinical Practice: Principles of Diagnosis and Management* (Bradley WG,

- Daroff RB, Fenichel GM, Marsden CD, eds). Boston: Butterworth-Heinemann 2000; pp759-776.
23. Singhal BS. Multiple sclerosis. *Neural India* 1999; **47**: 1-2.
 24. Siddiqui I, Aleem S, Kayani N, Baig S. CSF oligoclonal bands in multiple sclerosis. *J Pak Med Assoc* 2002; **52**: 351-353.
 25. Senanayake B, Ranawaka U, Wijesekera J. Multiple sclerosis in Sri Lanka. *Ceylon Med J* 2001; **46**: 159-160.
 26. Kalanie H, Kamgooyan M, Sadeghian H, Kalanie AR. Histocompatibility antigen (HLA) with multiple sclerosis in Iran. *Mult Scler* 2000; **6**: 317-319.
 27. Jain S, Maheshwari MC. Multiple sclerosis: Indian experience in the last thirty years. *Neuroepidemiology* 1985; **4**: 96-107.
 28. Syal P, Prabhakar S, Thussu A, Seghal S, Khandelwal N. Clinical profile of multiple sclerosis in north-west India. *Neural India* 1999; **47**: 12-17.
 29. Mani J, Chaudhary N, Ravat S, Shah PU. Multiple sclerosis: Experience in neuroimaging era from western India. *Neural India* 1999; **47**: 8-11.
 30. Bansil S, Singhal BS, Ahuja GK, Ladiwala U, Behari M, Friede R et al. Comparison between multiple sclerosis in India and United States: a case-control study. *Neurology* 1996; **46**: 385-387.
 31. Bharucha NE, Bharucha EP, Wadia NH, Singhal BS, Bharucha AE, Bhise AV et al. Prevalence of multiple sclerosis in the Parsis of Bombay. *Neurology* 1988; **38**: 727-729.
 32. Wadia NH, Bhatia K. Multiple sclerosis is prevalent in the Zoroastrians (Parsis) of India. *Ann Neurol* 1990; **28**: 177-179.
 33. Wadia NH, Trikannad VS, Krishnaswamy PR. Association of HLA-B12 with multiple sclerosis in India. *Tissue Antigens* 1980; **15**: 90-93.
 34. Wadia NH, Trikannad VS, Krishnaswamy PR. HLA antigens in multiple sclerosis amongst Indians. *J Neurol Neurosurg Psychiatry* 1981; **44**: 849-851.
 35. Razdan S, Kaul RL, Motta A, Kaul S, Bhatt RK. Prevalence and pattern of major neurological disorders in rural Kashmir (India) in 1986. *Neuroepidemiology* 1994; **13**: 113-119.
 36. Olek MS, Dawson DM. Multiple sclerosis and other inflammatory demyelinating diseases of the central nervous system. In: *Neurology in Clinical Practice; Principles of Diagnosis and Management* (Bradley WG, Daroff RB, Fenichel GM, Marsden CD, eds). Boston: Butterworth Heinemann, 2000; pp1431-1466.
 37. Bansil S, Singhal BS, Ahuja GK, Riise T, Ladiwala U, Behari M et al. Multiple sclerosis in India: a case control study of environmental exposures. *Acta Neurol Scand* 1997; **95**: 90-95.
 38. Kurtzke JF. A reassessment of the distribution of multiple sclerosis. *Acta Neurol Scand* 1975; **51**: 110-157.
 39. Al-Din AS, Khogali M, Poser CM, Al-Nassar KE, Shakir R, Hussain J et al. Epidemiology of multiple sclerosis in Arabs in Kuwait: a comparative study between Kuwaitis and Palestinians. *J Neurol Sci* 1990; **100**: 137-141.
 40. Al-Din AS. Multiple sclerosis in Kuwait: clinical and epidemiological study. *J Neurol Neurosurg Psychiatry* 1986; **49**: 928-931.
 41. Kantarci O, Siva A, Eraksoy M, Karabudak R, Sutlas N, Agaoglu J et al. Survival and predictors of disability in Turkish MS patients. Turkish Multiple Sclerosis Study Group (TUMSSG). *Neurology* 1998; **51**: 765-772.
 42. Al-Din AS, ElKhateeb M, Kurdi A, Mubaidin A, Wriekat A, Al-Shehab A et al. Multiple sclerosis in Arabs in Jordan. *J Neurol Sci* 1995; **131**: 144-149.
 43. Hamdi TI. Multiple sclerosis in Iraq: a clinical and geomedical survey. *J Postgr Med* 1975; **21**: 1-9.
 44. Al-Araji A, Mohammed AI. Multiple sclerosis in Iraq: does it have the same features encountered in Western countries? *J Neurol Sci* 2005; **234**: 47-71.
 45. Kurtzke JF. Multiple sclerosis in time and space - geographic clues to cause. *J Neurovirol* 2000; **6**: S134-S140.
 46. Kwon OJ, Karni A, Israel S, Brautbar C, Amar A, Meiner Z et al. HLA class II susceptibility to multiple sclerosis among Ashkenazi and non-Ashkenazi Jews. *Arch Neurol* 1999; **56**: 555-560.
 47. Karni A, Kahana E, Zilber N, Abramsky O, Alter M, Karussis D. The frequency of multiple sclerosis in Jewish and Arab populations in greater Jerusalem. *Neuroepidemiology* 2003; **22**: 82-86.
 48. al Rajeh S, Bademosi O, Ismail H, Awada A, Dawodu A, al-Freih H et al. A community survey of neurological disorders in Saudi Arabia: the Thugbah study. *Neuroepidemiology* 1993; **12**: 164-178.
 49. Daif AK, Al-Rajeh S, Awada A, Al Bunyan M, Ogunniyi A, Abduljabar M et al. Pattern of presentation of multiple sclerosis in Saudi Arabia: analysis based on clinical and paraclinical features. *Eur Neurol* 1998; **39**: 182-186.
 50. Yaqub BA, Daif AK. Multiple sclerosis in Saudi Arabia. *Neurology* 1988; **38**: 621-623.
 51. Kantor R, Bakhanashvili M, Achiron A. A mutated CCR5 gene may have favorable prognostic implications in MS. *Neurology* 2003; **61**: 238-240.
 52. Pulatov AM. Determinant factor and pathogenesis of multiple sclerosis. *Zh Nevropatol Psikhiatr Im S S Korsakova* 1979; **79**: 142-147.
 53. Alaev BA, Umanskii KG, Samibaev MKh. Clinical picture and course of multiple sclerosis among inhabitants of southwest regions of Uzbekistan. *Zh Nevropatol Psikhiatr Im S S Korsakova* 1985; **85**: 210-214.
 54. Alaev BA. Ethnic factors and their relation to the pathogenesis and geography of multiple sclerosis. *Zh Nevropatol Psikhiatr Im S S Korsakova* 1992; **92**: 9-12.
 55. Alaev BA, Aslanov AM. Climatic factors and clinical polymorphism of multiple sclerosis. *Zh Nevropatol Psikhiatr Im S S Korsakova* 1990; **90**: 9-12.
 56. Murray S, Bashir K, Penrice G, Womersley SJ. Epidemiology of multiple sclerosis in Glasgow. *Scott Med J* 2004; **49**: 100-104.
 57. Dean G, Elian M. Age at immigration to England of Asian and Caribbean immigrants and the risk of developing multiple sclerosis. *J Neurol Neurosurg Psychiatry* 1997; **63**: 565-568.
 58. Kira J. Multiple sclerosis in the Japanese population. *Lancet Neurol* 2003; **2**: 117-127.
 59. Cree BA, Khan O, Bourdette D, Goodin DS, Cohen JA, Marrie RA et al. Clinical characteristics of African Americans vs Caucasian Americans with multiple sclerosis. *Neurology* 2004; **63**: 2039-2045.
 60. Chan KH, Tsang KL, Fong GC, Cheung RT, Ho SL. Idiopathic severe recurrent transverse myelitis: a restricted variant of neuromyelitis optica. *Clin Neurol Neurosurg* 2005; **107**: 132-135.
 61. Ghezzi A, Bergamaschi R, Martinelli V, Trojano M, Tola MR, Merelli E et al. Clinical characteristics, course and prognosis of relapsing Devic's Neuromyelitis Optica. *J Neurol* 2004; **251**: 47-52.
 62. de Seze J, Lebrun C, Stojkovic T, Ferriby D, Chatel M, Vermersch P. Is Devic's neuromyelitis optica a separate disease? A comparative study with multiple sclerosis. *Mult Scler* 2003; **9**: 521-525.
 63. de Seze J, Stojkovic T, Ferriby D, Gaurvit JY, Montagne C, Mounier-Vehier F et al. Devic's neuromyelitis optica: clinical, laboratory, MRI and outcome profile. *J Neurol Sci* 2002; **197**: 57-61.
 64. Fujihara K, Nakashima I, Itoyama Y. Optic-spinal form of multiple sclerosis and immune-mediated myelopathy in Japan. *J Neurol Transm Suppl* 2000; **58**: 205-214.
 65. Wingerchuk DM, Hogancamp WF, O'Brien PC, Weinschenker BG. The clinical course of neuromyelitis optica (Devic's syndrome). *Neurology* 1999; **53**: 1107-1114.
 66. Ono T, Zambenedetti MR, Yamasaki K, Kawano Y, Kamikawaji N, Ito H et al. Molecular analysis of HLA class I (HLA-A and -B) and HLA class II (HLA-DRB1) genes in Japanese patients with multiple sclerosis (Western type and Asian type). *Tissue Antigens* 1998; **52**: 539-542.
 67. Kira J, Kanai T, Nishimura Y, Yamasaki K, Matsushita S, Kawano Y et al. Western versus Asian types of multiple sclerosis: immunogenetically and clinically distinct disorders. *Ann Neurol* 1996; **40**: 569-574.
 68. Haase CG. Devic's neuromyelitis optica. Disease or variants of multiple sclerosis? *Nervenarzt* 2001; **72**: 750-754.
 69. Wu XM, Osoegawa M, Yamasaki K, Kawano Y, Ochi H, Horiuchi I et al. Flow cytometric differentiation of Asian and Western types of multiple sclerosis, HTLV-1-associated myelopathy/tropical spastic paraparesis (HAM/TSP) and hyperIgEaemic myelitis by analyses of memory CD4 positive T cell subsets and NK cell subsets. *J Neurol Sci* 2000; **177**: 24-31.
 70. Weinschenker B. Western vs optic-spinal MS: two diseases, one treatment? *Neurology* 2005; **64**: 594-595.
 71. Weinschenker BG, Fujihara K, Pittock SJ. The Asian optic-spinal form of MS is the same entity as neuromyelitis optica in Caucasians: insights from a novel serum marker. *Neurology* 2004; **62** (Suppl 5): A 480. Abstract P06.061.
 72. Saida T, Tashiro K, Itoyama Y, Sato T, Ohashi Y, Zhao Z, et al. Interferon beta-1b is effective in Japanese RRMS patients: a randomized, multicenter study. *Neurology* 2005; **64**: 621-630.
 73. Fujihara K, Ong BK. MS across continents: challenging our knowledge of MS in Asia and the Middle East. Report from the 2nd MS Forum Pan-Asian Conference, 19-20 November 2004, Ho Chi Minh City, Vietnam. *Int MS J* 2005; **12**: 34-35.