



THE AGA KHAN UNIVERSITY

eCommons@AKU

Medical College Documents

Medical College, Pakistan

1-1-2021

Harassment and mental health in surgical training: A pilot survey of surgical trainees in Pakistan

Russell S. Martins

Aga Khan University, russell.martins@scholar.aku.edu

Syeda Amrah Hashmi

Aga Khan University, amrah.hashmi@scholar.aku.edu

Hina Inam

Aga Khan University, hina.inam@aku.edu

Mahin Janjua

Aga Khan University, mahin.janjua@aku.edu

Mahim Malik

Aga Khan University, mahim.malik@aku.edu

Follow this and additional works at: https://ecommons.aku.edu/pakistan_fhs_mc_mc



Part of the [Cardiovascular Diseases Commons](#), [Gender and Sexuality Commons](#), [Medical Education Commons](#), and the [Psychiatric and Mental Health Commons](#)

Recommended Citation

Martins, R. S., Hashmi, S. A., Inam, H., Janjua, M., Malik, M. (2021). Harassment and mental health in surgical training: A pilot survey of surgical trainees in Pakistan. *JPMA. The Journal of the Pakistan Medical Association*, 71(1(Suppl 1)), S23-S28.

Available at: https://ecommons.aku.edu/pakistan_fhs_mc_mc/188

Harassment and mental health in surgical training: a pilot survey of surgical trainees in Pakistan

Russell Seth Martins,¹ Syeda Amrah Hashmi,² Hina Inam,³ Mahin Binte Naeem Janjua,⁴ Mahim Akmal Malik⁵

Abstract

Objective: To assess harassment among surgical trainees and its effects on mental health, and to explore its association with gender.

Method: The nationwide cross-sectional pilot study was conducted by the Association of Women Surgeons of Pakistan from July to September 2019, and included surgical trainees of either gender working in both public and private hospitals. Data was collected using an anonymous online survey form to assess harassment and self-perceived burnout and depression. Data was analysed using SPSS 22.

Results: Of the 147 respondents, 49(33.3%) were males; 98(66.6%) were females; and 118(80.3%) were residents. Workplace harassment was reported by 80(54.4%) trainees. Among the males it was reported by 24(49%) and among the females by 56(57%) ($p=0.349$). Of those having faced harassment, 9(11.3%) reported it to the administration. Severe self-perceived burnout was reported by 102(69.4%) respondents, and severe self-perceived depression by 69(46.9%). Respondents experiencing bullying were more likely to report severe self-perceived burnout than those not experiencing bullying ($p=0.02$). Multivariable logistic regression showed female gender to be significantly associated with sexual harassment (odds ratio: 4.261 [95% confidence interval: 1.067-17.019]) and severe self-perceived depression (odds ratio: 5.052 [95% confidence interval: 1.187-21.503]). Need for a support group was identified by 134(91.2%) trainees.

Conclusion: An overwhelming need was found for trainee surgeon support groups and other interventions targeted at improving the workplace environment for surgical trainees in Pakistan.

Keywords: Sexual harassment, Nonsexual harassment, Graduate medical education, Mental health, Surgery, Cross-sectional study. (JPMA 70: S-23 [Suppl. 1]; 2021)

Introduction

Harassment, described as any unwelcome behaviour that makes a person feel humiliated, intimidated or offended, is a disturbingly common occurrence in the surgical workplace.¹ Harassment takes many forms and may be verbal, physical, emotional and sexual in nature. Particularly vulnerable due to a unique power differential are surgical trainees, with up to 47% experiencing workplace harassment,^{2,3} with female surgical trainees more likely to suffer harassment, especially of the sexual type.²⁻⁴ Understandably, harassment adversely affects job satisfaction, performance and mental well-being of the trainees, leading to increased risk of burnout, depression, substance abuse disorders and even suicidal ideation.^{2,5-9}

While harassment among surgical trainees has been extensively discussed in the developed countries,^{2,4} such literature is scarce from Pakistan. Surveys conducted by Avan et al. in 2006, Imran et al. in 2010, Nayer-ul-Islam et

al. in 2014, and Zubairi et al. in 2019 have described discrimination, bullying, harassment and mistreatment in doctors and postgraduate trainees overall, but did not focus on surgical trainees specifically.¹⁰⁻¹³ Additionally, Malik et al. in 2016 revealed that more than half of surgical residents experienced high levels of burnout, which negatively affected their job satisfaction.¹⁴ To make matters worse, there is a lack of adequate support to improve workplace environment for surgical trainees in Pakistan, and there is an urgent need for policy-influencing research in this regard.¹⁵ The current pilot survey was planned to assess harassment and its ensuing effects on mental health, and to explore its association with gender among surgical trainees in Pakistan.

Subjects and Methods

The nationwide cross-sectional pilot survey was conducted from July to September 2019 by the Association of Women Surgeons of Pakistan (AWSP). After getting approval from the ethics review committee of Aga Khan University (AKU), Karachi, the anonymous online survey using Google Forms was disseminated via social media platforms, such as Twitter, Facebook and Instagram, and to surgical programmes across Pakistan.

.....
^{1,2}Medical College, ^{3,5}Department of Cardiothoracic Surgery, Aga Khan University Hospital, Karachi, Pakistan, ⁴Department of Surgery, Howard University Hospital, Washington, USA.

Correspondence: Mahim Akmal Malik. Email: mahim.malik@aku.edu

The survey was designed by the research team and pretested for ambiguities prior to distribution to the participants. The target respondents were trainee surgeons in Pakistan, including surgical residents, fellows and senior medical officers.

The survey form had four sections: demographic and work characteristics; experience of harassment at workplace and response to harassment; perceived burnout and depression which asked the participants how burntout/depressed they felt, and their responses were scored on a 5-point Likert scale that were categorised as 1= none, 2-3 = mild-to-moderate, 4-5 = severe; and the perceived need for trainee surgeon support groups.

Data was analysed using SPSS 22. Qualitative variables were presented as frequencies and percentages and were assessed using Chi-squared/Fisher exact tests. Multivariable logistic regression, adjusted for age, position and specialty, was performed with gender as the dependent variable. Odds ratio (OR) and 95% confidence interval (CI) were calculated. $P < 0.05$ was considered statistically significant.

Results

Of the 147 respondents, 49(33.3%) were males; 98(66.6%) were females; 118(80.3%) were residents; and 132(89.8%) were aged 25-35 years (Table-1). Of the total, 68(45%) were from general surgery, followed by subspecialties neurosurgery 18(12.2%), plastic surgery 12(8.2%), orthopaedic surgery 12(7.5%), urology 9(6%),

Table-1: Demographics and work characteristics of the respondents.

Variable	Overall (N=147) n (%)	Males (N=49) n (%)	Females (N=98) n (%)	P-Value
Age				
< 25 Years	3 (2.0)	0 (0)	3 (3.1)	
25-35 Years	132 (89.8)	48 (98.0)	84 (85.7)	0.061
35.1-45 Years	11 (7.5)	1 (2.0)	10 (10.2)	
> 45 Years	1 (0.7)	0 (0)	1 (1.0)	
Marital Status				
Married	68 (46.3)	27 (55.1)	41 (41.8)	
Single	76 (51.7)	31 (42.9)	55 (56.1)	0.308
Divorced	3 (2.0)	1 (2.0)	2 (2.0)	
Province				
Sindh	111 (75.5)	42 (85.7)	69 (70.4)	
Punjab	32 (21.8)	7 (14.3)	25 (25.5)	0.179
Others	4 (2.7)	0 (0)	3 (3.1)	
Position				
Resident	118 (80.3)	41 (83.7)	77 (78.6)	
Senior Medical Officer	19 (12.9)	7 (14.3)	12 (12.2)	0.294
Fellow	3 (2.0)	1 (2.0)	2 (2.1)	
Other	7 (4.8)	0 (0)	7 (7.1)	
Sub-Specialty				
General Surgery	67 (45.6)	17 (34.7)	50 (51.0)	0.061
Other Subspecialty	80 (54.4)	32 (65.3)	48 (49.0)	
Hospital				
Private Sector	93 (63.3)	42 (85.7)	51 (52.0)	< 0.001
Public Sector	54 (36.7)	7 (14.3)	47 (48.0)	
Work Hours				
≤ 80 Hours	74 (50.3)	10 (20.4)	64 (65.3)	< 0.001
> 80 Hours	73 (49.7)	39 (79.6)	34 (34.7)	

Table-2: Harassment and mental health amongst surgical trainees in Pakistan.

Variable	Overall (N=147) n (%)	Males (N=49) n (%)	Females (N=98) n (%)	P-Value
Any Workplace Harassment	80 (54.4)	24 (49.0)	56 (57.1)	0.349
Types of Harassment				
Verbal	72 (49.0)	22 (44.9)	50 (51.0)	0.484
Mental	61 (41.5)	19 (38.8)	42 (42.9)	0.349
Bullying	45 (30.6)	15 (30.6)	30 (30.6)	> 0.999
Physical	17 (11.6)	5 (10.2)	12 (12.2)	0.715
Sexual	19 (12.9)	3 (6.1)	16 (16.3)	0.082
Reporting of Harassment				
Reported to Administration	N = 80 9 (11.3)	N = 24 5 (20.8)	N = 56 4 (7.1)	0.119
Did Not Report	71 (88.7)	19 (79.2)	52 (92.9)	
Burnout				
None	4 (2.7)	1 (2.0)	3 (3.1)	
Mild-to-Moderate	41 (27.9)	11 (22.4)	30 (30.6)	0.521
Severe	102 (69.4)	37 (75.5)	65 (66.3)	
Depression				
None	16 (10.9)	7 (14.3)	9 (9.2)	
Mild-to-Moderate	62 (42.2)	22 (44.9)	40 (40.8)	0.473
Severe	69 (46.9)	20 (40.8)	49 (50.0)	
Need for Support Groups	134 (91.2)	43 (87.8)	91 (92.9)	0.304

Table-3: Multivariable Logistic Regression.

*adjusted for Age, Marital Status, Province, Specialty, Residency Sector, Position.

Variable	Female Gender	
	Adjusted OR (95% CI)	P-Value
Any Harassment	1.570 [0.705-3.495]	0.27
Types of Harassment		
Verbal	1.565 [0.702-3.486]	0.273
Mental	1.509 [0.676-3.369]	0.315
Bullying	0.847 [0.357-2.012]	0.707
Physical	1.326 [0.389-4.521]	0.652
Sexual	4.261 [1.067-17.019]	0.04
Burnout		
None	Reference	-
Mild-to-Moderate	4.730 [0.190-117-663]	0.343
Severe	4.178 [0.178-98.331]	0.375
Depression		
None	Reference	-
Mild-to-Moderate	2.775 [0.648-11.860]	0.168
Severe	5.052 [1.187-21.503]	0.028

OR: Odds ratio; CI: Confidence interval.

cardiothoracic surgery 9(6%), ophthalmology 9(5.4%), otolaryngology 5(3.4%) and paediatric surgery 5(3.4%). The majority of trainees were from a private sector hospital (63.3%). There were more female trainees working at public sector hospitals compared to male trainees ($p < 0.001$).

Workplace harassment was reported by 80(54.4%) trainees. Among the males it was reported by 24(49%) and among the females by 56(57%) ($p = 0.349$). Of those having faced harassment, 9(11.3%) reported it to the administration. Severe self-perceived burnout was reported by 102(69.4%) respondents, and severe self-perceived depression by 69(46.9%). Need for a support group was identified by 134(91.2%) trainees (Table-2). Respondents experiencing bullying were more likely to report severe self-perceived burnout than those not experiencing bullying ($p = 0.02$). Self-perceived depression and burnout were strongly associated with each other ($p < 0.001$).

When in need of emotional support, the majority of trainees reported going to their friends 105(71.4%), followed by parents 76(51.7%) and mentors 34(23.1%). Amongst those experiencing harassment, the commonest response was to ignore the incident 45(56.3%), followed by sharing the incident with a friend 42(52.5%), or seeking support 29(36.3%). Trainees working at private sector hospitals were more likely to experience sexual harassment ($p = 0.042$) compared to those working in public sector hospitals.

Trainees living alone had a significantly higher prevalence of severe burnout compared to those living with their families ($p = 0.018$). Trainees who were currently married were more likely to report severe burnout than those not married ($p = 0.042$). Female trainees currently married were more likely to report severe depression than those not married ($p = 0.013$).

The experience of harassment was most common in paediatric surgery ($n = 4$; 80%), cardiothoracic surgery ($n = 7$; 77.8%), and neurosurgery ($n = 13$; 72.2%), and least common in plastic surgery ($n = 4$; 33.3%), general surgery ($n = 33$; 47.1%), and ophthalmology ($n = 4$; 50%). Sexual harassment was most prevalent in cardiothoracic surgery ($n = 2$; 22.2%) and neurosurgery ($n = 4$; 22.2%). Rates of severe burnout and severe depression were highest in paediatric surgery (both $n = 4$; 80%) and otolaryngology (severe burnout: $n = 5$; 100%; severe depression $n = 3$; 60%).

Multivariable logistic regression showed female gender to be significantly associated with sexual harassment (OR: 4.261 [95% CI: 1.067-17.019]) and severe self-perceived depression (OR: 5.052 [95% CI: 1.187-21.503]) (Table-3).

Discussion

Though the period of surgical education and training is amongst the most formative in a budding surgeon's career, it is also as trainees when young surgeons are most vulnerable to harassment in the workplace. Our pilot survey describes the prevalence and types of harassment faced by surgical trainees in Pakistan, and explores associations with burnout and depression. Our results show that more than half of surgical trainees experience some form of harassment during their training. Alarming, the vast majority of harassment has gone unreported. In addition, 69.4% of trainees reported experiencing severe burnout and 46.9% severe depression. The need for surgical trainee support groups was unanimous, with more than 90% respondents expressing a need for such groups.

In our study, 54.4% of trainees reported facing harassment, most frequently verbal (49%) and mental (41%) harassment, and bullying (30.6%). Rates of harassment were similarly high amongst trainees of both genders, though sexual harassment was significantly associated with female trainees (OR: 4.261). These findings align somewhat with previous literature. A systematic review by Huang et al. revealed a pooled incidence of harassment of 31.2%, with high heterogeneity, and incidences in individual studies ranging from 9.7%-71%¹. The same review also reported a pooled incidence of bullying between 37.7%-40.3%¹. In

Pakistan, previous surveys reveal incidences of bullying and harassment of around 63% amongst junior doctors and trainees, although these are not specific to surgical trainees¹¹⁻¹³. Furthermore, similar to our findings, female surgical trainees have been consistently found to be at higher risk of sexual harassment in the workplace²⁻⁴. The incidence of sexual harassment amongst female trainees in our study was 16.3%, which is similar to the 19.9% reported by Hu et al. in their national survey of surgical residents in the US².

Unfortunately, despite the high prevalence of harassment amongst trainees in our study, only 11.3% reported their experience to their hospital's or institution's administration. This is similar to a previous study in Pakistan, where the majority of victims of bullying did not complain to their administration after experiencing bullying¹¹. Major reasons for not reporting incidents range from being afraid of the consequences of complaining to trying to cope with it oneself¹¹. This only emphasizes the position of vulnerability of surgical trainees, as in many cases the perpetrator of harassment or bullying may be none other than surgical faculty and/or senior residents or fellows^{2,11}. Our results reiterate a call to action for institutions across the country to work toward establishing safe disclosure policies and formulating reporting mechanisms that ensure protection and anonymity for the victim. Lastly, our study found that sexual harassment was more commonly experienced by trainees at private sector hospitals compared to those at public sector/government hospitals and surgical specialties with the highest rates of harassment were Pediatric Surgery, Cardiothoracic Surgery, and Neurosurgery. To the best of our knowledge, there are no studies from Pakistan comparing harassment and mistreatment between private hospitals and public hospitals, or between surgical sub-specialties, and these findings highlight areas requiring further exploration.

Our results revealed that 69.4% of surgical trainees reported severe self-perceived burnout, and 46.9% severe self-perceived depression, with the risk of severe depression significantly higher in women (OR: 5.052). Although our survey did not use validated tools to measure burnout and depression, our results based on a self-perceived rating of burnout and depression are comparable to previous studies. A study by Bui et al. across urban academic surgery programs in the US reported incidences of burnout and depression to be 63.2% and 36.6%, respectively¹⁶. A meta-analysis by Low et al. similarly revealed a pooled incidence of burnout of 52.27% amongst surgical residents¹⁷. In the national survey of surgical residents in the US, Hu et al. reported a

prevalence of weekly burnout symptoms amongst 38.5% of trainees, with these symptoms being associated with harassment and mistreatment². This is corroborated by our study, as trainees experiencing bullying were more likely to report severe self-perceived burnout as compared to those not experiencing bullying. Furthermore, our study revealed self-perceived burnout and depression amongst surgical trainees to be strongly associated with each other, and this is also supported by previous literature^{18,19}. In addition, rates of severe self-perceived burnout and depression in our study were highest in Pediatric Surgery and Otolaryngology. This is in contrast to the study by Bui et al., which reported the highest rates of burnout and depression in Obstetrics and Gynecology, Orthopedic Surgery, and Neurosurgery¹⁶, and a meta-analysis by Low et al. which reported the highest incidence of burnout in General Surgery¹⁷.

When in need of emotional support, the majority of respondents in our study reported going to their friends (71.4%) or parents (51.7%), with less than a quarter going to their mentors. Similarly, responses to harassment also most commonly included ignoring the incident (56.3%) or sharing it with a friend (52.5%). Moreover, respondents wholeheartedly expressed the need for institutional trainee support groups to help with stress control (85.1%), mentoring (82.1%), and work-life balance (67.9%). These results serve to illustrate a very serious issue across most postgraduate training programs in Pakistan: the lack of institutional support towards trainees in matters such as reporting and taking action against harassment, safeguarding mental well-being, and imparting mentorship. Although the Protection Against Harassment of Women at Workplace Act (2010)²⁰, the Punjab Healthcare Commission Act (2010)²¹ and the Sindh Healthcare Commission Act (2013)²² emphasize on the need to report and act upon harassment faced in the healthcare workplace, very few hospitals and institutions have progressed towards developing formal reporting and investigatory systems. Moreover, most surgical training programs in the country lack platforms that make trainee well-being an institutional priority. When judiciously implemented, simple interventions such as mentorship and coaching programs for professional and personal development, faculty wellness champions, peer support programs, process groups, and stress-relieving group activities, may all help promote trainee mental wellness in the surgical workplace^{16,23}. It is of utmost importance that the medical and surgical community in Pakistan work at the institutional and national level towards the development of zero tolerance harassment policies and interventions to resolve harassment incidents and protect the mental well-being of surgical trainees. The College of

Physicians and Surgeons Pakistan (CPSP) is suitably placed to coordinate anti-harassment and mental health policy nationally across hospitals and institutions in Pakistan. Additionally, the CPSP must also use existing databases of trainee emails and contacts to conduct comprehensive, nationwide assessments of trainee mistreatment and mental health, so as to build on the results of this pilot survey to develop evidence-based action plans to protect surgical trainees across the country.

Limitations of our pilot survey include its sample size which was restricted due to the lack of access to a universal platform for survey dissemination amongst surgical trainees across the country. This was also the reason for inadequate representation from provinces other than Sindh and Punjab. Moreover, the majority of female respondents could be a result of a response bias due to the survey being conducted under an organization titled the Association of Women Surgeons in Pakistan. Lastly, our survey did not use any full validated tools to measure burnout and depression, but rather gauged respondents' perceptions about their own mental wellbeing. Nevertheless, the findings of our pilot survey lay a solid foundation and reference for future related research and policy, and represent an important preliminary step for a better workplace for surgical trainees in Pakistan.

Conclusion

Harassment, most frequently verbal and mental, is experienced by a large proportion of surgical trainees in Pakistan, with most cases passing unreported to the administrations. Although overall harassment is similarly prevalent amongst male and female trainees, women are at a greater risk of sexual harassment. In addition, many surgical trainees also face burnout and depression, which may be associated with certain forms of harassment, such as bullying. Lastly, there was an overwhelming need for trainee surgeon support groups, and other policies and interventions targeted at improving the workplace environment for surgical trainees in Pakistan.

Acknowledgements: We are grateful to Taleaa Masroor, a medical student at the Aga Khan University, for assisting in data collection and to the members of the Association of Women Surgeons of Pakistan (AWSP) for their assistance with the creation, piloting and dissemination of the survey at the national level.

Disclaimer: None.

Conflict of Interest: None.

Source of Funding: None.

References

- Huang Y, Chua TC, Saw RPM, Young CJ. Discrimination, Bullying and Harassment in Surgery: A Systematic Review and Meta-analysis. *World J Surg* 2018;42:3867-73. doi: 10.1007/s00268-018-4716-5.
- Hu YY, Ellis RJ, Hewitt DB, Yang AD, Cheung EO, Moskowitz JT, et al. Discrimination, Abuse, Harassment, and Burnout in Surgical Residency Training. *N Engl J Med* 2019;381:1741-52. doi: 10.1056/NEJMsa1903759.
- Nukala M, Freedman-Weiss M, Yoo P, Smeds MR. Sexual Harassment in Vascular Surgery Training Programs. *Ann Vasc Surg* 2020;62:92-7. doi: 10.1016/j.avsg.2019.05.011.
- Halim UA, Riding DM. Systematic review of the prevalence, impact and mitigating strategies for bullying, undermining behaviour and harassment in the surgical workplace. *Br J Surg* 2018;105:1390-97. doi: 10.1002/bjs.10926.
- Aldrees T, Hassouneh B, Alabdulkarim A, Asad L, Alqaryan S, Aljohani E, et al. Burnout among plastic surgery residents. National survey in Saudi Arabia. *Saudi Med J* 2017;38:832-6. doi: 10.15537/smj.2017.8.18346.
- Chow OS, Sudarshan M, Maxfield MW, Seese LM, Watkins AA, Fleishman A, et al. National Survey of Burnout and Distress among Cardiothoracic Surgery Trainees. *Ann Thorac Surg* 2020;2020:0003-4975(20)31358-8. doi: 10.1016/j.athoracsur.2020.06.062. [ahead of print]
- Elmore LC, Jeffe DB, Jin L, Awad MM, Turnbull IR. National Survey of Burnout among US General Surgery Residents. *J Am Coll Surg* 2016;223:440-51. doi: 10.1016/j.jamcollsurg.2016.05.014.
- Patti MG, Schlottmann F, Sarr MG. The Problem of Burnout Among Surgeons. *JAMA Surg* 2018;153:403-04. doi: 10.1001/jamasurg.2018.0047.
- Williford ML, Scarlet S, Meyers MO, Luckett DJ, Fine JP, Goettler CE, et al. Multiple-Institution Comparison of Resident and Faculty Perceptions of Burnout and Depression During Surgical Training. *JAMA Surg* 2018;153:705-11. doi: 10.1001/jamasurg.2018.0974.
- Avan BI, Raza SA, Khokhar S, Awan F, Sohail N, Rashid S, et al. Residents' perceptions of work environment during their postgraduate medical training in Pakistan. *J Postgrad Med* 2006;52:11-8.
- Imran N, Jawaid M, Haider II, Masood Z. Bullying of junior doctors in Pakistan: a cross-sectional survey. *Singapore Med J* 2010;51:592-5.
- Nayyer-ul-Islam, Yousuf-Islam M, Farooq MS, Mazharuddin SM, Hussain SA, Umair-ul-Islam. Workplace violence experienced by doctors working in government hospitals of Karachi. *J Coll Physicians Surg Pak* 2014;24:698-9.
- Zubairi AJ, Ali M, Sheikh S, Ahmad T. Workplace violence against doctors involved in clinical care at a tertiary care hospital in Pakistan. *J Pak Med Assoc* 2019;69:1355-9.
- Malik AA, Bhatti S, Shafiq A, Khan RS, Butt UI, Bilal SM, et al. Burnout among surgical residents in a lower-middle income country - Are we any different? *Ann Med Surg (Lond)* 2016;9:28-32. doi: 10.1016/j.amsu.2016.05.012.
- Shiwani MH, Elenin H. Bullying and harassment at workplace: are we aware? *J Pak Med Assoc* 2010;60:516-7.
- Bui AH, Ripp JA, Oh KY, Basloe F, Hassan D, Akhtar S, et al. The impact of program-driven wellness initiatives on burnout and depression among surgical trainees. *Am J Surg* 2020;219:316-21. doi: 10.1016/j.amjsurg.2019.10.027.
- Low ZX, Yeo KA, Sharma VK, Leung GK, McIntyre RS, Guerrero A, et al. Prevalence of Burnout in Medical and Surgical Residents: A Meta-Analysis. *Int J Environ Res Public Health* 2019;16:e1479. doi: 10.3390/ijerph16091479.

18. Dyrbye LN, West CP, Satele D, Boone S, Tan L, Sloan J, et al. Burnout among U.S. medical students, residents, and early career physicians relative to the general U.S. population. *Acad Med* 2014;89:443-51. doi: 10.1097/ACM.0000000000000134.
 19. Mata DA, Ramos MA, Bansal N, Khan R, Guille C, Di Angelantonio E, et al. Prevalence of Depression and Depressive Symptoms Among Resident Physicians: A Systematic Review and Meta-analysis. *JAMA* 2015;314:2373-83. doi: 10.1001/jama.2015.15845.
 20. Pakistan Go. Protection against Harassment of Women at Workplace Act (2010). [Online] 2010 [Cited 2020 September 29]. Available from URL:
 21. http://www.na.gov.pk/uploads/documents/1399368475_218.pdf Punjab Go. The Punjab Healthcare Commission Act 2010. [Online] 2010 [Cited 2020 September 29]. Available from URL: <http://punjablaws.gov.pk/laws/2434.html>
 22. Sindh Go. The Sindh Healthcare Commission Act 2013. [Online] 2014 [Cited 2020 September 29]. Available from URL: <http://www.pas.gov.pk/uploads/acts/Sindh%20Act%20No.VII%20of%202014.pdf>
 23. Baker K, Sen S. Healing Medicine's Future: Prioritizing Physician Trainee Mental Health. *AMA J Ethics* 2016;18:604-13. doi: 10.1001/journalofethics.2016.18.6.medu1-1606.
-