



12-2017

## Dementia survey among Attendees of a Dementia Awareness Event in Karachi, Pakistan

Qurat ul Ain Khan

*Aga Khan University, Karachi*

Yusra Hanif Khan

*Multan Medical and Dental College*

Mohammad Zaman Khan

*Aga Khan University, Karachi*

Shireen Najam

*Aga Khan University, Karachi*

Follow this and additional works at: <https://ecommons.aku.edu/pjns>

 Part of the [Neurology Commons](#)

### Recommended Citation

Khan, Qurat ul Ain; Khan, Yusra Hanif; Zaman Khan, Mohammad; and Najam, Shireen (2017) "Dementia survey among Attendees of a Dementia Awareness Event in Karachi, Pakistan," *Pakistan Journal of Neurological Sciences (PJNS)*: Vol. 12 : Iss. 4 , Article 8.

Available at: <https://ecommons.aku.edu/pjns/vol12/iss4/8>

# Dementia survey among Attendees of a Dementia Awareness Event in Karachi, Pakistan

Qurat ul Ain Khan<sup>1</sup>, Yusra Hanif Khan<sup>2</sup>, Mohammad Zaman Khan<sup>3</sup>, Shireen Najam<sup>4</sup>,

<sup>1,2,3,4</sup> Department of Psychiatry, Aga Khan University, Karachi, Department of Psychiatry, Multan Medical and Dental College

**Corresponding to:** Qurat ul Ain Khan M.D, Department of Psychiatry, Second floor, Faculty Office Building, Aga Khan University, Stadium Road, Karachi. Email address: qurat.khan@aku.edu

**Date of submission:** August 22, 2017 **Date of revision:** September 11, 2017 **Date of acceptance:** September 30, 2017

## ABSTRACT

### BACKGROUND

There is little information available about peoples' knowledge regarding symptoms of dementia, dementia diagnosis, and care arrangements for dementia patients in Pakistan.

**Aims:** We aimed to find out if patients with dementia and their caregivers were aware of diagnosis, the most common and most distressing symptoms, and care arrangement for these patients.

### METHOD

We conducted a pilot survey among the attendees of a dementia awareness session held at the Aga Khan University Hospital in Karachi, Pakistan by using a self-reported questionnaire.

**Results:** Out of the 38 participants who returned filled questionnaires, about half identified the type of dementia as "Alzheimer's" and memory loss was reported as the most prominent and most distressing symptom. "Watching TV" was reported as the most common activity of patients with dementia. Almost all caregivers identified themselves as primary caregivers, living in the same house with the patient, and bearing financial cost of care. About 50% of all participants reported the activity as helpful and showed willingness to participate in such activities in future.

### CONCLUSIONS

The results from this study provide important information about common man's knowledge about dementia symptoms and care arrangement in Pakistan, and has important implications for stakeholders.

### DECLARATION OF INTEREST

Authors report no financial relationship or any conflict of interest. This is a non-funded study.

### KEYWORDS

Dementia, survey, lower and middle income countries, Karachi, Pakistan

### INTRODUCTION

According to Alzheimer's Disease International World Report 2016 over 47 million people worldwide have dementia and this number is expected to rise to more than 131 million by 2050. (Alzheimer's Disease International, 2016) In 2015, 58% of the total dementia population was estimated to reside in low and middle income countries (LMICs) which is expected to rise to 63% by 2030 and 68% by 2050. (Prince et al., 2015) Dementia and related issues in LMICs are different than in the West including assessment, diagnosis, awareness, caregiver characteristics, resources, and policy and strategies. Therefore research done in the

West may not be applicable to LMICs in all aspects and contextualized research is needed for tailored solutions in such settings. (Ferri et al., 2017) Pakistan is a low middle income country in South Asia with a rapidly aging population. It is currently the sixth most populous country in the world and is expected to become the third most populous by 2050 (Khan, 2014) with 10% of the country's total population estimated to be over 60 years of age (WHO World Aging Report, 2013). Islam is the religion of the state with 97 % of the population being Muslim (Khan et al., 2016) with a very low literacy rate of 56% and low GDP per capita (Rehman et al., 2015). Among several factors identified as contributors to peoples' understanding of different medical conditions, religious and

cultural beliefs, level of education, and living arrangements are also important ones that play a role in explanatory models of disease symptomatology (Mahoney et al., 2005 Dec) and outcomes (Zheng et al., 2011). In Pakistan, joint family system prevails and people with dementia are mostly cared for at home by family members and informal caregivers. A qualitative study conducted in Lahore, Pakistan interviewed 8 dementia caregivers to explore thinking patterns and physical health, and identified maladaptive cognitive patterns, fatigue, and sleep disturbances as causes of caregiver distress (Ali, S. et al., 2015). Caregiver distress is also often associated with type, severity, and frequency of dementia symptoms (Dawood, S. 2016).

Caregiver characteristics and relationship with patient may also differ in different geographical regions, affecting the dynamics, quality of care, and care burden (Ali, S. et al., 2015). In Pakistan caregiver role is usually assigned to women (Ahmad, K 2012) who may be spouses, daughters, or daughter in laws of the affected person. Due to the unique characteristics of this society and dearth of dementia related research, it is important to study peoples' knowledge about dementia, cultural beliefs, most common and difficult to manage symptoms, and resources available.

We designed a 15 item questionnaire survey to be filled out by the participants of a dementia awareness program in Karachi to study patients' knowledge, care arrangement, and caregiver characteristics.

## METHODS

A dementia public awareness event was organized on International Alzheimers Dementia Day in October 2015 at the Aga Khan University Hospital in Karachi. The event was open and free for public with no prior registration required.

Each attendee was handed a paper questionnaire at the entrance of the auditorium which were collected by the volunteers at the end of the session. Participants were also asked to write name, age, and contact number/email on the questionnaire. The questionnaire included 15 questions, written in English and each question was followed by multiple choices to choose from; some questions were required to be answered as "yes" or "no". 5 out of 15 items were about dementia patients such as the type of dementia, most prominent, and most distressing symptoms. 2 questions were about the hobbies or activities that the patient engaged in. The next 7 questions were about the care arrangement such as identifying the primary caregiver, living arrangement, time spent with the patient, person bearing the financial needs of the patient, any other relatives involved in patient care, and paid help available. 2 questions were about the participants' opinion about the awareness session.

The study was approved by the Aga Khan University Research Ethics Review Committee.

Data collected was entered and assessed via SPSS 14.0.

## RESULTS

Fifty people attended the session out of which 38 returned filled questionnaires. 11 out of 38 identified themselves as dementia patients and 8 identified themselves as caregivers (50% of the total sample). 19 participants (50%) did not identify with either group and were people from the general public. 9 out of 19 (47%) patients/caregivers identified the type of dementia as Alzheimers and the rest were not aware with the type of dementia. All 19 patients/caregivers with dementia identified forgetfulness as the most prominent as well as the most distressing symptom. Anger/aggression was marked as the second most prominent symptom by all 8 caregivers and 5 dementia patients. Question about engaging in hobbies/activities was marked positive by 7 patients and 5 caregivers (12/19, 63%). The most common activity reported was "watching TV" (10/12) (Table 1).

Among the caregiver related questions, almost all identified themselves as primary caregivers and half of them identified patient as their mothers. 7/8 caregivers reported bearing the financial cost related to patient care and living in the same house with the patient. 3 patients reported bearing financial cost themselves. 7/8 caregivers also reported spending more than 2 hours with the patient every day, and involvement of other people in patient's care. 6/19 patients/caregivers reported having paid help to care for the patient. The activity was reported to be helpful by 17/38 (45%) among which 6 were patients, 5 were caregivers and 6 people from general public. 55% of the participants showed willingness to participate in such activities in future. (Table 2).

## DISCUSSION

This survey was conducted to study knowledge about dementia and care arrangements among the attendees of a dementia awareness event held at a tertiary care university hospital in Karachi. The information obtained from this study reflects that patients/caregivers may be more familiar with Alzheimers dementia than other dementias. It is however also known that Alzheimers is the most common type of dementia (Karantzoulis et al., 2011). It may also mean that due to lack of expertise among health care providers in this domain many people are diagnosed with Alzheimers dementia than other types of dementias. This information is important in devising public awareness campaigns and improving healthcare provider education. Forgetfulness was identified as the most common and most distressing symptom of dementia by all patients and caregivers in our survey. This is similar to the results from dementia

registry study published in Islamabad, Pakistan (Ahmad, A. et al., 2011) which reported 81% people with dementia presenting with memory loss as the main symptom. This provides good insight into peoples' knowledge about recognizing forgetfulness as dementia symptom, and interventions or management strategies based around this symptom would also be desirable as it is also reported to be the most distressing symptom. This is an interesting finding as some data from western literature suggests that forgetfulness is the most frequent but least distressing symptom (Fauth et al., 2014), whereas disruptive behavior was least frequent but most distressing. In joint family socio-cultural setting in Pakistan and other South Asian countries where there may be multiple caregivers available, disruptive behavior may be less distressful as compared to a single caregiver taking care of the patient (Ali, S. et al., 2015). Anger/aggression was marked as dementia symptom by all caregivers and less number of patients themselves where as low mood/anxiety was reported as prominent and also distressing symptom by more number of patients than caregivers. World literature also suggests that anger/aggression contributes significantly to caregiver burden (Cheng, ST. 2017), and in a country like Pakistan where almost all dementia carers are informal carers and family members of patients, it is not surprising that this is reported by caregivers more than patients themselves. Prevalence of depression (García-Alberca, JM., 2017) and anxiety (Orgeta, V. et al., 2015) is also reported to be high in dementia patients in world literature and compromises the quality of life of these patients. Difficulty in performing ADLs is reported as equally distressful by patients and caregivers in our study. This was also identified as a theme in another study addressing caregiver distress in families of dementia patients in Rawalpindi, Pakistan (Qadiret al., 2013). This is known to increase dependence, caregiver burden, and mortality in general (Bürgeet al., 2017) and may be more difficult to manage in a resource poor setting such as Pakistan and other LMICs (Maestre, 2012).

More than half of patients/caregivers in our survey reported that the patient engaged in hobbies or activities, and watching TV was reported to be the most common activity. Exercise and listening to music were reported by the least number of people. This is in keeping with the sedentary life style of elderly population in Pakistan and is particularly the case in people with dementia. According to the conventional role of "being old" in Pakistani society people are generally expected to stay at home after retirement,

spend time praying and performing religious activities, "look religious" such as by growing beard, giving up "young attire" and sticking to more conservative forms of dressing, spend time with grand kids, stay passive, give up responsibilities, and have the next generation take over before they die (Vertejee et al). Community organizations, social clubs, libraries, parks, gymnasiums, old age societies, and day care centers are almost nonexistent (Alam et al., 2016). Elderly men, however usually go to the local mosque for prayers five times a day. Watching TV is the most accessible and affordable activity available to most especially in urban setting. This information also brings to attention a very important area to explore in caring for such patients and to devise methods at the community level to identify and engage them with the health care system such as through community health workers, workers in local mosques, or by using electronic media. Physical exercise was only reported by 3 participants of the survey as an activity. The role of physical exercise in prevention or modification of dementia and brain aging is well recognized (Ahlskoget al., 2011) (Ruthirakuhanet al., 2012); however, there are multiple cultural and social factors, non-availability of space, unsafe environment etc. which make this an unfavorable practice in local setting. Similarly listening to music was also reported by just 3 participants as an activity. This may also have cultural and social explanations among others. This is important contextualized information and emphasizes the need of devising culturally acceptable interventions to prevent and manage cognitive decline in such settings.

Among the caregiver related questions about half identified themselves as children of patients with dementia in our study. A study from Lahore, Pakistan (Ali, S. et al., 2015) reported spouse followed by children as primary caregivers. In a WHO report, children/children in law were identified as primary caregivers of most dementia patients in rural and urban areas of India (World Health Organization (WHO), 2017). In our study most caregivers reported bearing financial cost related to patient care, living in the same house with the patient, spending more than 2 hours with the patient every day, and involvement of other people in patient's care. In traditional Pakistani society (Ali, S. et al., 2015) as well as in India (World Health Organization WHO, 2017) majority of patients with dementia and their adult children live in the same house and may have more than one family members involved in their care. Due to lack of social security, financial support, and public health insurance systems, older people in general and people with dementia in particular are financially dependent on their caregivers as indicated in our survey. About one third patients in our survey reported to have paid help available, however as this was a self-reported, non-

descriptive questionnaire we are not able to comment on the type of help, but having maids for domestic help in Karachi and other urban areas is very common and available at low wages; it is highly likely that these are unprofessional and untrained domestic workers. This scenario may be completely different in low socio-economic and rural areas of Pakistan, as also reported in a study in rural India with 0% of people having paid help available (Brinda et al., 2014). These are informal and untrained caregivers and have limited understanding of dementia symptoms. They may also have other professional and domestic responsibilities which may compromise patient care and vice versa. Impact of joint family system, family dynamics, and financial needs of caregiving are important aspects to explore in future research.

About half of the total number of participants reported that awareness session was helpful and showed willingness to participate in such activities in future. This is important to recognize information needs of caregivers and to design effective supportive interventions and educational activities. Based on this finding, we started a Dementia Education and Support Program for Caregivers at the AKUH which is ongoing. A study conducted in Southern India also revealed that family caregivers of people with dementia reported feeling distressed and wished to participate in educational programs but many had unrealistic expectations for what they wanted to learn (Narayan et al., 2015).

This study has many limitations. First is the small number of participants. The total number of people (patients/caregivers/general public) who attended the session was 50. This is a small number keeping in view the fact that Karachi is a very large and populous city and the event was advertised in two leading newspapers in both English and Urdu languages, the hospital is located in an accessible and densely populated area, and the event was free with no prior registration required. This could have many explanations such as less awareness about dementia among public with common belief in Pakistani society that forgetfulness is a part of normal aging, and stigma attached to the issue. Due to small sample size we recommend replicating the findings by large scale surveys. Second the findings from these participants may not be generalizable as the survey is done in an urban setting, and in a private hospital which caters to affluent class. However this was a public event which was advertised in local newspapers and was free for all to make it easy for general public to attend. Third, the questionnaire was written in English language and people with no or low formal education or those unable to read and understand English may not have participated in the survey. Urdu is the national language of the country and is read and understood more commonly than English which is considered a

foreign or second language. Fourth people with dementia may have had difficulty reading, understanding, remembering, or providing correct information due to cognitive deficits. Fifth we think it would have been useful to design separate questionnaires for patients, caregivers, and general public to reduce ambiguity and missing information and also because of the fact that people with dementia and caregivers may have different opinions and perceptions about symptoms, burden etc. Adding open ended or descriptive answer choices and using mixed method study design would be more useful for such research in future studies. However we think that such surveys may be very useful in exploring awareness about dementia and care arrangements for those suffering from dementia and need to be replicated in different settings in the country. For future work, questions pertaining to attitudes and awareness of advance care planning could be explored as well. Advance care planning would be especially relevant for the early stages of dementia, when patients still retain mental decision capacity. Advance care planning has also been shown to reduce decisional conflict and caregiver stress (Ng, QX et al., 2017).

**Conclusion:** The results from this study provide important information about common man's knowledge about dementia symptoms and care arrangement in Pakistan, and has important implications for stakeholders.

**Table 1. DEMENTIA SYMPTOMS**

<b>Variables</b>	<b>N=38 n (%)</b>
1. Informant Dementia patients Caregivers of dementia patients Others	11 (29.9) 8 (19.4) 19 (50.6)
2. Type of Dementia Alzheimer's Lewy Body dementia Fronto-temporal dementia Vascular dementia Don't know Not specified/Don't Know	9 (23.7) 0 0 0 17 (44.7) 12 (31.6)
3. Most prominent symptoms Memory problem/forgetfulness Anger/aggression Low mood/anxiety Suspiciousness Hearing voices/seeing things	26 (68.4) 13 (34.2) 13 (34.2) 5 (13.2) 9 (27.7)
4. Most difficult to manage/distressful symptoms Memory problem/forgetfulness Depression/anxiety Suspiciousness Hearing voices/seeing things Anger/aggression Confusion/wandering/leaving the house Inability to walk/falls Difficulty/inability to control urine/bowel, bathing/changing clothes	23 (60.5) 15 (39.5) 5 (13.2) 7 (18.4) 6 (15.8) 7 (18.4) 4 (10.5) 8 (21.1)
5. Does patient engage in any hobbies/activities he/she likes? Yes No Missing data	12(31.6) 8(21.1) 18(47.4)
6. If yes, which one? Watching TV Listening to music Reading Going out for shopping/eating Exercise	10(26.3) 03(7.9) 07(18.4) 08(21.5) 03(7.9)

**Table 2. CAREGIVER RELATED QUESTIONS**

<b>Variables</b>	<b>N=38 n (%)</b>
1. Are you the primary caregiver for the patient? Yes No Missing data	07 (18.4) 12 (31.6) 19 (50.0)
2. Relationship of patient with caregiver Wife Father Mother Daughter Father/mother in law Any other	1 (14.2) 1 (14.2) 4 (57.1) 1 (14.2) 1 (14.2) 1 (14.2)
3. Do you bear financial costs related to patient care? Yes No	10 (26.3) 6 (15.8)
4. Do you and the patient live in the same house? Yes No Missing data	14 (36.8) 09 (23.7) 15 (39.5)
5. Number of hours spent with patient every day Less than 30 mins 31-60 mins Two hours More than two hours	5 (13.2) 2 (5.3) 1 (2.6) 8 (21.1)
6. Other than you how many other people are involved in patient's care? None One Two Three Four Five	0 2 (5.3) 1 (2.6) 1 (2.6) 3 (7.9) 2 (5.3)
7. Do you have servant/maid/paid help to care for patient? Yes No Missing data	06 (15.8) 16 (42.1) 16 (42.1)
8. Do you think today's activity was helpful? Yes No Missing data	17 (44.7) 03 (7.9) 18 (47.4)
9. Would you like to participate in such activities in future? Yes Missing data	21 (55.3) 17 (44.7)

## REFERENCES:

1. Ahmad A, Khatri I, Siddiqui M, Khan N, Kamal S, Mehboob N. Dementia in Pakistan: initial results from our registry at a tertiary care hospital. *Alzheimer's & Dementia*. 2011 Jul 1;7(4):S767. Ahmad K. Informal Caregiving to Chronically Ill Older Family Members: Caregivers' Experiences and Problems. *South Asian Studies*. 2012 Jan 1;27(1):101.
2. Ahlskog JE, Geda YE, Graff-Radford NR, Petersen RC. Physical exercise as a preventive or disease-modifying treatment of dementia and brain aging. In *Mayo Clinic Proceedings* 2011 Sep 1 (Vol. 86, No. 9, pp. 876-884). Elsevier.
3. Alam A, Ibrar M, Khan P. Socio-Economic and Psychological Problems of the Senior Citizens of Pakistan.
4. Ali S, Bokharey IZ. Maladaptive cognitions and physical health of the caregivers of dementia: An interpretative phenomenological analysis. *International journal of qualitative studies on health and well-being*. 2015 Jan 1;10(1):28980.
5. World Alzheimer Report 2016. London: Alzheimer's Disease International (ADI); September 2016. Available from: <https://www.alz.co.uk/research/WorldAlzheimerReport2016.pdf>.
6. Brinda EM, Rajkumar AP, Enemark U, Attermann J, Jacob KS. Cost and burden of informal caregiving of dependent older people in a rural Indian community. *BMC health services research*. 2014 Dec;14(1):207.
7. Bürge E, Berchtold A, Maupetit C, Bourquin NM, von Gunten A, Ducraux D, Zumbach S, Peeters A, Kuhne N. Does physical exercise improve ADL capacities in people over 65 years with moderate or severe dementia hospitalized in an acute psychiatric setting? A multisite randomized clinical trial. *International psychogeriatrics*. 2017 Feb;29(2):323-32.
8. Cheng ST. Dementia caregiver burden: a research update and critical analysis. *Current psychiatry reports*. 2017 Sep 1;19(9):64.
9. Dawood S. Caregiver burden, quality of life and vulnerability towards psychopathology in caregivers of patients with dementia/Alzheimer's disease. *J. Coll. Physicians Surg. Pak*. 2016 Nov 1;26:892-5.
10. Dias A, Patel V. Closing the treatment gap for dementia in India. *Indian journal of psychiatry*. 2009 Jan;51(Suppl1):S93.
11. Fauth EB, Gibbons A. Which behavioral and psychological symptoms of dementia are the most problematic? Variability by prevalence, intensity, distress ratings, and associations with caregiver depressive symptoms. *International journal of geriatric psychiatry*. 2014 Mar 1;29(3):263-71.
12. Ferri CP, Jacob KS. Dementia in low-income and middle-income countries: Different realities mandate tailored solutions. *PLoS Medicine*. 2017 Mar 28;14(3):e1002271.
13. García-Alberca JM. Cognitive-behavioral treatment for depressed patients with Alzheimer's disease. An open trial. *Archives of gerontology and geriatrics*. 2017 Jul 1;71:1-8.
14. Karantzoulis S, Galvin JE. Distinguishing Alzheimer's disease from other major forms of dementia. *Expert review of neurotherapeutics*. 2011 Nov 1.
15. Khan Q. Dementia: Challenges of practice in Pakistan. *Neurology*. 2014 Nov 25;83(22):2091-2.
16. Khan QU, Sanober A. "Jinn Possession" and Delirious Mania in a Pakistani Woman. *American Journal of Psychiatry*. 2016 Feb 19;173(3):219-20.
17. Maestre GE. Assessing dementia in resource-poor regions. *Current neurology and neuroscience reports*. 2012 Oct 1;12(5):511-9.
18. Mahoney DF, Clutterbuck J, Neary S, Zhan L. African American, Chinese, and Latino family caregivers' impressions of the onset and diagnosis of dementia: cross-cultural similarities and differences. *The Gerontologist*. 2005 Dec 1;45(6):783-92.
19. Narayan SM, Varghese M, Hepburn K, Lewis M, Paul I, Bhimani R. Caregiving experiences of family members of persons with dementia in south India. *American Journal of Alzheimer's Disease & Other Dementias®*. 2015 Aug;30(5):508-16.
20. Ng QX, Kuah TZ, Loo GJ, Ho WH, Wagner



- NL, Sng JG, Yang GM, Tai BC. Awareness and Attitudes of Community-Dwelling Individuals in Singapore towards Participating in Advance Care Planning. *Annals of the Academy of Medicine, Singapore*. 2017 Mar 1;46(3):84-90.
21. Orgeta V, Qazi A, Spector A, Orrell M. Psychological treatments for depression and anxiety in dementia and mild cognitive impairment: systematic review and meta-analysis. *The British Journal of Psychiatry*. 2015 Oct 1;207(4):293-8.
  22. Prince MJ. World Alzheimer Report 2015: the global impact of dementia: an analysis of prevalence, incidence, cost and trends. *Alzheimer's Disease International*; 2015.
  23. Prince M, Comas-Herrera A, Knapp M, Guerchet M, Karagiannidou M. World Alzheimer report 2016: improving healthcare for people living with dementia: coverage, quality and costs now and in the future.
  24. Qadir F, Gulzar W, Haqqani S, Khalid A. A pilot study examining the awareness, attitude, and burden of informal caregivers of patients with dementia. *Care Management Journals*. 2013 Dec 1;14(4):230-40.
  25. Rehman A, Jingdong L, Hussain I. The province-wise literacy rate in Pakistan and its impact on the economy. *Pacific Science Review B: Humanities and Social Sciences*. 2015 Nov 1;1(3):140-4.
  26. Ruthirakuhan M, Luedke AC, Tam A, Goel A, Kurji A, Garcia A. Use of physical and intellectual activities and socialization in the management of cognitive decline of aging and in dementia: a review. *Journal of aging research*. 2012;2012.
  27. Vertejee S, Karamali NN. Active ageing in Pakistan: challenges and opportunities. *JPMA: Journal of Pakistan Medical Association*. 2014;64(1):P-76.

Conflict of interest: Author declares no conflict of interest.  
Funding disclosure: Nil

**Author's contribution:**

Dr Qurat ul Ain Khan was involved in formulating study concept and design including designing the survey questionnaire.

Dr Yusra Hanif Khan was involved in acquisition of subjects.

Mohammad Zaman Khan performed data analysis.

Dr Shireen Najam helped in preparation of manuscript.