



THE AGA KHAN UNIVERSITY

eCommons@AKU

Internal Medicine, East Africa

Medical College, East Africa

7-2021

Mental Disorders Among Health Care Workers at the Early Phase of COVID-19 Pandemic in Kenya; Findings of an Online Descriptive Survey

Edith Kamaru Kwobah

Ann Mwangi

Kirtika Patel

Thomas Mwogi

Robert Kiptoo

See next page for additional authors

Follow this and additional works at: https://ecommons.aku.edu/eastafrica_fhs_mc_intern_med



Part of the [Psychiatry Commons](#)

Authors

Edith Kamaru Kwobah, Ann Mwangi, Kirtika Patel, Thomas Mwogi, Robert Kiptoo, and Lukoye Atwoli



Mental Disorders Among Health Care Workers at the Early Phase of COVID-19 Pandemic in Kenya; Findings of an Online Descriptive Survey

Edith Kamaru Kwobah^{1*}, Ann Mwangi², Kirtika Patel³, Thomas Mwogi⁴, Robert Kiptoo¹ and Lukoye Atwoli^{5,6}

¹ Department of Mental Health, Moi Teaching and Referral Hospital, Eldoret, Kenya, ² Department of Mathematics, Physics and Computing, Moi University, Eldoret, Kenya, ³ Department of Immunology, Moi University, Eldoret, Kenya, ⁴ Directorate of Informatics, Moi Teaching and Referral Hospital, Eldoret, Kenya, ⁵ Department of Medicine, Aga Khan University Medical College, East Africa, Nairobi, Kenya, ⁶ Department of Mental Health and Behavioral Sciences, Moi University, Eldoret, Kenya

OPEN ACCESS

Edited by:

Eric Hahn,
Charité—Universitätsmedizin
Berlin, Germany

Reviewed by:

Zeinab Abbas,
Lebanese International
University, Lebanon
Agata Benfante,
University of Turin, Italy

*Correspondence:

Edith Kamaru Kwobah
eckamaru@gmail.com

Specialty section:

This article was submitted to
Public Mental Health,
a section of the journal
Frontiers in Psychiatry

Received: 08 February 2021

Accepted: 24 June 2021

Published: 22 July 2021

Citation:

Kwobah EK, Mwangi A, Patel K, Mwogi T, Kiptoo R and Atwoli L (2021) Mental Disorders Among Health Care Workers at the Early Phase of COVID-19 Pandemic in Kenya; Findings of an Online Descriptive Survey. *Front. Psychiatry* 12:665611. doi: 10.3389/fpsy.2021.665611

Background: Healthcare workers responding to the Corona Virus Pandemic (COVID-19) are at risk of mental illness. Data is scanty on the burden of mental disorders among Kenyan healthcare workers responding to the pandemic that can inform mental health and psychosocial support. The purpose of this study was to establish the frequency and associated factors of worry, generalized anxiety disorder, depression, posttraumatic stress disorder and poor quality of sleep among Kenyan health care workers at the beginning of COVID-19 pandemic.

Methods: We conducted an online survey among 1,259 health care workers in Kenya. A researcher developed social demographic questionnaire and several standardized tools were used for data collection. Standardized tools were programmed into Redcap, (Research Electronic Data Capture) and data analysis was performed using R Core Team. In all analysis a p -value < 0.05 was considered significant.

Results: 66% of the participants reported experiencing worry related to COVID-19. 32.1% had depression, 36% had generalized anxiety, 24.2% had insomnia and 64.7% scored positively for probable Post Traumatic Stress Disorder (PTSD). Depression was higher among females compared to men (36.5 vs. 26.9%, $p = 0.003$), workers < 35 years old compared to older ones (38.1 vs. 26.4%, $p < 0.001$), and those who were not married compared to those who were married (40.6 vs. 27.6%, $p < 0.001$). Generalized anxiety was commoner among workers aged < 35 years (43.5 vs. 29.3%, $p < 0.001$), females (41.7 vs. 29.2%, $p < 0.001$), those who were not married compared to the married (45.2 vs. 31.2%, $p < 0.001$) and those with < 10 years working experience (41.6 vs. 20.5%, $p < 0.001$). Younger health care professional had a higher proportion of insomnia compared to the older ones (30.3 vs. 18.6%, $p < 0.001$). Insomnia was higher among those with < 10 years' experience compared to those with more than 20 years' experience (27.3 vs. 17.6%, $p = 0.043$).

Conclusion: Many Kenyan healthcare workers in the early phase of COVID-19 pandemic suffered from various common mental disorders with young, female professionals who are not married bearing the bigger burden. This data is useful in informing interventions to promote mental and psychosocial wellbeing among Kenyan healthcare workers responding to the pandemic.

Keywords: prevalence, health care workers, Kenya, mental disorders, COVID-19

BACKGROUND

The mental health impact of the Corona Virus disease of 2019 (COVID-19) is expected to be huge due to the nature of the disease and the containment measures imposed (1). There is sufficient evidence to indicate that during any epidemic or pandemic, healthcare workers are affected mentally to a great extent (2). Unique to the case of COVID-19 pandemic in the early phase was the widespread media coverage, high rates of human to human transmission, and the fact that a lot is still unknown about this new virus, which may all contribute to poor mental health among these health care professionals. The ever-increasing number of confirmed and suspected cases, overwhelming workload, depletion of personal protection equipment, lack of specific curative drugs, and healthcare workers feelings of being inadequately supported significantly contribute to the risk of their mental health illness (3).

An increase in mental and psychological distress during epidemics has been documented in literature. As an example, the health care workers who responded to the 2014-2015 Ebola epidemic in Africa reported high levels of psychological symptoms (4). Similarly, during the outbreak of Severe Acute Respiratory Syndrome (SARS) in Hong Kong and Beijing, 20% of the total number of reported cases in the end were of frontline healthcare workers, who had to face a tremendous mental burden in addition to physical strain in taking care of patients with the highly contagious disease (5). Further, a previous study found that during the outbreak of infectious diseases, the affected hospitals experienced severe staff shortage as a result of personal or family health concerns, child care issues, quarantine measures or inability to get to work, and that health care workers were particularly worried for both their own and their family's health, and they experienced significant psychosocial trauma (6). In Singapore following the SARS outbreak, doctors had to be more vigilant when examining and reviewing patients, and it is reported that 20% of the doctors and nurses suffered PTSD -Post Traumatic Stress Disorder (7).

During a pandemic, healthcare workers are likely to express worry, besides fulfilling criteria for specific mental health disorders. A rapid review of literature done between March and April 2020 reported that health care workers working during the current pandemic reported frequent worry regarding their own health and the fear of infecting their families, friends and

colleagues (8). A web based survey that included 31 countries across the globe, including Kenya, conducted between April and May 2021 across the globe reported that 60% of health care workers had anxiety and 53% had mild to moderate depression (9). A review of the immediate impact of Covid-19 revealed that up to 35% of healthcare workers are likely to report symptoms of traumatic stress (10).

There has been a call across the globe to put in all possible measures to preserve and enhance resilience of healthcare workers (11) and for managers to proactively take steps to protect the mental well-being of staff (12). Such efforts need to be backed by evidence of the burden of the mental illness among these front liners, but unfortunately there is scanty data from the Kenyan context and other low resource settings to guide interventions to improve mental health and psychological support among healthcare workers. Such evidence would be useful not only for Kenya but for similar contexts in low and middle income countries where human resource for health is especially constrained, hence are likely to be stretched some more by the current pandemic, thus increasing their risk of mental ill health (13). This study therefore sought to 1. Determine the Frequency of worry, generalized anxiety disorder, depression, posttraumatic stress disorder (PTSD) and poor quality of sleep among health care professionals at the beginning of the COVID-19 pandemic- [these conditions were selected as they have been demonstrated as being highly prevalent among healthcare workers even before the pandemic (14, 15)], and 2. To establish the association between socio-demographic characteristics and mental disorders among health care professionals at the beginning of COVID-19 pandemic in Kenya.

MATERIALS AND METHODS

A cross-sectional descriptive online survey was used for this study. Trained healthcare professionals working in healthcare settings during the COVID-19 pandemic were included—these included, nurses, doctors, clinical officers and public health officers interacting with patients at the time of the study. Trained nurses, doctors, clinical officers and public health officers who do not interact with patients on day to day basis such as professionals working with insurance companies, those working in administration settings only or those who work exclusively in teaching institutions were excluded.

The survey questionnaires were programmed into Redcap, (Research Electronic Data Capture) a secure, web-based software platform designed to support data capture for research studies (16). A researcher developed social demographic questionnaire

Abbreviations: AOR, adjusted odds ratio; COVID-19, corona virus disease of 2019; DSMV, diagnostic statistical manual version V; GAD, generalized anxiety disorder; PHQ, patient health questionnaire; PTSD, post-traumatic stress disorder.

was used to collect data on age, sex, cadre, place of work, role during the pandemic and years of experience. Standardized tools were used for data collection. For anxiety we used Generalized Anxiety Disorder 7 (GAD 7) which we scored 0–4 = no anxiety, 5–9 = mild anxiety, 10–14 = moderate anxiety and ≥ 15 = severe anxiety (17). GAD 7 has demonstrated good internal consistency and convergent validity in heterogeneous samples (18) and has been used within Kenyan settings (19, 20). For depression we used Patient Health Questionnaire (PHQ 9) for depression which we scored 0–5 = Mild depression, 6–10 = moderate depression, 11–15 = moderately severe and 16–20 = severe depression (21). PHQ 9 is a widely used tool for screening for depression with a good reliability and convergent validity (22) and has been used previously in Kenya, having demonstrated good content validity in a study in Western Kenya (23). For PTSD we used Primary Care- Post traumatic Stress disorder (PC- PTSD) for Diagnostic Statistical Manual (DSM) V—those who responded positively to 3 out of 5 questions about how the traumatic event(s) had affected them over the past month was regarded as probable PTSD (24). Though this has not been validated in a Kenyan context it has been shown to have high diagnostic accuracy and is highly acceptable due to its simplicity and brevity- having only five questions (25) and has shown good reliability even outside the US setting, as is the case of the Korean validation (26). We used the term Probable PTSD because to make a diagnosis of PTSD we would need a more comprehensive assessment by a clinician. For sleep difficulties we used Pittsburg Sleep Quality Index (PSQI), a self-rated questionnaire which assesses sleep quality and disturbances over a 1-month time interval, and used the following cut-offs; 0–7 = no clinically significant insomnia; 8–14 = subthreshold insomnia; 15–21 = clinical insomnia of moderate severity; 21–28 = severe clinical insomnia (27). Though not validated for use in Kenya it has been validated for other African settings such as Ethiopia (28) and Nigeria (29).

We used the Cochran formulae for sample size calculation for a survey, assuming a 50% prevalence of common mental disorders and a 95% confidence level: $\text{sample Size} = 1.96^2 \cdot [0.5 \times (1-0.5)] / [(0.05)^2]$. The minimum sample for each of the cadres was 384 (30). To allow for incomplete data, we planned to recruit at least 400 participants for each of the cadres.

A virtual snowball convenient sampling technique was utilized to recruit participants given. This was necessitated by limitation of a clear database for healthcare workers' contacts which would allow randomization. The online survey was sent to different healthcare workers in various networks on Facebook, WhatsApp and emails. A weekly reminder requesting workers to participate was sent between April 27th and June 5th 2020. The healthcare workers were requested to respond to the survey and share with their colleagues while a track of responses was kept using the Redcap software until there were no new responses for a period of 2 weeks, at which point we closed the survey.

Descriptive statistics were used to summarize the socio demographic characteristics of the participants. Chi square test was used in the bivariate analysis to assess categorical factors associated with the various mental health disorders and significant variables at 0.20 were considered in the multivariable logistic regression analysis and presented as adjusted odds

ratios (AORs) with the corresponding 95% Confidence Intervals. The variables considered in the regression analysis were the various social demographics characteristics (age, gender, years of experience, cadre), having existing medical conditions, having contacts with a confirmed COVID-19 patient and the type of facility one was working in. Data analysis was performed using R Core Team (2013). In all analysis a p -value < 0.05 was considered significant.

RESULTS

Socio Demographic Characteristics

Though we initially hoped for 2,000 participants (400 per cadre for 5 cadres) only a total of 1,259 participants opened the survey after 6 weeks of sharing the link. A total of 69 did not consent to participate in the study. A further 233 consented but didn't respond to the various sections of the questionnaire including the socio demographic and were thus excluded from the analysis. Nine hundred fifty seven participants completed at least one or more components of the questionnaire, giving a response rate of 76%.

The median age of the participants was 35 years (IQR: 30–42). Majority 522 (54.5%) were females, married 619 (64.7%), non-specialist doctors 378 (39.5%), Christians 794 (83.0%) had an undergraduate level of education 433 (45.5) and had < 10 years of experience in the medical field 546 (57%). Majority were working in a public facility 672 (70.2%) and most were in level 6 facility 306 (32%). Most of the participants reported that they did not have a medical condition 739 (77.2%) and also that they had never been treated for a mental illness 739 (77.2%) (Table 1).

Frequency and Factors Associated With Mental Disorders Worry

Overall, 66% of the participants were "Quite a bit" or "Very worried" about at least one of the various issues related to COVID-19. Participants reported being "Quite a bit" or "Very worried" about: contracting COVID-19 (71%) being hospitalized for COVID-19 (58%), dying of COVID-19 (43%), losing a loved one due to COVID-19 (66%), being rejected due to COVID-19 (42%), infecting others with COVID-19 (65%) and not being able to do what they know best (61%). At least a third of the HCWs were worried about each of the COVID-19 related issues, including contacting the disease, dying and infecting others.

Being worried about COVID-19 was less likely to be reported among participants who had more years of experience compared to those who had less year (64 vs. 69%, $p = 0.026$), specialist doctors compared to other cadres (56 vs. 73% $p = 0.006$) and those who were working in private facilities compared to working in public facilities (61 vs. 68 % $p = 0.024$). Those who had contact with COVID-19 patients had higher likelihood of reporting worry than those who had not had contact (71 vs. 64%, $p = 0.040$) (Table 2).

Being worried about COVID-19 was less likely to be reported among participants who had more years of experience compared to those who had less year (64 vs. 69%, $p = 0.026$), specialist doctors compared to other cadres (56 vs. 73% $p =$

TABLE 1 | Sociodemographic characteristics of the participants.

	(N = 957)
Age (yrs)	
Mean (SD)	37.39 (10.18)
Range	20.00 - 83.0
Gender	
Male	435 (45.5%)
Female	522 (54.5%)
Marital status	
Married	619 (64.7%)
Single	272 (28.4%)
Widowed/Separated/Divorced	47 (4.9%)
Other	19 (2.0%)
Religion	
Muslim	118 (12.3%)
Christian	794 (83.0%)
Other	45 (4.7%)
Cadre	
Doctor non-specialist	378 (39.5%)
Doctor specialist	156 (16.3%)
Nurse	186 (19.5%)
Other	236 (24.7%)
Education level	
Diploma	151 (15.8%)
Higher diploma	59 (6.2%)
Undergraduate	435 (45.5%)
Masters	270 (28.2%)
Phd/Fellowship	42 (4.4%)
Years of experience	
0–10	546 (57.0%)
11–20	239 (25.0%)
20+	108 (18.0%)
Facility	
Public	672 (70.2%)
Private	285 (29.8%)
Level of facility	
Level 6 National	306 (32.0%)
Level 5 county referral	273 (28.5%)
Level 4 Subcounty	217 (22.7%)
Level 3 Health Center	112 (11.7%)
Level 2 Dispensary	49 (5.1%)
Have a known medical condition	
Yes	218 (22.8%)
No	739 (77.2%)
Ever been treated for mental illness	
Yes	58 (6.1%)
No	899 (93.9%)
Currently working directly with COVID-19 clients	
Yes	231 (24.1%)
No	726 (75.9%)

0.006) and those who were working in private facilities compared to working in public facilities (61 vs. 68 % $p = 0.024$). Those who had contact with COVID-19 patients had higher likelihood of reporting worry than those who had not had contact (71 vs. 64%, $p = 0.040$) (Table 2).

Depression

One third of the participants 283/599 (32.1%) scored positively for depression, with a majority having moderate depression. On bivariate analysis, age, sex, marital status and years of experiences were significantly associated with depression. A higher proportion of female (36.5%) had moderate/sever symptoms of depression compared to males (26.9%), $p = 0.003$. In terms of age there was a higher proportion with depression among those <35 years (38.1%) compared to those above 35 years (26.4%), $p < 0.001$. Health care providers who were married reported lower rates of depression (27.6%) compared to those who were not married (40.6%), $p < 0.001$. While the proportion with depression decreased as the number of years of experience in the medical increased; 37% for those with <10 years and 20% for those with above 20 years of experience, $p < 0.001$ (Table 3).

Adjusting for age, years of experience, cadre and type of facility on a multivariate logistic regression, sex and marital status were statistically significantly associated with depression. The odds of moderate/severe depression was 1.5 times in females compared to males (AOR = 1.47; 95% CI: 1.09, 2.00, $p = 0.012$). The odds of depression in health workers who were not married was 1.5 times that of those who were married (AOR = 1.47; 95%CI: 1.07, 2.01, $p = 0.016$). HCWS with more years of experience were less likely to score positively for depression (AOR 0.50, 95% CI 0.28–0.88, $p = 0.016$) (Table 4).

Generalized Anxiety Disorder

About a third of the participants, 291/807 (36%) scored positively for generalized anxiety. Those who experienced sleeping difficulties were 24.2% (189/780). On bivariate analysis, age, sex, marital status, years of experience in medical profession and cadre were statistically significantly associated with generalized anxiety disorder. Younger health care profession reported a higher proportion with generalized anxiety (43.5%) compared to those aged more than 35 years (29.3%), $p < 0.001$. Females had a higher proportion with generalized anxiety compared to males (41.7 vs. 29.2%), $p < 0.001$, while married had a lower proportion compared to those who were not married (31.2 vs. 45.2%), $p < 0.001$. In terms of experience as the number of years of experience in the health profession increased from <10 years to more than 20 years the proportion with generalized anxiety decreased from 41.6 to 20.5% $p < 0.001$, and the Doctor specialist reported the lowest proportions of generalized anxiety (28.4%) compared to other workers; non-specialist doctors 36.8%, nurses 30.9, and other cadres 43.9 %, $p = 0.013$ (Table 3).

Adjusting for age, years of experience, cadre and having contact with COVID-19 patients' on a multivariate logistic regression analysis, factors associated with generalized anxiety were sex and, marital status and the type of facility were associated with generalized anxiety. Females had a higher odds of having anxiety compared to males (AOR = 1.72; 95%CI: 1.25,

TABLE 2 | Factors associated with Worry among Healthcare workers.

Variable	Levels	Worry (N = 957)		
		Not very much (n = 322)	Very much (n = 635)	P-value
Age in years	<35	135 (29.2)	327 (70.8)	0.262
	≥35	187 (37.8)	308 (62.2)	
Sex	Male	155 (35.6%)	280 (64.4%)	0.235
	Female	167 (32.0%)	355 (68.0%)	
Marital status	Married	214 (34.6%)	405 (65.4%)	0.828
	Not married	86 (31.6%)	186 (68.4%)	
Years of experience	0–10	165 (30.2%)	381 (69.8%)	0.026
	11–20	95 (39.7%)	144 (60.3%)	
	20+	62 (36.0%)	110 (64.0%)	
Cadre	Specialist	68 (43.6%)	88 (56.4%)	0.006
	Doctor	132 (34.9%)	246 (65.1%)	
	Nurse	59 (31.7%)	127 (68.3%)	
	Other	63 (26.7%)	173 (73.3%)	
Facility	Public	211 (31.4%)	461 (68.6%)	0.024
	Private	111 (38.9%)	174 (61.1%)	
Have known medical condition	Yes	67 (30.7%)	151 (69.3%)	0.300
	No	255 (34.5%)	484 (65.5%)	
Contact COVID-19 clients	Yes	65 (28.1%)	166 (71.9%)	0.040
	No	257 (35.4%)	469 (64.6%)	

2.37, $p = < 0.001$). Not being married and working in private facility increased the odds of having generalized anxiety (AOR = 1.41; 95%CI: 1.02, 1.96, $p = 0.036$), and (AOR = 1.57; 95%CI: 1.13, 2.20, $p = 0.008$), respectively. HCWS with more years of experience were less likely to score positively for GAD (AOR 0.49, 95% CI 0.27–0.97, $p = 0.021$) (Table 4).

Insomnia (Pittsburg Sleep Quality Index)

15/957 (1.9%) of the healthcare workers had clinically moderate insomnia while 174/957 (22.3%) had subthreshold insomnia. On bivariate analysis, age, cadre and years of experience in the health profession were statistically significantly associated with insomnia. Younger health care professional had a higher proportion with insomnia (30.3%) compared to 18.6% among those over 35 years old, $p < 0.001$. Proportion with Insomnia was higher among those with less than 10 years' experience (27.3%) compared to those with more than 20 years' experience in the health profession (17.6), $p = 0.043$. Other cadres had higher proportions of insomnia (30.7) compared to specialist doctors (17.5), $p = 0.022$ (Table 3).

Adjusting for sex, marital status, years of experience, cadre and having a known medical condition in a multivariate logistic regression, health professionals aged 35 years and above had a reduced odds of reporting insomnia compared to the younger ones (AOR = 0.49; 95%CI: 0.28, 0.83) (Table 4).

PTSD

For PTSD majority of the health workers didn't respond to the questions. Of the ones who responded 225/348 (64.7%) had some probable PTSD (Table 5). Due to the few number of people

responding to the trauma questions, we did not do further analysis on this item.

DISCUSSIONS

This study adds to the existing literature on the frequency of mental illness among health care workers responding to COVID-19 in Kenya. We found a substantial degree of worry related to COVID-19 among healthcare workers in the early phase of the pandemic. The frequency of depression, anxiety, insomnia and probable PTSD were high. Young female workers with less experience in the healthcare profession were more likely to score positively for mental illness.

Worry

In our study two thirds of the participants were "Quite a bit" or "Very worried" about at least one of the various issues related to COVID-19. The leading three issues participants worried about were losing a loved one to COVID-19, infecting others, and getting infected with COVID-19. While worry may not be classified as a mental illness, it is of importance as it could be a precursor of more severe mental illness if not addressed (31). These findings are comparable to those of a Japanese study that showed that majority (78%) of the healthcare workers were seriously worried about COVID-19 and being infected was one of the leading concerns (32). The high frequency of worry in our setting is in keeping with a recent review that indicated that other than the main diagnostic categories of mental illness, many healthcare workers have various concerns regarding COVID-19 that increase their risk of experiencing psychological distress (33).

TABLE 3 | Factors associated with mental disorders on bivariate analysis.

Variable	Levels	PHQ (N = 882)			GAD (N = 807)			PSQI (N = 780)		
		Mild	Moderate/ severe	P-value	Present	None	P-value	Insomnia	No insomnia	P-value
Age in years	<35	265 (61.9)	163 (38.1)	<0.001	167 (43.5)	217 (56.5)	<0.001	114 (30.3)	262 (69.7)	<0.001
	≥35	334 (73.6)	120 (26.4)		124 (29.3)	299 (70.7)		75 (18.6)	329 (81.4)	
Sex	Male	296 (73.1)	109 (26.9)	0.003	106 (29.2)	257 (70.8)	<0.001	73 (20.8)	278 (79.2)	0.052
	Female	303 (63.5)	174 (36.5)		185 (41.7)	259 (58.3)		116 (27.0)	313 (73.0)	
Marital status	Married	419 (72.4)	160 (27.6)	<0.001	165 (31.2)	363 (68.8)	<0.001	110 (21.9)	392 (78.1)	0.052
	Not married	180 (59.4)	123 (40.6)		126 (45.2)	153 (54.8)		79 (28.4)	199 (71.6)	
Years of experience	0-10	320 (62.9)	189 (37.1)	<0.001	191 (41.6)	268 (58.4)	<0.001	123 (27.3)	327 (72.7)	0.043
	11–20	155 (71.4)	62 (28.6)		70 (34.7)	132 (65.3)		42 (21.6)	152 (78.4)	
	20+	124 (79.5)	32 (20.5)		30 (20.5)	116 (79.5)		24 (17.6)	112 (82.4)	
Cadre	Specialist	106 (70.7)	44 (29.3)	0.454	40 (28.4)	101 (71.6)	0.013	24 (17.5)	113 (82.5)	0.022
	Doctor	241 (67.7)	115 (32.3)		116 (36.8)	199 (63.2)		80 (25.5)	234 (74.5)	
	Nurse	116 (70.7)	48 (29.3)		47 (30.9)	105 (69.1)		27 (19.4)	112 (80.6)	
	Other	135 (64.0)	76 (36.0)		87 (43.9)	111 (56.1)		58 (30.7)	131 (69.3)	
Facility	Public	426 (69.4)	188 (30.6)	0.182	190 (33.7)	373 (66.3)	0.046	133 (24.8)	404 (75.2)	0.668
	Private	173 (64.6)	95 (35.4)		101 (41.4)	143 (58.6)		56 (23.0)	187 (77.0)	
Have known medical condition	Yes	138 (69.7)	60 (30.3)	0.600	73 (39.0)	114 (61.0)	0.378	50 (28.6)	125 (71.4)	0.155
	No	461 (67.4)	223 (32.6)		218 (35.2)	402 (64.8)		139 (23.0)	466 (77.0)	
Contact COVID-19 clients	Yes	141 (68.1)	66 (31.9)	1.000	76 (39.8)	115 (60.2)	0.253	47 (25.1)	140 (74.9)	0.816
	No	458 (67.9)	217 (32.1)		215 (34.9)	401 (65.1)		142 (23.9)	451 (76.1)	

Participants who had more years of experience, were specialist doctors, or were working in private facilities had lower likelihood of reporting worry, while those who had had contact with COVID-19 patients had higher likelihood of reporting worry. Such worry has been associated with the fact that this disease is new, is highly contagious, has no cure and there is rapidly evolving information about its outcomes (34). This is also likely due to the several reports of healthcare workers contracting the disease and a good number succumbing to it in various part of the world (35).

Depression

Our study established high rates of depression among healthcare workers; 15.4% having moderately severe to severe depression

and 16% having moderate depression. Our findings differ from a similar study done by Onchonga et al. in the same setting which reported that 53% of healthcare workers had mild depression while 9.2% had severe depression (36). This difference may be attributed to the timing of data collection during the pandemic as different waves may present with different levels of psychological distress. These findings are higher than that of a similar study involving 150 healthcare workers from Nepal which reported that 5.3% had moderate depression while 2.7% had moderately severe to severe depression (37). A study done among 906 healthcare workers from Singapore and India reported that 5.3% had moderate to severe depression (38). Our findings are also higher than a recent systematic review and meta-analysis done during the COVID-19 season that

TABLE 4 | Multivariate logistic regression.

Variable	PTSD			GAD			PSQI		
	AOR	95% CI	P-value	AOR	95% CI	P-value	AOR	95% CI	P-value
Age in years	1			1			1		
<35									
≥35	0.83	0.53, 1.28	0.400	0.76	0.47, 1.20	0.200	0.49	0.28, 0.83	0.010
Sex	1			1			1		
Male									
Female	1.47	1.09, 2.00	0.012	1.72	1.25, 2.37	<0.001	1.39	0.98, 1.99	0.067
Marital status	1			1			1		
Married									
Not married	1.47	1.07, 2.01	0.016	1.41	1.02, 1.96	0.036	1.15	0.80, 1.64	0.500
Years of experience	1			1			1		
0–10									
11–20	0.75	0.46, 1.21	0.200	0.92	0.56, 1.52	0.700	1.27	0.70, 2.33	0.400
20+	0.50	0.28, 0.88	0.016	0.49	0.27, 0.90	0.021	1.12	0.55, 2.28	0.800
Cadre	1			1			1		
Specialist									
Doctor	0.70	0.43, 1.14	0.150	0.94	0.57, 1.56	0.800	1.20	0.68, 2.17	0.500
Nurse	0.85	0.51, 1.43	0.500	1.01	0.59, 1.74	0.900	0.92	0.49, 1.73	0.800
Other	1.04	0.64, 1.70	0.900	1.59	0.96, 2.63	0.072	1.72	0.98, 3.10	0.064
Facility	1			1					
Public									
Private	1.29	0.94, 1.78	0.120	1.57	1.13, 2.20	0.008	-		
Known Medical condition									
Yes							1		
No	-						0.66	0.45, 0.99	0.045
Contact COVID-19 clients									
Yes			1						
No			0.75	0.53, 1.07	0.110	-			

included 13 studies with total of 33,062 participants reported a pooled prevalence of 22.8% for depression, although it is not clear the cut offs for depression that were used (39). These differences may be attributed to the differences in the settings of these studies and the social demographic characteristics of the participants.

Several factors were associated with Depression in our study. A higher proportion of female had moderate/severe symptoms of depression compared to males. This is similar to findings of a nationwide study done in India among 433 healthcare workers that reported that women were twice likely to have depressive symptoms requiring treatment (40). This may be attributed to the fact that even in the general population, females are at a higher risk of depression owing to various psychological and

cultural factors (41). A study done in Chinese healthcare workers also found being male to be protective for depression (42). Health care providers who were married reported lower rates of depression compared to those who were not married and this is likely because marriage provides a support system that reduces the risk of depression. There was a higher proportion with depression among those <35 years compared to those above 35 years and the proportion with depression decreased as the number of years of experience in the medical field increased. This is in line with a study done among South African doctors that demonstrated that junior doctors are likely to experience burnout and depression and this could partly be attributed to the resilience that is built over years of practice resulting in less emotional exhaustion (43).

TABLE 5 | Frequency of various mental disorders among Kenyan Healthcare workers.

	Overall (N = 957)
Depression (PHQ)	
N-Miss	75
Mild	599 (67.9%)
Moderate	147 (16.7%)
Moderately severe	93 (10.5%)
Severe	43 (4.9%)
Generalized Anxiety Disorder	
N-Miss	150
None	516 (63.9%)
Mild	163 (20.2%)
Moderate	69 (8.6%)
Severe	59 (7.3%)
PC-PTSD (DSMV)	
N-Miss	609
None	123 (35.3%)
Probable PTSD	225 (64.7%)
Pittsburg Sleep Quality Index	
N-Miss	177
No Insomnia	591 (75.8%)
Clinically moderate Insomnia	15 (1.9%)
Subthreshold Insomnia	174 (22.3%)

Anxiety

In our study, a third of the participants (36%) had generalized anxiety. This is higher than the recent meta-analytic evidence which reported a pooled prevalence of 23% (39). These findings are comparable to those of the Nepal study that reported that 34% had anxiety but differed among those who had severe anxiety in Nepal where only 2% had moderately severe to severe anxiety (37).

Younger health care profession reported a higher proportion of generalized anxiety compared to those aged more than 35 years. This is in agreement with a Chinese study which also showed higher rates of anxiety among the younger healthcare workers compared to the older ones (44). Females had a higher proportion with generalized anxiety compared to males. This is in agreement with A Turkish study which indicated that females were at higher risks of generalized and health related anxiety compared to their male counterparts (45). These findings may be because, female gender has been associated with anxiety even in the general population. Those who were married had a lower proportion of anxiety compared to those who were not married. This may be explained by the social support that marriage brings to an individual. In terms of experience as the number of years of experience in the health profession increased from <10 years to more than 20 years the proportion with generalized anxiety decreased from and the doctor specialist reported the lowest proportion with generalized anxiety and these two may be associated with the confidence that comes with practice and managing more complex patients.

Sleeping Difficulties

Upto 24.2% of the healthcare workers reported sleeping difficulties. This is comparable to findings of a study done in China including both the health care workers and the general population which reported that of the 5641 respondents, 20% had clinically significant insomnia (46). It was also higher than that of an Ethiopian study that reported a prevalence of 12.4%, but lower than the pooled prevalence of insomnia among healthcare workers from a meta-analytic evidence which was reported to be 38.9% (39).

In our study, younger health care professionals had a higher proportion with insomnia compared to those over 35 years old. Proportion with Insomnia decreased with increased experience in the health profession and those in other cadres other than nurses and doctors had a higher proportion with insomnia. Sleep difficulties among healthcare workers during this pandemic are important as they may not only increase vulnerability to the virus but also affect productivity of the workers who need to be at their optimal performance as they combat the virus (47).

Probable PTSD

It is worth noting that many participants did not respond to the trauma question. The reason for this is not clear and would warrant further evaluation of the understanding of trauma concepts in the Kenyan setting. Of those who responded, two thirds had been exposed to potentially traumatic events that could result in posttraumatic stress disorder. This is slightly higher than findings of a study done among 863 medical care workers from seven provinces in China using the Impact of Event Scale-6 which reported a prevalence of 40.2% (48). Similar high rates of PTSD have been shown in previous Flu out breaks. For example a study done among nurses working in south Korea during the Middle East Respiratory Syndrome (MERS) reported that more than half of the develop PTSD (49). The higher rates may be explained by the high mortality rates associated with Covid-19 being published throughout the globe that increases the risk of vicarious trauma among the providers.

Relevance to Policy and Practice

This study provides an evidence base for the burden of mental illness among healthcare workers responding to COVID-19 pandemic in Kenya. The burden which is likely to worsen as the pandemic lingers on is likely to further compromise the healthcare systems which were already understaffed prior to the pandemic. This evidence put together is a call to prioritize Kenyan healthcare worker wellness by putting in all possible measures to preserve and enhance resilience of healthcare workers and to link to care those who are affected in order to sustain productivity and ensure continuity.

A global survey involving 32 countries already reported that healthcare workers are employing various coping strategies to cope such as positive thinking and getting family support (50). Government as well as institutional efforts to help healthcare workers implement these coping strategies in order to enable them continue working with

COVID-19 patients are highly recommended. These efforts include managers taking proactive steps to protect the mental well-being of staff, to be frank about the challenging situations, and to allow teams to come together often to discuss their experiences and check on each other's well-being as well as link to care those who actually develop mental illness.

STUDY LIMITATIONS

While these findings are novel, a few limitations must be born in interpreting these findings. First this was an online survey and may have response bias where by the non-responders might be more stressed health care professionals who did not respond to the online survey. Second Online surveys are less accessible to some groups of people (the older professionals especially) hence creating a bias. Third, cross-sectional data can identify associations, not causal relationships with COVID-19. Fourth we used convenient sampling hence the results may not be generalizable to other settings. Finally, there are several personal factors that could influence mental health of individuals that were not considered in this study. Nevertheless our study provides more evidence for the burden of mental ill health among healthcare workers that would inform efforts to promote mental well-ness among healthcare workers during the current pandemic and in future similar situations.

CONCLUSIONS

This study established that there is a high frequency of depression, anxiety, insomnia and probable PTSD among healthcare workers responding to COVID-19 pandemic in Kenya with the highest burden being among young, less experienced female professionals. This evidence is a call to all relevant authorities to put in place measures to enhance healthcare workers mental and psychosocial well-ness in order to enable healthcare workers participate optimally in the fight against COVID-19. Further, longitudinal studies are recommended to better understand the causal relationship between COVID-19 and mental ill health among healthcare workers in Kenya and similar settings.

REFERENCES

1. Semo B, Frissa SM. The Mental Health Impact of the COVID-19 Pandemic: Implications for Sub-Saharan Africa. *Psychol Res Behav Manag.* (2020) 13:713–20. doi: 10.2147/PRBM.S264286
2. Preti E, Di Mattei V, Perego G, Ferrari F, Mazzetti M, Taranto P, et al. The psychological impact of epidemic and pandemic outbreaks on healthcare workers: rapid review of the evidence. *Curr Psychiatry Rep.* (2020) 22:43. doi: 10.1007/s11920-020-01166-z
3. Lai J, Ma S, Wang Y, Cai Z, Hu J, Wei N, et al. Factors associated with mental health outcomes among health care workers exposed to coronavirus disease 2019. *JAMA Netw Open.* (2020) 3:e203976. doi: 10.1001/jamanetworkopen.2020.3976
4. Ji D, Ji Y-J, Duan X-Z, Li W-G, Sun Z-Q, Song X-A, et al. Prevalence of psychological symptoms among Ebola survivors and healthcare workers

DATA AVAILABILITY STATEMENT

The original contributions presented in the study are included in the article/**Supplementary Material**, further inquiries can be directed to the corresponding author/s.

ETHICS STATEMENT

The studies involving human participants were reviewed and approved by Moi University/MTRH Institutional Research and Ethics Committee. The ethics committee waived the requirement of written informed consent for participation.

AUTHOR CONTRIBUTIONS

KP and EK developed the initial concept. EK, LA, KP, TM, AM, and RK were involved in concept development, review of manuscript, and approval for final submission. AM did the statistical analysis. TM programmed the survey into Redcap and performed data cleaning. LA sourced for funding. All authors contributed to the article and approved the submitted version.

FUNDING

This work was supported by Kenya Medical Association, Equity project.

ACKNOWLEDGMENTS

The authors would like to acknowledge all Kenyan healthcare workers who took time to fill in the survey. A report on this work was shared with Kenya Medical Association. We acknowledge the coordination role played by Julius Barasa.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fpsy.2021.665611/full#supplementary-material>

during the 2014–2015 Ebola outbreak in Sierra Leone: a cross-sectional study. *Oncotarget.* (2017) 8:12784–91. doi: 10.18632/oncotarget.14498

5. Wong TW, Yau JKY, Chan CLW, Kwong RSY, Ho SMY, Lau CC, et al. The psychological impact of severe acute respiratory syndrome outbreak on healthcare workers in emergency departments and how they cope. *Eur J Emerg Med.* (2005) 12:13–8. doi: 10.1097/00063110-200502000-00005
6. Martinese F, Keijzers G, Grant S, Lind J. How would Australian hospital staff react to an avian influenza admission, or an influenza pandemic? *Emerg Med Australas.* (2009) 21:12–24. doi: 10.1111/j.1742-6723.2008.01143.x
7. Chan AOM, Chan YH. Psychological impact of the 2003 severe acute respiratory syndrome outbreak on health care workers in a medium size regional general hospital in Singapore. *Occup Med.* (2004) 54:190–6. doi: 10.1093/occmed/kqh027
8. Barello S, Falcó-Pegueroles A, Rosa D, Tolotti A, Graffigna G, Bonetti L. The psychosocial impact of flu influenza pandemics on healthcare workers

- and lessons learnt for the COVID-19 emergency: a rapid review. *Int J Public Health*. (2020) 1–12:1205–16. doi: 10.1007/s00038-020-01463-7
9. Htay MNN, Marzo RR, AlRifai A, Kamberi F, El-Abasiri RA, Nyamache JM, et al. Immediate impact of COVID-19 on mental health and its associated factors among healthcare workers: A global perspective across 31 countries. *J Glob Health*. (2020) 10:020381. doi: 10.7189/jogh.10.020381
 10. Benfante A, Di Tella M, Romeo A, Castelli L. Traumatic stress in healthcare workers during covid-19 pandemic: a review of the immediate impact. *Front Psychol*. (2020) 11. doi: 10.3389/fpsyg.2020.569935
 11. Santarone K, McKenney M, Elkbuli A. Preserving mental health and resilience in frontline healthcare workers during COVID-19. *Am J Emerg Med*. (2020) 38:1530–1. doi: 10.1016/j.ajem.2020.04.030
 12. Greenberg N, Docherty M, Gnanapragasam S, Wessely S. Managing mental health challenges faced by healthcare workers during covid-19 pandemic. *BMJ*. (2020) 368:m1211. doi: 10.1136/bmj.m1211
 13. WHO. *Priorities for research into human resources for health in low- and middle-income countries*. WHO. Available online at: <https://www.who.int/bulletin/volumes/88/6/09-066290/en/> (accessed March 31, 2021).
 14. Kim M-S, Kim T, Lee D, Yook J, Hong Y-C, Lee S-Y, et al. Mental disorders among workers in the healthcare industry: 2014 national health insurance data. *Ann Occup Environ Med*. (2018) 30:31. doi: 10.1186/s40557-018-0244-x
 15. Ruitenburg MM, Frings-Dresen MH, Sluiter JK. The prevalence of common mental disorders among hospital physicians and their association with self-reported work ability: a cross-sectional study. *BMC Health Serv Res*. (2012) 12:292. doi: 10.1186/1472-6963-12-292
 16. Software—REDCap. Available online at: <https://projectredcap.org/software/> (accessed November 16, 2020).
 17. Spitzer RL, Kroenke K, Williams JBW, Löwe B. A Brief Measure for Assessing Generalized Anxiety Disorder: The GAD-7. *Arch Intern Med*. (2006) 166:1092. doi: 10.1001/archinte.166.10.1092
 18. Johnson SU, Ulvenes PG, Øktedalen T, Hoffart A. Psychometric properties of the general anxiety disorder 7-Item (GAD-7) Scale in a heterogeneous psychiatric sample. *Front Psychol*. (2019) 10:1713. doi: 10.3389/fpsyg.2019.01713
 19. Nyongesa MK, Mwangi P, Koot HM, Cuijpers P, Newton CRJC, Abubakar A. The reliability, validity and factorial structure of the Swahili version of the 7-item generalized anxiety disorder scale (GAD-7) among adults living with HIV from Kilifi, Kenya. *Ann Gen Psychiatry*. (2020) 19:62. doi: 10.1186/s12991-020-00312-4
 20. Osborn TL, Venturo-Conerly KE, Wasil AR, Schleider JL, Weisz JR. Depression and anxiety symptoms, social support, and demographic factors among kenyan high school students. *J Child Fam Stud*. (2020) 29:1432–43. doi: 10.1007/s10826-019-01646-8
 21. Kroenke K, Spitzer RL, Williams JBW. The PHQ-9. *J Gen Intern Med*. (2001) 16:606–13. doi: 10.1046/j.1525-1497.2001.016009606.x
 22. Cameron IM, Crawford JR, Lawton K, Reid IC. Psychometric comparison of PHQ-9 and HADS for measuring depression severity in primary care. *Br J Gen Pract*. (2008) 58:32–6. doi: 10.3399/bjgp08X263794
 23. Monahan PO, Shacham E, Reece M, Kroenke K, Ong'or WO, Omollo O, et al. Validity/Reliability of PHQ-9 and PHQ-2 depression scales among adults living with HIV/AIDS in Western Kenya. *J Gen Intern Med*. (2009) 24:189–97. doi: 10.1007/s11606-008-0846-z
 24. Prins A, Bovin MJ, Smolenski DJ, Marx BP, Kimerling R, Jenkins-Guarnieri MA, et al. The Primary Care PTSD Screen for DSM-5 (PC-PTSD-5): Development and evaluation within a veteran primary care sample. *J Gen Intern Med*. (2016) 31:1206–11. doi: 10.1007/s11606-016-3703-5
 25. Bovin MJ, Kimerling R, Weathers FW, Prins A, Marx BP, Post EP, et al. Diagnostic Accuracy and Acceptability of the Primary Care Posttraumatic Stress Disorder Screen for the Diagnostic and Statistical Manual of Mental Disorders (Fifth Edition) among US veterans. *JAMA Netw Open*. (2021) 4:e2036733. doi: 10.1001/jamanetworkopen.2020.36733
 26. Jung Y-E, Kim D, Kim W-H, Roh D, Chae J-H, Park JE. A Brief Screening Tool for PTSD: Validation of the Korean Version of the Primary Care PTSD Screen for DSM-5 (K-PC-PTSD-5). *J Korean Med Sci*. (2018) 33:e338. doi: 10.3346/jkms.2018.33.e338
 27. Carpenter JS, Andrykowski MA. Psychometric evaluation of the pittsburgh sleep quality index. *J Psychosom Res*. (1998) 45:5–13. doi: 10.1016/S0022-3999(97)00298-5
 28. Salahuddin M, Maru TT, Kumalo A, Pandi-Perumal SR, Bahammam AS, Manzar MD. Validation of the pittsburgh sleep quality index in community dwelling ethiopian adults. *Health Qual Life Outcomes*. (2017) 15:58. doi: 10.1186/s12955-017-0637-5
 29. Aloba OO, Adewuya AO, Ola BA, Mapayi BM. Validity of the pittsburgh sleep quality index (psqi) among nigerian university students. *Sleep Med*. (2007) 5:266–70. doi: 10.1016/j.sleep.2006.08.003
 30. Cochran_1977_Sampling_Techniques_Third_Edition.pdf. Available online at: https://glad.geog.umd.edu/Potapov/Library/Cochran_1977_Sampling_Techniques_Third_Edition.pdf (accessed April 7, 2021).
 31. *Worry and its Psychological Disorders: Theory, Assessment and Treatment—Google Books*. Available online at: [https://books.google.co.ke/books?hl=en&lr=&id=a34y-oRuMOIC&oi=fnd&pg=PR5&dq=worry\\$+\\$and\\$+\\$mental\\$+\\$health&ots=we5DU6wq8u&sig=KLeCnBH5A70LXL6gN_Z5g4_WKOI&redir_esc=y#v=onepage&q=worry%20and%20mental%20health&f=false](https://books.google.co.ke/books?hl=en&lr=&id=a34y-oRuMOIC&oi=fnd&pg=PR5&dq=worry$+$and$+$mental$+$health&ots=we5DU6wq8u&sig=KLeCnBH5A70LXL6gN_Z5g4_WKOI&redir_esc=y#v=onepage&q=worry%20and%20mental%20health&f=false) (accessed April 12, 2021).
 32. Sahashi Y, Endo H, Sugimoto T, Nabeta T, Nishizaki K, Kikuchi A, et al. Worries and concerns among healthcare workers during the coronavirus 2019 pandemic: A web-based cross-sectional survey. *Humanit Soc Sci Commun*. (2021) 8:1–8. doi: 10.1057/s41599-021-00716-x
 33. Spoorthy MS, Pratapa SK, Mahant S. Mental health problems faced by healthcare workers due to the COVID-19 pandemic—a review. *Asian J Psychiatry*. (2020) 51:102119. doi: 10.1016/j.ajp.2020.102119
 34. Onigbinde OA, Babatunde O, Ajagbe AO. The Welfare of Healthcare Workers amidst COVID-19 pandemic in Sub-Sahara Africa: A call for concern. *Ethics Med Public Health*. (2020) 15:100555. doi: 10.1016/j.jemep.2020.100555
 35. Neto MLR, Almeida HG, Esmeraldo JD, Nobre CB, Pinheiro WR, de Oliveira CRT, et al. When health professionals look death in the eye: the mental health of professionals who deal daily with the 2019 coronavirus outbreak. *Psychiatry Res*. (2020) 288:112972. doi: 10.1016/j.psychres.2020.112972
 36. Onchonga D, Ngetich E, Makunda W, Wainaina P, Wangeshi D, Viktoria P. Anxiety and depression due to 2019 SARS-CoV-2 among frontier healthcare workers in Kenya. *Heliyon*. (2021) 7:e06351. doi: 10.1016/j.heliyon.2021.e06351
 37. Gupta AK, Mehra A, Niraula A, Kafle K, Deo SP, Singh B, et al. Prevalence of anxiety and depression among the healthcare workers in Nepal during the COVID-19 pandemic. *Asian J Psychiatry*. (2020) 54:102260. doi: 10.1016/j.ajp.2020.102260
 38. Chew NWS, Lee GKH, Tan BYQ, Jing M, Goh Y, Ngiam NJH, et al. A multinational, multicentre study on the psychological outcomes and associated physical symptoms amongst healthcare workers during COVID-19 outbreak. *Brain Behav Immun*. (2020) 88:559–65. doi: 10.1016/j.bbi.2020.04.049
 39. Pappa S, Ntella V, Giannakas T, Giannakoulis VG, Papoutsis E, Katsaounou P. Prevalence of depression, anxiety, and insomnia among healthcare workers during the COVID-19 pandemic: a systematic review and meta-analysis. *Brain Behav Immun*. (2020) 88:901–7. doi: 10.1016/j.bbi.2020.05.026
 40. Wilson W, Raj JP, Rao S, Ghiya M, Nedungalaparambil NM, Mundra H, et al. Prevalence and predictors of stress, anxiety, and depression among healthcare workers managing COVID-19 pandemic in India: a nationwide observational study. *Indian J Psychol Med*. (2020) 42:353–8. doi: 10.1177/0253717620933992
 41. Albert PR. Why is depression more prevalent in women? *J Psychiatry Neurosci*. (2015) 40:219–21. doi: 10.1503/jpn.150205
 42. Zhu J, Sun L, Zhang L, Wang H, Fan A, Yang B, et al. Prevalence and Influencing Factors of Anxiety and Depression Symptoms in the First-Line Medical Staff Fighting Against COVID-19 in Gansu. *Front Psychiatry*. (2020) 11:386. doi: 10.3389/fpsyg.2020.00386
 43. Rossouw L, Seedat S, Emsley R, Suliman S, Hagemester D. The prevalence of burnout and depression in medical doctors working in the Cape Town Metropolitan Municipality community healthcare clinics and district hospitals of the Provincial Government of the Western Cape: a cross-sectional study. *South Afr Fam Pract*. (2013) 55:567–73. doi: 10.1080/20786204.2013.10874418
 44. Huang Y, Zhao N. Generalized anxiety disorder, depressive symptoms and sleep quality during COVID-19 outbreak in China: a web-based cross-sectional survey. *Psychiatry Res*. (2020) 288:112954. doi: 10.1016/j.psychres.2020.112954

45. Özdin S, Bayrak Özdin S. Levels and predictors of anxiety, depression and health anxiety during COVID-19 pandemic in Turkish society: The importance of gender. *Int J Soc Psychiatry*. (2020) 66:504–11. doi: 10.1177/0020764020927051
46. Morin CM, Carrier J. The acute effects of the COVID-19 pandemic on insomnia and psychological symptoms. *Sleep Med*. (2020) 77:346–7. doi: 10.1016/j.sleep.2020.06.005
47. Ferini-Strambi L, Zucconi M, Casoni F, Salsone M. Covid-19 and sleep in medical staff: reflections, clinical evidences, and perspectives. *Curr Treat Options Neurol*. (2020) 22:29. doi: 10.1007/s11940-020-00642-4
48. Si M-Y, Su X-Y, Jiang Y, Wang W-J, Gu X-F, Ma L, et al. Psychological impact of COVID-19 on medical care workers in China. *Infect Dis Poverty*. (2020) 9:113. doi: 10.1186/s40249-020-00724-0
49. Jung H, Jung SY, Lee MH, Kim MS. Assessing the presence of post-traumatic stress and turnover intention among nurses post-middle east respiratory syndrome outbreak: the importance of supervisor support. *Workplace Health Saf*. (2020) 68:337–45. doi: 10.1177/2165079919897693
50. Htay MNN, Marzo RR, Bahari R, AlRifai A, Kamberi F, El-Abasiri RA, et al. How healthcare workers are coping with mental health challenges during COVID-19 pandemic?—a cross-sectional multi-countries study. *Clin Epidemiol Glob Health*. (2021) 11:100759. doi: 10.1016/j.cegh.2021.100759

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2021 Kwobah, Mwangi, Patel, Mwogi, Kiptoo and Atwoli. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.