



THE AGA KHAN UNIVERSITY

eCommons@AKU

Department of Radiology

Medical College, Pakistan

January 2015

Pseudo-thyroid lobe: A diagnostic conundrum caused by ossified anterior longitudinal ligament on bone scan

Maseeh Uz Zaman

Aga Khan University, maseeh.uzzaman@aku.edu

Nosheen Fatima

Aga Khan University, nosheen.fatima@aku.edu

Zafar Sajjad

Unaiza Zaman

Areeba Zaman

See next page for additional authors

Follow this and additional works at: https://ecommons.aku.edu/pakistan_fhs_mc_radiol



Part of the [Radiology Commons](#)

Recommended Citation

Zaman, M. U., Fatima, N., Sajjad, Z., Zaman, U., Zaman, A., Tahseen, R. (2015). Pseudo-thyroid lobe: A diagnostic conundrum caused by ossified anterior longitudinal ligament on bone scan. *Indian Journal of Nuclear Medicine*, 30(1), 78-79.

Available at: https://ecommons.aku.edu/pakistan_fhs_mc_radiol/247

Authors

Maseeh Uz Zaman, Nosheen Fatima, Zafar Sajjad, Unaiza Zaman, Areeba Zaman, and Rabia Tahseen

“Pseudo-thyroid lobe”: A diagnostic conundrum caused by ossified anterior longitudinal ligament on bone scan

Maseeh Uz Zaman, Nosheen Fatima¹, Zafar Sajjad, Unaiza Zaman², Areeba Zaman², Rabia Tahseen²

Department of Radiology, Aga Khan University Hospital, ¹Department of Nuclear Medicine, Dr. Ziauddin Hospital, ²Dow University of Health Sciences, Karachi, Pakistan

ABSTRACT Radionuclide bone imaging is one of the most commonly performed nuclear medicine procedure around the world and characterized by its high sensitivity and relatively low specificity. False positive findings on a bone scan are very common; however, dense uptake over unilateral ossified anterior longitudinal ligament appearing as single thyroid lobe on a bone scan has not been described in the literature.

Keywords: Bone scan, ossified anterior longitudinal ligament, pseudo-thyroid, thyroid uptake

A 75-year-old male, clinically asymptomatic diagnosed case of carcinoma prostate since 2000 has had a radionuclide bone scan with 740 MBq of ^{99m}Tc-methylene diphosphonate (^{99m}Tc-MDP) for disease surveillance. Delayed spot images showed an elongated area of abnormal tracer uptake over right thyroid bed mimicking a thyroid lobe [Figure 1]. There was also evidence of degenerative changes over right shoulder, mid-cervical region on the left side and focal uptakes over right body of mandible and upper jaw due to associated dental infections. No gastric or salivary glands tracer uptake suggestive of free pertechnetate due to inadequate labeling was noted. Subsequently, a computed tomography scan was performed which revealed ossification of right anterior longitudinal ligament (OALL) involving (C4-T1) with fractures and associated pseudo-articulations [Figure 2]. There was no evidence of ossification of posterior longitudinal ligament (OPLL) and spinal stenosis. OALL is scarcely described in the literature as it is rarely symptomatic and dysphagia being the most common complication.^[1,2] OPLL with radiculopathy is more common entity than OALL.^[3]

OALL is most commonly seen in the thoracic spine followed by cervical and lumbar regions. In this case, it was found on the right side which is postulated to be due to the protective effect of the pulsatile aorta on the left of the thoracic spine.^[4] Uptake of ^{99m}Tc-MDP by the OALL over lower cervical and upper dorsal spine could create a diagnostic conundrum and must be included in the differential diagnosis of abnormal radiotracer uptake over thyroid bed in a bone scan.




Figure 1: Three hours delayed anterior bone scan images (a: Head straight; b and c: Head rotated to left and right, respectively) showing an elongated area of increased tracer uptake over right cervical region mimicking right thyroid lobe (pseudo-thyroid) with focal uptakes over jaws secondary to dental infection and arthritic changes over right shoulder

Access this article online

Quick Response Code:



Website:
www.ijnm.in

DOI:
10.4103/0972-3919.147554

Address for correspondence:

Dr. Maseeh Uz Zaman, Department of Radiology, Aga Khan University Hospital, Karachi, Pakistan.
E-mail: maseeh.uzzaman@aku.edu




Figure 2: (a) Sagittal, (b) coronal, and (c) axial nonenhanced computed tomography scan images showing right-sided ossified anterior longitudinal ligament extending from C4 to T1. There are also fractures and evidence of pseudo-articulation (arrows). Note the relationship to the thyroid gland and cartilage that lies immediately anterior to the bulky osteophytes

REFERENCES

1. McCafferty RR, Harrison MJ, Tamas LB, Larkins MV. Ossification of the anterior longitudinal ligament and Forestier's disease: An analysis of seven cases. *J Neurosurg* 1995;83:13-7.
2. Epstein NE, Hollingsworth R. Ossification of the cervical anterior longitudinal ligament contributing to dysphagia. Case report. *J Neurosurg* 1999;90:261-3.
3. Choi BW, Song KJ, Chang H. Ossification of the posterior longitudinal ligament: A review of literature. *Asian Spine J* 2011;5:267-76.
4. Resnick D, Shapiro RF, Wiesner KB, Niwayama G, Utsinger PD, Shaul SR. Diffuse idiopathic skeletal hyperostosis (DISH) [ankylosing hyperostosis of Forestier and Rotes-Querol]. *Semin Arthritis Rheum* 1978;7:153-87.

How to cite this article: Zaman MU, Fatima N, Sajjad Z, Zaman U, Zaman A, Tahseen R. "Pseudo-thyroid lobe": A diagnostic conundrum caused by ossified anterior longitudinal ligament on bone scan. *Indian J Nucl Med* 2015;30:78-9.

Source of Support: Nil. **Conflict of Interest:** None declared.

