



THE AGA KHAN UNIVERSITY

eCommons@AKU

---

Community Health Sciences

Department of Community Health Sciences

---

January 2017

# Effects of bidi smoking on all-cause mortality and cardiorespiratory outcomes in men from south Asia: an observational community-based substudy of the Prospective Urban Rural Epidemiology Study (PURE)

MyLinh Duong

*McMaster University and Hamilton Health Sciences, Hamilton, ON, Canada*

Sumathy Rangarajan

*McMaster University and Hamilton Health Sciences, Hamilton, ON, Canada*

Xiaohe Zhang

*McMaster University and Hamilton Health Sciences, Hamilton, ON, Canada*

Kieran Killian

*McMaster University and Hamilton Health Sciences, Hamilton, ON, Canada*

Prem Mony

*St John's Research Institute, Bengaluru, India*

---

## Recommended Citation

Duong, M., Rangarajan, S., Zhang, X., Killian, K., Mony, P., Swaminathan, S., Bharathi, A. V., Nair, S., Vijayakumar, K., Mohan, I., Iqbal, R., Kazmi, K. (2017). Effects of bidi smoking on all-cause mortality and cardiorespiratory outcomes in men from south Asia: an observational community-based substudy of the Prospective Urban Rural Epidemiology Study (PURE). *Lancet Global Health*, 5(2), 168-176.

Available at: [https://ecommons.aku.edu/pakistan\\_fhs\\_mc\\_chs\\_chs/351](https://ecommons.aku.edu/pakistan_fhs_mc_chs_chs/351)

*See next page for additional authors*

Follow this and additional works at: [https://ecommons.aku.edu/pakistan\\_fhs\\_mc\\_chs\\_chs](https://ecommons.aku.edu/pakistan_fhs_mc_chs_chs)

 Part of the [Cardiology Commons](#), and the [Public Health Commons](#)

---

---

**Authors**

MyLinh Duong, Sumathy Rangarajan, Xiaohe Zhang, Kieran Killian, Prem Mony, Sumathi Swaminathan, Ankalmadagu Venkatsubbareddy Bharathi, Sanjeev Nair, Krishnapillai Vijayakumar, Indu Mohan, Romaina Iqbal, and Khawar Kazmi



# Effects of bidi smoking on all-cause mortality and cardiorespiratory outcomes in men from south Asia: an observational community-based substudy of the Prospective Urban Rural Epidemiology Study (PURE)

MyLinh Duong, Sumathy Rangarajan, Xiaohe Zhang, Kieran Killian, Prem Mony, Sumathi Swaminathan, Ankalmadagu Venkatsubbareddy Bharathi, Sanjeev Nair, Krishnapillai Vijayakumar, Indu Mohan, Rajeev Gupta, Deepa Mohan, Shanthi Rani, Viswanathan Mohan, Romaina Iqbal, Khawar Kazmi, Omar Rahman, Rita Yusuf, Lakshmi Venkata Maha Pinnaka, Rajesh Kumar, Paul O'Byrne, Salim Yusuf



## Summary

**Background** Bidis are minimally regulated, inexpensive, hand-rolled tobacco products smoked in south Asia. We examined the effects of bidi smoking on baseline respiratory impairment, and prospectively collected data for all-cause mortality and cardiorespiratory events in men from this region.

**Methods** This substudy of the international, community-based Prospective Urban Rural Epidemiology (PURE) study was done in seven centres in India, Pakistan, and Bangladesh. Men aged 35–70 years completed spirometry testing and standardised questionnaires at baseline and were followed up yearly. We used multilevel regression to compare cross-sectional baseline cardiorespiratory symptoms, spirometry measurements, and follow-up events (all-cause mortality, cardiovascular events, respiratory events) adjusted for socioeconomic status and baseline risk factors between non-smokers, light smokers of bidis or cigarettes ( $\leq 10$  pack-years), heavy smokers of cigarettes only ( $>10$  pack-years), and heavy smokers of bidis ( $>10$  pack-years).

**Findings** 14 919 men from 158 communities were included in this substudy (8438 non-smokers, 3321 light smokers, 959 heavy cigarette smokers, and 2201 heavy bidi smokers). Mean duration of follow-up was 5.6 years (range 1–13). The adjusted prevalence of self-reported chronic wheeze, cough or sputum, dyspnoea, and chest pain at baseline increased across the categories of non-smokers, light smokers, heavy cigarette smokers, and heavy bidi smokers ( $p < 0.0001$  for association). Adjusted cross-sectional age-related changes in forced expiratory volume in 1 s (FEV<sub>1</sub>) and FEV<sub>1</sub>/forced vital capacity (FVC) ratio were larger for heavy bidi smokers than for the other smoking categories. Hazard ratios (relative to non-smokers) showed increasing hazards for all-cause mortality (light smokers 1.28 [95% CI 1.02–1.62], heavy cigarette smokers 1.59 [1.13–2.24], heavy bidi smokers 1.56 [1.22–1.98]), cardiovascular events (1.45 [1.13–1.84], 1.47 [1.05–2.06], 1.55 [1.17–2.06], respectively) and respiratory events (1.30 [0.91–1.85], 1.21 [0.70–2.07], 1.73 [1.23–2.45], respectively) across the smoking categories.

**Interpretation** Bidi smoking is associated with severe baseline respiratory impairment, all-cause mortality, and cardiorespiratory outcomes. Stricter controls and regulation of bidis are needed to reduce the tobacco-related disease burden in south Asia.

**Funding** Population Health Research Institute, Canadian Institutes of Health Research, and Heart and Stroke Foundation of Ontario.

**Copyright** © The Author(s). Published by Elsevier Ltd. This is an Open Access article under the CC BY-NC-ND license.

## Introduction

Tobacco use is the world's leading preventable cause of premature death.<sup>1</sup> The harmful effects of cigarette smoking have been extensively studied and are universally accepted. Less is known about the health effects of non-cigarette tobacco products, such as bidis (also known as beedis), which are commonly used in populations of low socioeconomic status.

Bidis are inexpensive, small, hand-rolled tobacco products commonly smoked in south Asia. Estimates suggest that there were 53 million users of bidis in India

and Bangladesh in 2003–09.<sup>2</sup> Outside this region, bidi use has mainly been reported in young adults in developed countries,<sup>3</sup> where unregulated marketing over the internet and in ethnic stores has allowed easy access to this vulnerable population. Bidis are manufactured in south Asia by a cottage industry that has avoided many of the local and international tobacco regulations and taxes enforced on factory-made cigarettes. Consequently, bidis are sold cheaply, at various prices and in packaging with poorly visible health warnings.<sup>4–6</sup> Furthermore, herbal and flavoured varieties are made to appeal to young adults as

*Lancet Glob Health* 2017;  
5: e168–76

Population Health Research Institute (M Duong MBBS, S Rangarajan MSc, X Zhang MSc, Prof S Yusuf DPhil) and Department of Medicine (M Duong, Prof K Killian PhD, Prof P O'Byrne MB, Prof S Yusuf), McMaster University and Hamilton Health Sciences, Hamilton, ON, Canada; Division of Epidemiology and Population Health, St John's Research Institute, Bengaluru, India (P Mony MD, S Swaminathan PhD, A V Bharathi PhD); Department of Pulmonary Medicine, Medical College, Thiruvananthapuram, Kerala, India (Prof S Nair MD); Health Action by People, Thiruvananthapuram, Kerala, India (Prof S Nair, Prof K Vijayakumar MD); Dr Somervell Memorial CSI Medical College, Karakonam, Thiruvananthapuram, Kerala, India (Prof K Vijayakumar); Fortis Escorts Hospitals, JLN Marg, Jaipur, India (I Mohan MD, R Gupta PhD); Madras Diabetes Research Foundation, Chennai, India (D Mohan PhD, S Rani PhD, Prof V Mohan MD); Department of Community Health Sciences (R Iqbal PhD) and Department of Medicine (R Iqbal, K Kazmi MD), Aga Khan University, Karachi, Pakistan; Independent University, Dhaka, Bangladesh (Prof O Rahman DSc, R Yusuf PhD); and Post Graduate Institute of Medical Education and Research (PGIMER) School of Public Health, Chandigarh, India (L V M Pinnaka MD, Prof R Kumar MD)

Correspondence to:  
Dr MyLinh Duong, Department  
of Medicine and Respiriology,  
McMaster University and  
Hamilton Health Sciences,  
Hamilton, ON, L8V 1C3, Canada  
duongmy@mcmaster.ca

### Research in context

#### Evidence before this study

We searched PubMed, Embase, the Cochrane database, and bibliographies of retrieved articles for relevant reports published in English between Jan 1, 1960, and Jan 1, 2016. We used key search terms “beedi”, “beedis”, “bidi”, “bidis”, “tobacco-smoking”, and “India”, “south Asia” to identify reports of bidi smoking on health outcomes including mortality, cardiorespiratory health, and lung function in south Asia. We found few reports published after 2000. Previous publications were not methodologically robust, and reported data derived from retrospective, cross-sectional, or case-control studies with limited and variable adjustments for potential confounders such as differences in socioeconomic and baseline risk factors. Furthermore, very few studies directly compared the effects of bidi smoking with cigarette smoking on cardiorespiratory outcomes and lung function.

#### Added value of this study

To our knowledge, this is the largest prospective multisite study with standardised approaches for data collection

(covariates, cardiorespiratory symptoms, and lung function measurements) and ascertainment of follow-up outcomes (deaths and cardiorespiratory events) in India, Bangladesh, and Pakistan. We examined the effects of cigarette and bidi smoking on several outcomes, adjusting for important differences in socioeconomic and baseline risk factors. A coherent pattern of worst outcomes was seen for bidi smokers, including highest self-reported baseline respiratory symptoms, obstructive ventilatory impairment, and follow-up mortality and cardiorespiratory events compared with cigarette smokers and non-smokers.

#### Implications of all the available evidence

Our findings fill an important gap in knowledge about the many harmful health effects of bidi smoking. These findings can be used for evidence-based practice and policy making that will help bring about greater controls on bidi use in south Asia.

“natural and safe” tobacco alternatives to cigarettes.<sup>3</sup> This claim contrasts with the toxicology data, which have shown higher nicotine, tar, and carbon monoxide levels delivered for a lower content of tobacco in bidis than in cigarettes.<sup>7</sup> However, there is a paucity of clinical data on the health effects of bidis. Most of these data are derived from retrospective, cross-sectional, or case-control studies, most with small sample sizes and limited adjustments for a wide range of potential confounders.<sup>8–11</sup>

We prospectively assessed the effects of bidi and cigarette smoking on mortality, respiratory, and cardiovascular outcomes in an unselected community-based cohort of men in south Asia. All comparisons were adjusted for differences in socioeconomic status, user-specific characteristics, and baseline risk factors. Furthermore, cross-sectional comparisons of self-reported baseline respiratory symptoms and spirometry measurements were done to provide information about baseline respiratory morbidity. The high prevalence of low-intensity smoking (ie,  $\leq 10$  pack-years) in this cohort provided an opportunity to assess the effects of low-intensity tobacco smoking on health outcomes.

## Methods

### Study design and participants

This substudy of the Prospective Urban Rural Epidemiology (PURE) study included community-based participants from India (five centres), Pakistan (one centre), and Bangladesh (one centre). Details of the overall PURE study design have been described elsewhere<sup>12</sup> and are summarised in the appendix (pp 5–9). The centres were purposely chosen to provide a diverse range of economic, physical, and sociocultural

environments, balanced by the feasibility of centres to achieve long-term follow-up. Standardised approaches were used for the enumeration of households, identification of individuals, recruitment, and data collection. The methods of approaching households differed between countries, but aimed to avoid biases in participant selection. Households with at least one member aged 35–70 years who were intending to stay locally for more than 4 years were approached. The final sample size for analysis varied by the outcome of interest and included only men with no missing data relevant for the outcome of interest. Only men were selected because the rate of smoking in women in south Asia were low. Similarly, former and current smokers were combined as ever-smokers for all analyses.

All eligible individuals who provided written informed consent were enrolled. Baseline data were collected from Jan 1, 2003, to Dec 30, 2009, and follow-up data from Jan 1, 2008, to Dec 30, 2013. The study was coordinated by the Population Health Research Institute (Hamilton, ON, Canada) and approved by the Hamilton Health Sciences Research Ethics Board and by the local ethics committee at each site.

### Procedures

Standardised interview-based questionnaires adapted from previous cohort studies (appendix p 10) were administered by trained personnel to household members aged 35–70 years. The questionnaire elicited demographic, household, behavioural, and medical information (risk factors, symptoms, comorbid disorders). Bidi and cigarette use was defined as self-reported duration of use more than 0 days or quantity


See Online for appendix

more than none per day. Former smokers were defined as individuals who had not smoked for at least 12 months. Participants were classified into four categories based on the number of pack-years of smoking (duration [years]×quantity [sticks per day] divided by 20) and tobacco type: non-smokers of bidis or cigarettes (never-smokers), light smokers of either or both products ( $\leq 10$  pack-years), heavy smokers of cigarettes only ( $>10$  pack-years), and heavy smokers of bidis (including concurrent use of cigarettes;  $>10$  pack-years). Other relevant data included cooking fuel (solid or kerosene vs gas or electricity), education, asset index, proportion of income spent on food, physical activity, dietary intake, and cardiorespiratory symptoms (dyspnoea with usual activity; wheeze; cough or sputum; chest pain) occurring at least weekly in the previous 6 months.

Household owned items were used to generate an asset index, an indicator of wealth. Physical activity was assessed with the International Physical Activity Questionnaire (IPAQ).<sup>13</sup> Dietary intake was assessed with a validated food frequency questionnaire.<sup>14</sup> Physical measurements were collected by standardised methods

for anthropometrics, blood pressure, handgrip strength, and spirometry. An individualised INTERHEART risk score was calculated with the version excluding cholesterol concentration. This score provided a validated and quantitative measure of the risk-factor burden for cardiovascular disease, which incorporates information about self-reported age, sex, cigarette use, diabetes, family history of cardiovascular disease, psychosocial factors, diet, physical activity, and measured waist-to-hip ratio and hypertension ( $\geq 140/90$  mm Hg).<sup>15</sup>

Lung function was measured with a portable spirometer (MicroGP, MicroMedical Ltd, Chatham, IL, USA), which did not generate spirograms. Each participant attempted up to six forced prebronchodilator manoeuvres while standing and wearing a nose clip. Each manoeuvre was closely observed for maximal effort, with exhalation time 6 s or more and without coughing. Spirometers were calibrated monthly (3 L syringe) or before each use in extreme temperature or handling. For analyses, we selected participants with at least two forced expiratory volume in 1 s (FEV<sub>1</sub>) and forced vital capacity (FVC) measurements within 200 mL variability. The quality of the spirometry data in PURE have previously been



**Figure 1: Participant selection for the substudy**

PURE=Prospective Urban Rural Epidemiology Study. \*Participants with poor-quality lung function data were those with less than two measurements of forced expiratory volume in 1 s (FEV<sub>1</sub>) or forced vital capacity (FVC), or with variability between two the highest FEV<sub>1</sub> or FVC measurements of  $>200$  mL. †Participants with self-reported diagnoses of cardiorespiratory disease, stroke, cancer, tuberculosis, or HIV infection at baseline were excluded from the analysis of follow-up events.

validated for external, internal, and face validity (appendix pp 11–16).<sup>16</sup> The highest FEV<sub>1</sub>, FVC, and FEV<sub>1</sub>/FVC ratio recorded for each individual was included in the analysis.

Participants were followed up with yearly telephone calls and face-to-face interviews every 3 years. At each contact, participants, or close relatives in cases of deaths (via verbal autopsy<sup>17</sup>), were questioned as to whether any clinical events had occurred and if so, documentation was obtained for event adjudication by the site investigators using standard definitions (appendix pp 17–20). Furthermore, a random subset of events from each site was assessed centrally (Population Health Research Institute) to ensure consistency in the adjudicated events.

To ensure standardisation and data quality, comprehensive operations manuals, reinforced by periodic training workshops, training DVDs, and regular communication were used in all sites. Data were entered locally by each site into a customised database programmed

with range and consistency checks and transmitted electronically to the coordinating centre, where further quality control measures were implemented.

### Statistical analysis

We used multilevel marginal regression to estimate the effect size by smoking category on outcomes relative to non-smokers adjusted for age, body-mass index, asset index, education, cooking fuel, INTERHEART risk score, diabetes, hypertension, and centre (except for spirometry measurements and respiratory events). Community was treated as a random effect to account for data clustering. Logistic regression provided estimates on the adjusted prevalence and odds ratios (ORs) for cross-sectional baseline symptoms. We used linear regression to compare the cross-sectional age-related changes on baseline FEV<sub>1</sub> and FEV<sub>1</sub>/FVC ratio by smoking category adjusted for height, weight, centre, and education (FEV<sub>1</sub>) or age, height, and centre (FEV<sub>1</sub>/FVC ratio). We used Cox proportional hazards models to estimate the incidences and hazard ratios (HRs) for all-cause mortality and cardiovascular events (myocardial infarction, stroke, heart failure, sudden death, cardiovascular-related death, and cardiovascular-related hospital admission). For respiratory events (chronic obstructive pulmonary disease, asthma, pneumonia, tuberculosis), the same Cox model was used but without INTERHEART risk score, diabetes, and hypertension as covariates. To compare the effect size estimates by smoking category, smoking category was fitted as a categorical variable and its coefficient reflects the nature of association between effect size and smoking category.

There was no formal sample size calculation for this substudy. We assessed the adequacy of the sample size using guidelines proposed by Concato and colleagues,<sup>18</sup> which recommend that for Cox regression at least ten events for each degree of freedom (df) are needed to provide stable models. There were 685 deaths, 552 cardiovascular events, and 269 respiratory events recorded during follow-up. The model for deaths and cardiovascular events contained nine covariates including the smoking categories, giving a total of 22 df. For the respiratory event model, there were five covariates giving a total of 16 df. Thus the events per df for mortality, cardiovascular, and respiratory events were 31 (685/22), 25 (552/22), and 16 (269/16), respectively. These estimates are all greater than ten events per df, indicating the sample size was adequate to provide stable models for all three event outcomes.

Some sensitivity analyses were carried out to examine for any changes to our main findings on outcome events using different criteria for selection of study population (including participants with significant baseline comorbidities); classification of smoking categories (excluding former smokers or smokers of both bidis and cigarettes) and using different socioeconomic status covariates (education, rural or urban location, percent income spent of food, and the combination of these). All

	Non-smokers (n=8438)	Light smokers (n=3321)	Heavy cigarette smokers (n=959)	Heavy bidi smokers (n=2201)
<b>Tobacco type</b>				
Cigarettes only	..	1811 (55%)	959 (100%)	0
Bidis only	..	1330 (40%)	0	1627 (74%)
Both	..	180 (5%)	0	574 (26%)
Smokeless tobacco use*	696 (8%)	227 (7%)	57 (6%)	121 (5%)
Age (years)	49.3 (10.6)	47.9 (10.0)	53.0 (9.1)	51.9 (10.1)
Height (cm)	164.9 (7.4)	165.1 (7.2)	165.4 (6.9)	164.2 (6.7)
Weight (kg)	63.6 (13.7)	59.8 (13.4)	63.2 (13.6)	54.0 (10.8)
Body-mass index (kg/m <sup>2</sup> )	23.4 (4.4)	22.0 (4.3)	23.0 (3.9)	20.2 (3.5)
Urban residency	4649 (55%)	1414 (43%)	578 (60%)	504 (23%)
<b>Education</b>				
Secondary or higher	5595 (67%)	1836 (56%)	633 (66%)	615 (28%)
Primary or none	2695 (33%)	1456 (44%)	321 (34%)	1573 (72%)
Missing data	148 (2%)	29 (1%)	5 (1%)	13 (1%)
Asset index†	-0.8 (1.0)	-1.3 (0.8)	-0.7 (0.7)	-1.5 (0.5)
Proportion of income spent on food (%)‡	54.3% (26.2)	62.3% (25.0)	58.5% (23.1)	68.3% (22.4)
<b>Cooking fuel</b>				
Gas or electricity	4735 (61%)	1338 (43%)	583 (62%)	394 (21%)
Kerosene or solid fuel	3021 (39%)	1751 (57%)	353 (38%)	1502 (79%)
Missing data	682 (8%)	232 (7%)	23 (2%)	305 (14%)
Manual labour occupations§	2267/8438 (45%)	987/3321 (63%)	421/959 (60%)	1114/ (87%)
<b>Location¶</b>				
Bangladesh	572 (7%)	312 (9%)	211 (22%)	238 (11%)
Chandigarh, India	1392 (16%)	89 (3%)	18 (2%)	127 (6%)
Chennai, India	1265 (15%)	473 (14%)	116 (12%)	306 (14%)
Thiruvananthapuram, India	985 (12%)	549 (17%)	309 (32%)	228 (10%)
Bengaluru, India	2487 (29%)	1426 (43%)	146 (15%)	792 (36%)
Jaipur, India	1190 (14%)	344 (10%)	13 (1%)	507 (23%)
Pakistan	547 (6%)	128 (4%)	146 (15%)	3 (<1%)

(Table 1 continues on next page)

analyses were done with SAS version 9.2. No adjustments for multiplicity of analysis were made.

### Role of the funding source

The funders of the study had no role in study design, data collection, data analysis, data interpretation, or writing of the report. The corresponding author had full access to all the data in the study and had final responsibility for the decision to submit for publication.

### Results

14 919 men from 158 communities across seven centres in south Asia were included in this substudy (figure 1). Bidi smoking was more common in India, whereas cigarette smoking was more common in Bangladesh and Pakistan (table 1). A small proportion of smokers reported using both tobacco types; bidis were the predominant type. The demographics of participants who smoked both bidis and cigarettes were similar to the demographics of those who smoked bidis only; we therefore reclassified this group as bidi smokers (appendix p 21). A small and similar proportion of participants across the smoking categories reported the use of smokeless tobacco. There was a small proportion of former smokers and the numbers were evenly distributed across the smoking categories. Overall, heavy bidi smokers were more likely to come from rural communities and have a low socioeconomic status than were men in all other smoking categories (lower education and asset index; higher percentages of income spent on food, manual labour occupations, and use of solid or kerosene cooking fuels).

The adjusted prevalence of self-reported chronic cardiorespiratory symptoms at baseline was significantly higher in heavy cigarette smokers and heavy bidi smokers than in non-smokers (table 2, figure 2). The largest effect was seen for chronic cough and sputum. Light smokers showed a modest increase in adjusted prevalence of chronic symptoms relative to non-smokers; however, because of low reported rates of wheeze and dyspnoea, the increase in these symptoms was not significant. The prevalence of chronic symptoms at baseline increased across the categories of non-smokers, light smokers, heavy cigarette smokers, and heavy bidi smokers ( $p < 0.001$  for association).

For the cross-sectional analysis of spirometry measurements, centres from Pakistan, Jaipur (India), and Bangladesh were excluded because of high proportions of participants with missing spirometry data (figure 1). We therefore examined spirometry data from the five centres in India. The adjusted cross-sectional age-related changes in FEV<sub>1</sub> and FEV<sub>1</sub>/FVC ratio were significantly larger for heavy bidi smokers than for men in the other smoking categories (table 3, figure 3). This finding suggests increasingly lower lung function in the older age groups of bidi smokers. By contrast, age-related changes in FEV<sub>1</sub> did not differ between non-smokers, light smokers, and heavy cigarette smokers. However, similar pairwise comparison

	Non-smokers (n=8438)	Light smokers (n=3321)	Heavy cigarette smokers (n=959)	Heavy bidi smokers (n=2201)
(Continued from previous page)				
Smoking history				
Age of smoking initiation (years)	..	31.0 (11.6)	22.7 (8.4)	22.7 (9.1)
Duration of smoking (years)	..	15.7 (10.6)	29.2 (10.1)	28.7 (11.0)
Tobacco sticks smoked per day	..	6.1 (5.0)	18.2 (13.6)	21.0 (11.7)
Tobacco stick-years¶	..	83.0 (61.4)	494.3 (378.6)	575.4 (393.3)
Pack-years	..	4.2 (3.1)	24.7 (18.9)	28.8 (19.7)
Current smokers**	..	2934 (88%)	807 (84%)	1995 (91%)
Former smokers	..	364 (11%)	145 (15%)	192 (9%)
Missing data	..	23 (1%)	7 (1%)	14 (1%)

Data are n (%) or mean (SD). Non-smokers self-reported no bidi or cigarette use at baseline survey; light smokers reported smoking ten or fewer pack-years of either bidis, cigarettes, or both; and heavy smokers (>10 pack-years) are divided into those who smoked cigarettes only and those who smoked bidis with or without cigarettes. ..not applicable. \*Smokeless tobacco use was defined as self-reported duration of use more than 0 days or quantity more than none per day of smokeless tobacco (chewed tobacco, snuff, or rolled tobacco leaves). †Asset index is the non-monetary aspect of wealth based on the number and type of household items owned (a high positive value indicates greater wealth). ‡Lower-income families spend a greater percentage of total income on food. §Manual labour includes workers in agricultural, fishery, and craft industries, plant/machine operators, assemblers, and elementary workers; missing data mainly due to retirement. ¶Tobacco stick-years=number of tobacco sticks smoked per day × duration of smoking (years). ||Pack-years=tobacco stick-years/20. \*\*Current smokers were defined as individuals who reported use of at least one tobacco stick per day within 12 months.

**Table 1: Baseline characteristics**

showed a significantly reduced FEV<sub>1</sub>/FVC ratio, suggesting mild obstructive ventilatory impairment, in light smokers and heavy cigarette smokers compared with non-smokers.

Mean duration of follow-up was 5.6 years (range 1–13). 13 264 (98%) participants completed follow-up, with a similar proportion in each smoking category (7554 [98%] non-smokers; 2966 [98%] light smokers; 819 [99%] heavy cigarette smokers; 1925 [98%] heavy bidi smokers). Participants with self-reported baseline cardiovascular disease, respiratory disease, cancer, or HIV infection were excluded from the analysis of follow-up events (figure 1). The demographics of the excluded and included participants were similar (appendix pp 21). In the analysis population, there were 685 deaths, 552 cardiovascular events, and 269 respiratory events (table 4). Heavy bidi smokers had the highest incidence of follow-up events compared with men in the other smoking categories (table 4, figure 4). The largest effect of heavy bidi smoking was on respiratory events. The incidence of follow-up events in light smokers and heavy cigarette smokers was intermediate between that for non-smokers and heavy bidi smokers; however, the increase in incidence of respiratory events in light smokers and heavy cigarette smokers compared with non-smokers was not significant. The HRs for follow-up events increased across the categories of light smokers, heavy cigarette smokers, and heavy bidi smokers.


We did sensitivity analyses that included all participants with complete vital statistics; adjusted for other indicators



	Non-smokers (n=8438)	Light smokers (n=3321)	Heavy cigarette smokers (n=959)	Heavy bidi smokers (n=2201)
<b>Wheeze</b>				
Number	425	234	69	242
Adjusted prevalence (95% CI)	4.3% (3.4-5.4)	5.1% (3.9-6.6)	6.1% (4.2-8.7)	8.0% (6.2-10.2)
Adjusted OR (95% CI)	1.0	1.20 (0.97-1.50)	1.45 (1.10-1.92)	1.95 (1.59-2.39)
p value	..	0.092	0.009	<0.0001
<b>Cough or sputum</b>				
Number	1094	686	214	696
Adjusted prevalence (95% CI)	12.9% (11.0-15.2)	17.9% (15.2-20.8)	21.4% (18.1-25.1)	26.7% (23.2-30.5)
Adjusted OR (95% CI)	1.0	1.46 (1.27-1.69)	1.83 (1.49-2.24)	2.45 (2.07-2.89)
p value	..	0.0001	0.0001	0.0001
<b>Dyspnoea with usual activity</b>				
Number	966	469	170	508
Adjusted prevalence (95% CI)	12.8% (11.4-14.3)	15.0% (13.3-16.8)	19.7% (16.4-23.4)	19.6% (16.6-23.0)
Adjusted OR (95% CI)	1.0	1.20 (1.05-1.39)	1.67 (1.36-2.06)	1.67 (1.36-2.04)
p value	..	0.010	<0.0001	<0.0001
<b>Chest pain</b>				
Number	1241	677	181	612
Adjusted prevalence (95% CI)	15.9% (14.5-17.5)	19.9% (17.9-22.1)	20.2% (17.5-23.1)	23.6% (20.9-26.5)
Adjusted OR (95% CI)	1.0	1.31 (1.16-1.48)	1.33 (1.09-1.63)	1.63 (1.40-1.89)
p value	..	0.0001	0.006	0.0001

Self-reported baseline symptoms defined as those occurring at least weekly in the 6 months before the baseline survey. ORs (relative to non-smokers) were estimated with multilevel marginal logistic regression with age, asset index, body-mass index, and centre as covariates. Community was treated as a random effect to account for data clustering. OR=odds ratio.

**Table 2: Self-reported cardiorespiratory symptoms at baseline**



**Figure 2: Self-reported cardiorespiratory symptoms at baseline**  
 Self-reported baseline symptoms defined as those occurring at least weekly in the 6 months before the baseline survey. p value for association examined the order effect in effect size by different smoking groups (as categorical variable). Error bars represent 95% CI.

of socioeconomic status alone or in combination; and excluded former smokers and smokers of both bidis and cigarettes (appendix p 23–24). Results of sensitivity analyses showed the conclusions remained unchanged.

**Discussion**

In this prospective community-based cohort study of unselected men in south Asia, we recorded a high prevalence of bidi and cigarette use. Bidi smokers were more likely to be from rural areas and have low socioeconomic status. Accounting for these differences

and additional baseline risk factors, bidi use was consistently associated with significantly increased prevalences and relative risks of baseline cardiorespiratory symptoms, low ventilatory capacity, and follow-up mortality and cardiorespiratory outcomes. Light smokers and heavy cigarette smokers also showed increased risks of death and cardiovascular events relative to non-smokers, but not for respiratory events. The observed pattern of greater baseline respiratory morbidity and higher risks of mortality and cardiorespiratory outcomes in heavy bidi smokers suggest that bidis are at least as harmful as cigarettes and contribute to the burden of tobacco-related disease and deaths in south Asia.

Our reported rates and pattern of bidi use are consistent with previous data, indicating a high prevalence of bidi smoking in south Asia, with geographical<sup>19</sup> and socioeconomic variation.<sup>20,21</sup> These baseline differences could potentially confound the relation between bidi smoking and health outcomes. Our large sample size and data on socioeconomic status and other characteristics made it possible to adjust for a large number of potentially important confounders, in order to derive an unbiased estimate of the independent effect of bidi smoking on cardiorespiratory health and mortality. In addition to the covariates included in the final model, we also explored other potential confounders in several sensitivity analyses and found their effects were small and did not substantially change the overall model or conclusions.

	Non-smokers (n=6701)	Light smokers (n=2849)	Heavy cigarette smokers (n=800)	Heavy bidi smokers (n=1691)
<b>FEV<sub>1</sub></b>				
Change per year, mL/s (95% CI)	-22.1 (-23.8 to -20.4)	-22.5 (-24.2 to -20.8)	-22.7 (-24.7 to -20.8)	-23.8 (-25.6 to -22.0)
Difference from non-smokers	0	-0.3 (-0.9 to 0.2)	-0.6 (-1.5 to 0.3)	-1.7 (-2.3 to -1.0)
p value	..	0.198	0.191	<0.0001
<b>FEV<sub>1</sub>/FVC ratio</b>				
Change per year (95% CI)	-0.145% (-0.171 to -0.119)	-0.160% (-0.184 to -0.136)	-0.158% (-0.184 to -0.132)	-0.180% (-0.207 to -0.153)
Difference from non-smokers	0	-0.015% (-0.027 to -0.003)	-0.013% (-0.028 to 0.002)	-0.035% (-0.049 to -0.022)
p value	..	0.012	0.097	<0.0001


Multilevel marginal linear regression was used to estimate the cross-sectional age-related change in forced expiratory volume in 1 s (FEV<sub>1</sub>) or FEV<sub>1</sub>/forced vital capacity (FVC) ratio by smoking category adjusted for height, weight, centre, and education (FEV<sub>1</sub>) or age, height, and centre (FEV<sub>1</sub>/FVC ratio). Community was treated as a random effect. Differences in the cross-sectional age-related change relative to non-smokers were also adjusted for the same covariates.

**Table 3: Cross-sectional age-related changes in FEV<sub>1</sub> and FEV<sub>1</sub>/FVC ratio**

Few contemporary data exist about the respiratory effects of bidis. Early studies showed an association between bidi smoking with chronic cough and sputum.<sup>22–25</sup> In keeping with these findings, we noted that the prevalence and risk of cardiorespiratory symptoms were consistently higher in heavy bidi smokers than in men in the other smoking categories; bidi smokers also had the lowest adjusted ventilatory capacity. The finding for ventilatory capacity had not been consistently documented across earlier studies, which were limited by small sample sizes and variability in their adjustments of potential confounders.<sup>24,26,27</sup> Our study is the largest study so far to compare spirometry data between bidi and cigarette smokers. Our findings suggest lower ventilatory capacity and greater airflow obstruction in heavy bidi smokers, and together with a higher prevalence of cardiorespiratory symptoms indicate substantially higher rates of obstructive respiratory impairment in bidi smokers compared with other smokers. Heavy cigarette smokers and light smokers also showed lower adjusted lung function than did non-smokers, but the magnitude of this difference did not reach significance. This finding might relate to the smaller sample size in these subgroups or the greater variability in the effect of cigarette smoking on lung function.

Large population-based studies have examined the incidence of cancer, cardiovascular disease, and all-cause mortality in tobacco users in south Asia.<sup>9,10,28–31</sup> To our knowledge, only one other publication has reported on the effect of bidis separately from cigarettes on all-cause mortality (HR relative to never-smokers 1.64 [95% CI 1.47–1.81] for bidi smokers vs 1.37 [1.23–1.53] for cigarette smokers).<sup>9</sup> Our work supports and extends this finding by showing that the risks of cardiorespiratory events are also significantly increased with bidi smoking independent of socioeconomic status.

Several aspects of this study are worthy of discussion. First, our study population included a large number of light smokers, particularly of cigarettes. Separating out the light smokers provided a more balanced and matched distribution of smoking intensity and smoking pattern



**Figure 3: Mean adjusted FEV<sub>1</sub> and FEV<sub>1</sub>/FVC ratio by age group**


Multilevel marginal linear regression was used to model forced expiratory volume in 1 s (FEV<sub>1</sub>) by age group and smoking category with height, weight, centre, and education as covariates (reference: non-smokers) and community as a random effect. A similar model was used for the comparison of FEV<sub>1</sub>/forced vital capacity (FVC) ratio by age group and smoking category with age, height, and centre as covariates (reference: non-smokers) and community as a random effect.

between heavy bidi smokers and heavy cigarette smokers for comparison. We found that low-intensity smoking can also be associated with respiratory impairment at an

	Non-smokers (n=7554)	Light smokers (n=2966)	Heavy cigarette smokers (n=819)	Heavy bidi smokers (n=1925)
<b>Cardiovascular events</b>				
Number	234	144	65	109
Adjusted incidence (95% CI)	3.1% (2.4-3.9)	4.6% (3.4-6.3)	4.4% (3.0-6.3)	5.1% (3.5-7.3)
Adjusted HR (95% CI)	1.0	1.45 (1.13-1.84)	1.47 (1.05-2.06)	1.55 (1.17-2.06)
p value	..	0.003	0.023	0.002
<b>Respiratory events</b>				
Number	99	61	27	82
Adjusted incidence (95% CI)	0.7% (0.4-1.1)	0.9% (0.6-1.5)	0.8% (0.4-1.5)	1.3% (0.8-2.2)
Adjusted HR (95% CI)	1.0	1.30 (0.91-1.85)	1.21 (0.70-2.07)	1.73 (1.23-2.45)
p value	..	0.154	0.496	0.002
<b>Deaths</b>				
Number	279	165	58	183
Adjusted incidence (95% CI)	2.3% (1.7-3.0)	3.1% (2.1-4.5)	3.3% (2.2-5.0)	3.8% (2.7-5.5)
Adjusted HR (95% CI)	1.0	1.28 (1.02-1.62)	1.59 (1.13-2.24)	1.56 (1.22-1.98)
p value	..	0.034	0.008	<0.0003

New cardiovascular events (cardiovascular-related death, myocardial infarction, heart failure, stroke, and cardiovascular-related hospital admission), respiratory events (chronic obstructive pulmonary disease [COPD], pneumonia, tuberculosis, asthma), and all-cause mortality at a mean follow-up of 5.6 years. Incidence and HRs (relative to non-smokers) for all-cause mortality and cardiovascular events were adjusted for age, body-mass index, asset index, education, handgrip, INTERHEART risk score, diabetes, and hypertension, and centre as fixed-effect covariates, and community as a random effect in a multilevel marginal Cox proportional hazards model. For respiratory events, the same model was used but without INTERHEART risk score, diabetes, and hypertension as covariates in the model. HR=hazard ratio. \*Participants with missing data or with self-reported diagnoses of stroke, heart disease, cancer, COPD, asthma, tuberculosis, or HIV infection were excluded from the analysis.

**Table 4: Respiratory events, cardiovascular events, and deaths during follow-up\***



**Figure 4: Respiratory events, cardiovascular events, and deaths during follow-up**  
New cardiovascular events (cardiovascular-related death, myocardial infarction, heart failure, stroke, and cardiovascular-related hospital admission), respiratory events (chronic obstructive pulmonary disease, pneumonia, tuberculosis, asthma), and all-cause mortality at a mean follow-up of 5.6 years. Hazard ratios (relative to non-smokers) for all-cause mortality and cardiovascular events were estimated with a marginal Cox proportional hazards model with age, education, asset index, body-mass index, handgrip, self-reported diabetes, hypertension, INTERHEART risk score, and centre as fixed effects and community as a random effect. For respiratory events, the same model was used but without INTERHEART risk score, diabetes, and hypertension as covariates in the model. p value for association examined the order effect in effect size by different smoking groups (as categorical variable). Error bars represent 95% CI.

exposure level that is often considered to be clinically trivial. Furthermore, this level of exposure was associated with increased risks of cardiovascular events and mortality compared with not smoking. This finding suggests that there is no threshold that can be considered safe from the harmful effects of tobacco smoking. Second, bidis are smaller with less tobacco content and are generally sold in different quantities from cigarettes. These differences make comparison of the two tobacco types using a common unit of exposure such as pack-years difficult. To maintain consistency and allow ease of comparison with the literature on tobacco, we continued to use pack-years to define the groups. However, any interpretation of the effect size of bidi use on outcomes must take into account of the lower tobacco content in bidis compared with cigarettes.<sup>7</sup> Finally, only participants with no previous cardiorespiratory morbidity, cancer, or HIV infection were analysed for follow-up events to avoid the effect of reverse causality on our findings.

There are limitations and strengths to our study. It was not feasible to aim for strict proportionate sampling in this large prospective cohort study. The design did not use standard random sampling but adopted a design that avoided biases in levels of risk factors and prevalence of disease conditions. Second, lung function was measured with a portable spirometer that did not provide flow-volume curves and therefore verification of individual effort was not possible. However, we had previously validated this method by comparing data obtained in certified pulmonary function laboratories with data obtained by PURE methods for 531 participants from participating sites including south Asia; we noted high correlations without biases.<sup>16</sup> Further, there is no a-priori reason to expect differential effects in methods on spirometry measurements between different smoking groups. The major strengths of our study include the large sample size, the prospective and standardised approach to data collection and outcome ascertainment, all of which provide for a robust and systematic analysis, and adjustment for a large number of potential confounders.

Our findings have important public health implications. South Asia is the second largest consumer of tobacco in the world, with more than 130 million tobacco smokers.<sup>2</sup> More than half these smokers use bidis, particularly among the poorest and most vulnerable sectors of the population. Therefore the health impact of bidis is highly relevant in this region. Furthermore, as the current trend of bidi exportation continues, the global impact of bidis will rise, particularly among young adults. Our findings suggest that bidis are at least as harmful as cigarettes on cardiorespiratory health and mortality, despite having substantially less tobacco content than cigarettes. Control of bidis should be an integral part of any anti-tobacco framework, both regionally and globally, with the greatest potential effects to be seen in poor and young people.

**Contributors**

All authors contributed to the intellectual conceptualisation of PURE, study design, planning, and collection of PURE data. MD, SR, XZ, KK,

and SY contributed to the statistical analysis and wrote the report. All authors contributed to the final approval of the report. MD, KK, POB, and SY take full responsibility for the overall content of this work.

#### Declaration of interests

We declare no competing interests.

#### Acknowledgments

The main PURE study and its components are funded by the Population Health Research Institute, the Canadian Institutes of Health Research, Heart and Stroke Foundation of Ontario, and through unrestricted grants from several pharmaceutical companies (with major contributions from AstraZeneca [Canada], Sanofi-Aventis [France and Canada], Boehringer Ingelheim [Germany and Canada], Servier, and GlaxoSmithKline), and additional contributions from Novartis and King Pharma. Various national or local organisations in participating countries also contributed: Fundacion ECLA (Argentina); Independent University, Bangladesh, and Mitra and Associates (Bangladesh); Unilever Health Institute (Brazil); Public Health Agency of Canada and Champlain Cardiovascular Disease Prevention Network, Canadian Institutes of Health Research, Heart and Stroke Foundation of Canada, and Population Health Research Institute (Canada); Universidad de la Frontera (Chile); National Center for Cardiovascular Diseases (China); Colciencias (grant 6566-04-18062; Colombia); Indian Council of Medical Research (India); Ministry of Science, Technology and Innovation of Malaysia (grant 100 - IRDC / BIOTEK 16/6/21 [13/2007], grant 07-05-IFN-BPH 010), Ministry of Higher Education of Malaysia (grant 600-RMI/LRGS/5/3 [2/2011]), Universiti Teknologi MARA, and Universiti Kebangsaan Malaysia (UKM-Hejima-Komuniti-15-2010; Malaysia); Polish Ministry of Science and Higher Education (grant 290/W-PURE/2008/0) and Wroclaw Medical University (Poland); North-West University, SANPAD (South Africa and Netherlands Programme on Alternatives in Development), National Research Foundation, Medical Research Council of South Africa, South African Sugar Association (SASA), and Faculty of Community and Health Sciences (UWC; South Africa); AFA Insurance, Swedish Council for Working Life and Social Research, Swedish Research Council for Environment, Agricultural Sciences and Spatial Planning, Swedish Heart and Lung Foundation, Swedish Research Council, the Swedish State under LäkarUtbildningsAvtalet agreement, and the Västra Götaland Region (FOUU; Sweden); Metabolic Syndrome Society, AstraZeneca, and Sanofi-Aventis (Turkey); and Sheikh Hamdan Bin Rashid Al Maktoum Award For Medical Sciences and Dubai Health Authority (United Arab Emirates). SY holds the Heart and Stroke Foundation/Marion W Burke Chair in Cardiovascular Disease. We would like to acknowledge the following individuals for their input and assistance in preparation of this report: Justina Greene for data cleaning and Shofiqul Islam, Amparo Casanova, and Lehana Thabane for statistical advice and guidance.

#### References

- World Health Organization. WHO report on the global tobacco epidemic, 2015: the MPOWER package. [http://www.who.int/tobacco/global\\_report/2015/en/](http://www.who.int/tobacco/global_report/2015/en/) (accessed Nov 28, 2016).
- Giovino GA, Mirza SA, Samet JM, et al, for the GATS Collaborative Group. Tobacco use in 3 billion individuals from 16 countries: an analysis of nationally representative cross-sectional household surveys. *Lancet* 2012; **380**: 668–879.
- Centers for Disease Control and Prevention. Smoking & tobacco use: bidis and kreteks. [http://www.cdc.gov/tobacco/data\\_statistics/fact\\_sheets/tobacco\\_industry/bidis\\_kreteks/index.htm](http://www.cdc.gov/tobacco/data_statistics/fact_sheets/tobacco_industry/bidis_kreteks/index.htm) (accessed Nov 28, 2016).
- Kostova D, Chaloupka FJ, Yurekli A, et al, for the GATS Collaborative Group. A cross-country study of cigarette prices and affordability: evidence from the Global Adult Tobacco Survey. *Tob Control* 2014; **23**: e3.
- Jha P. Deaths and taxes: stronger global tobacco control by 2025. *Lancet* 2015; **385**: 918–20.
- Sankaran S, Hillamo H, Glantz SA. Implementation of graphic health warning labels on tobacco products in India: the interplay between the cigarette and the bidi industries. *Tob Control* 2015; **24**: 547–55.
- Watson CH, Polzin GM, Calafat AM, Ashley DL. Determination of tar, nicotine and carbon monoxide yields in the smoke of bidi cigarettes. *Nicotine Tob Res* 2003; **5**: 747–53.
- Gajalakshmi V, Peto R, Kanak TS, Jha P. Smoking and mortality from tuberculosis and other diseases in India: retrospective study of 43 000 adult male deaths and 35 000 controls. *Lancet* 2003; **362**: 507–15.
- Gupta PC, Pednekar MS, Parkin DM, Sankaranarayanan R. Tobacco associated mortality in Mumbai (Bombay) India. Results of the Bombay Cohort Study. *Int J Epidemiol* 2005; **34**: 1395–1402.
- Rahman M, Sakamoto J, Fujui T. Bidi smoking and oral cancer: a meta-analysis. *Int J Cancer* 2003; **106**: 600–04.
- Teo KK, Ounpuu S, Hawken S, et al, for the INTERHEART Study Investigators. Tobacco use and risk of myocardial infarction in 52 countries in the INTERHEART study: a case-control study. *Lancet* 2006; **368**: 647–58.
- Teo KK, Chow CK, Vaz M, Rangarajan S, Yusuf S, for the PURE Investigators-Writing Group. The Prospective Urban Rural Epidemiology (PURE) study: examining the impact of social influences on chronic non-communicable diseases in low-, middle-, and high-income countries. *Am Heart J* 2009; **158**: 1–7e1.
- Craig CL, Marshall AL, Sjostrom M, et al. International physical activity questionnaire: 12-country reliability and validity. *Med Sci Sports Exerc* 2003; **35**: 1381–95.
- Iqbal R, Ajayan K, Bharathi AV, Zhang X, Islam S, Soman CR, Merchant AT. Refinement and validation of an FFQ developed to estimate macro- and micronutrient intakes in a south Indian population. *Public Health Nutr* 2009; **12**: 12–16.
- McGorrigan C, Yusuf S, Islam S, et al, for the INTERHEART Investigators. Estimating modifiable coronary heart disease risk in multiple regions of the world: the INTERHEART Modifiable Risk Score. *Eur Heart J* 2011; **32**: 581–89.
- Duong M, Islam S, Rangarajan S, et al, for the PURE-BREATH Study Investigators. Global differences in lung function by region (PURE): an international community-based prospective study. *Lancet Respir Med* 2013; **8**: 599–609.
- Gajalakshmi V, Peto R. Verbal autopsy of 80,000 adult deaths in Tamilnadu, South India. *BMC Public Health* 2004; **4**: 47.
- Concato J, Peduzzi P, Holford TR, Feinstein AR. Importance of events per independent variable in proportional hazards analysis. I. Background, goals and general strategy. *J Clin Epidemiol* 1995; **48**: 1495–501.
- Mony PK. Geographical epidemiology of cardiovascular disease in India: an exploratory study. MSc thesis, University of Toronto, 2009.
- National Sample Survey Organization. National sample surveys: 1987-88 to 2000-01. New Delhi, India: Department of Statistics, Ministry of Planning and Programme Implementation, Government of India.
- Nichter M, Nichter M, Van Sickle D. Popular perceptions of tobacco products and patterns of use among male college students in India. *Soc Sci Med* 2004; **59**: 415–31.
- Malik SK, Singh K. Smoking habits, chronic bronchitis and ventilatory impairment in rural males. *Indian J Chest Dis Allied Sci* 1978; **12**: 54–57.
- Malik SK, Wahi P. Prevalence of chronic bronchitis in a group of North Indian adults. *J India Med* 1978; **70**: 6–8.
- Chhabra SK, Rajpal S, Gupta R. Patterns of smoking in Delhi and comparison of chronic respiratory morbidity among beedi and cigarette smokers. *Indian J Chest Dis Allied Sci* 2001; **43**: 19–26.
- Jindal SK, Aggarwal AN, Chaudhry SK, et al. A multicentric study on epidemiology of chronic obstructive pulmonary disease and its relationship with tobacco smoking and environmental tobacco smoke exposure. *Indian J Chest Dis Allied Sci* 2006; **48**: 23–29.
- Dhand R, Malik S, Sharda PK. Long term effects of tobacco smoking: results of a spirometric study in 300 old men. *Indian J Chest Dis Allied Sci* 1985; **27**: 44–47.
- Khan MM, Tandon SN, Khan MT, Pandey US, Idris MZ. A comparative study of effects of cigarette and bidi smoking on respiratory function tests. *J Environ Biol* 2002; **23**: 89–93.
- Gupta PC, Bhonsle RB, Mehta FS, Pindborg JJ. Mortality experience in relation to tobacco chewing and smoking habits from 10-years follow-up study in Ernakulam district, Kerala. *Int J Epidemiol* 1984; **13**: 184–87.
- Gupta PC, Mehta F, Irani RR. Comparison of mortality rates among bidi smokers and tobacco chewers. *Indian J Cancer* 1980; **17**: 149–52.
- Pednekar MS, Gupta PC, Yeole BB, Hebert JR. Association of tobacco habits, including bidi smoking, with overall and site-specific cancer incidence: results from the Mumbai cohort study. *Cancer Causes Control* 2011; **22**: 859–68.
- Wu F, Chen Y, Parvez F, et al. A prospective study of tobacco and mortality in Bangladesh. *PLoS One* 2013; **8**: e58516.