



THE AGA KHAN UNIVERSITY

eCommons@AKU

Department of Radiology

Medical College, Pakistan

8-2012

Observer variation in MRI evaluation of patients with suspected lumbar disc herniation and nerve root compression: Comparison of neuroradiologist and neurosurgeon's interpretations

Imaad-ur Rehman

Rana Shoaib Hamid

Waseem Akhtar

Muhammad Shahzad Shamim

Rohana Naqi

See next page for additional authors

Follow this and additional works at: https://ecommons.aku.edu/pakistan_fhs_mc_radiol



Part of the [Neurosurgery Commons](#), and the [Radiology Commons](#)

Authors

Imaad-ur Rehman, Rana Shoaib Hamid, Waseem Akhtar, Muhammad Shahzad Shamim, Rohana Naqi, and
Haji Ismail Siddiq

Observer variation in MRI evaluation of patients with suspected lumbar disc herniation and nerve root compression: Comparison of Neuroradiologist and Neurosurgeon's interpretations

Imaad-ur-Rehman,¹ Rana Shoaib Hamid,² Waseem Akhtar,³ Muhammad Shahzad Shamim,⁴ Rohana Naqi,⁵ Haji Ismail Siddiq⁶

Department of Radiology,¹⁻³ Department of Neurosurgery,^{4,5} Aga Khan University Hospital, Medical Graduate, Karachi Medical and Dental College,⁶ Karachi.

Corresponding Author: Rana Shoaib Hamid. Email: drschoaib@hotmail.com

Abstract

Objective: To analyse inter-observer variation between a neuroradiologist and neurosurgeon in the MRI diagnosis of lumbar nerve root compression. Although lumbar MFI is primarily analyzed and reported by a radiologist, neurosurgeons often analyse it independently as they have sufficient clinical background as well as radiological expertise to diagnose most spinal pathologies on Magnetic Resonance Imaging (MRI).

Methods: Retrospective analysis was carried out for images of 54 patients who underwent MRI between March and July 2010 of lumbar spine with suspected lumbar disc herniation and nerve root compression, at Aga Khan Hospital, Karachi, Pakistan. One fellowship trained neuroradiologist and one neurosurgeon evaluated the images on PACS system separately. Both observers were unaware of the patient's clinical history and each other's findings. Lumbar discs at L3-L4, L4-L5 and L5-S1 levels were evaluated by both observers for disc disease and nerve compression. Findings were recorded on a proforma and analysed with SPSS Version 16.

Results: Total 162 lumbar discs were studied by both readers in 54 patients. Excellent inter-observer agreement was seen for the presence or absence of nerve root compression (Percentage agreement = 88.89%; $k = 0.774$; $p = 0.737$). For disc bulge, inter-observer agreement was fair but statistically insignificant (Percentage agreement = 72.84%; $k = 0.414$; $p = 0.132$). In case of disc herniation, although inter-observer agreement was fair, but the difference was statistically significant (Percentage agreement = 84.57%; $k = 0.511$; $p = 0.002$).

Conclusion: Inter-observer agreement between neuroradiologist and neurosurgeon in diagnosing nerve root compression due to lumbar disc disease was excellent. Agreement regarding disc bulge and herniation was fair.

Keywords: Lumbar disc herniation, Nerve root compression, Neuroradiologist, Neurosurgeon. (JPMA 62: 826; 2012)

Introduction

Lower back pain is the most common symptom-related reason, after common cold and headaches, for clinician visits and accounts for more sick leave and disability than any other single medical condition.^{1,2} The lifetime prevalence of back pain is estimated around 65 to 80%, though a specific diagnosis for back pain cannot be reached in approximately 80% of these cases.^{3,4} Further management of patients with lower back pain, either surgical or conservative, primarily depends upon the presence and severity of nerve root compression.²

For radiological evaluation of lumbar disc disease both MRI and Computerised Tomography (CT) have been used.^{5,6} In a series of comparative radiological studies on the evaluation of lumbar herniated discs, the difference between MRI and CT was not significant.⁷⁻¹⁰ However, for evaluation of nerve root compression, MRI was found to be superior to CT because of its ability to demonstrate excellent soft tissue

details.¹¹ The reported sensitivity and specificity of MRI for detection of nerve root compression is 80.65% and 100% respectively.¹² Since nerve root compression is the most important factor that determines the course of further management, therefore, according to current guidelines, MRI is the first choice investigation and CT should be used as the alternative in the evaluation of the lumbar back if MRI is contraindicated or unavailable.¹³⁻¹⁵

In addition to the presence and severity of nerve root compression assessed radiologically, the management of patients with or without surgery also depends upon patient evaluation for the degree of pain and disability produced by nerve root compression. Although the reporting radiologist has the required radiological and anatomical knowledge, but he is unaware of the patient's actual condition which renders him at a deficit in comparison to the treating neurosurgeon who has the added advantage of knowing the complete clinical picture.

Around the world it is common practice that neurosurgeons analyse MR images independently as they believe that they have sufficient radiological expertise with the additional benefit of being better aware of the patients' clinical picture as compared to the reporting radiologist. Also because backache forms one of the most common causes of referral to neurosurgeons, it becomes essential to know the basic interpretation of a lumbar MRI. Moreover, there exists intra-observer variability between reporting radiologists.¹⁶ Therefore, clinical decision making requires self interpretation of images on neurosurgeon's part. In addition it saves time.

The purpose of the study was to analyse inter-observer variability between a qualified neuro-radiologist and a neurosurgeon in the evaluation of nerve root compression to assess if there exists any discrepancy between neuroradiologists and neurosurgeons in the interpretations of MRI lumbar spine for the diagnosis of this common pathology.

Patients and Methods

This retrospective study was carried out at the Radiology Departments of Aga Khan University Hospital in Karachi. Medical records and images of those patients were retrospectively reviewed who underwent MRI of lumbar spine with suspicion of spinal stenosis from March to July 2010 at the MRI section. All patients who were referred for MRI lumbar spine with clinical suspicion of disc herniation and lumbar radiculopathy were also included. Patients less than 18 years of age, patients with history of surgery, spinal infections or tumours were excluded from the study. The final sample comprised of 54 patients.

All images were acquired with a 1.5 T MRI system (Magnetom Avanto, Siemens Corporation, USA). The standard imaging protocol included T1 and T2 weighted sagittal images (Slice thickness 4 mm, FOV 350x350mm, Image Matrix 672x896), T2 weighted axial images (Slice thickness 4mm, FOV 230x 230mm, Image Matrix 314x448). T2 weighted fat suppressed sagittal images (Slice thickness 4 mm, FOV 350 x 350 mm, Image Matrix 214 x 256).

One fellowship trained neuroradiologist and one neurosurgeon evaluated the images on picture archiving and communication system (PACS). Both observers were blinded to patient's clinical history and each other's findings. The scans were interpreted in sagittal and axial planes. Lumbar discs at L3-L4, L4-L5 and L5-S1 levels were evaluated by both observers. Disc at each level was evaluated for the presence or absence of disc bulge and disc herniation separately. No distinction was made between disc protrusion or extrusion and both were included in the term herniation.

Nerve roots from L3 to S1 levels were also evaluated for nerve compression. For nerve root compression a 5-point scale

was used as proposed by Van Rijn et al¹⁶ 1: No nerve compression, 2: Possibly no root compression, 3: Indeterminate, 4: Possible nerve compression, 5: Definite nerve compression. For the purpose of analyses, these grades of nerve compression were simplified as 'nerve compression' (possibly or definitely) or 'no nerve compression' (all other categories).

Findings of both observers were recorded on a proforma, entered and analysed in SPSS 16. Inter-observer agreement between neuroradiologist and neurosurgeon for evaluation of bulging and herniated discs and nerve root compression was calculated by applying Kappa statistics. Interpretation of Kappa was done as proposed by Cohen¹⁷ i.e. value less than 0.4 was considered poor, value between 0.4 to 0.75 was considered fair to good, and value above 0.75 was considered excellent. Chi-square test was applied to see the association between neuro-radiologist and neurosurgeon in image interpretations. P-value less than 0.05 was considered significant.

Results

Initially 62 patients were included, but 8 were excluded from the study based on the exclusion criteria and the final sample consisted of 54 patients (31 female, 23 male, age range 19-76 years, mean age 46.1 ± 15.1 years). Total 162 lumbar discs were studied by both readers in these 54 patients. For determining presence or absence of disc bulge, the observers agreed in MRI findings in 118 (72.84%) readings and disagreed in 44 (27.16%). Inter-observer agreement for the assessment of disc bulge was fair (k =

Table: Inter-observer agreement in evaluation of disc bulge, herniation and nerve compression.

Disc Bulge				
		Neuroradiologist		
Neurosurgeon	No	Yes		Total
No	82	15		97
Yes	29	36		65
Total	111	51		162
Inter-observer Agreement	(Percentage agreement = 72.84%; (118/162); k = 0.414; p = 0.132)			
Disc Herniation				
		Neuroradiologist		
Neurosurgeon	No	Yes		Total
No	119	24		143
Yes	1	18		19
Total	120	42		162
Inter-observer Agreement	(Percentage agreement = 84.57%; (137/162); k = 0.511; p = 0.002)			
Nerve Root Compression				
		Neuroradiologist		
Neurosurgeon	No	Yes		Total
No	82	7		89
Yes	11	62		73
Total	93	69		162
Inter-observer Agreement	(Percentage agreement = 88.89%; (144/162); k = 0.774; p = 0.737)			

0.414; $p = 0.132$), but statistically insignificant.

For assessment of disc herniation, the observers agreed on imaging findings in 137 (84.57%) cases and disagreed in 25 (15.43%). Inter-observer agreement for the assessment of disc herniation was fair, but clinically significant. ($k = 0.511$; $p = 0.002$).

The agreement between the two observers for the presence or absence of nerve root compression was achieved in 144 (88.89%) readings while they disagreed in 18 (11.11%). Inter-observer agreement was excellent. ($k = 0.774$; $p = 0.737$) (Table).

Discussion

Our study showed substantial disagreement between the observers regarding disc morphology. For disc bulge inter-observer agreement was fair ($k = 0.414$) but statistically insignificant ($p = 0.132$). However, in case of disc herniation although inter-observer agreement was fair ($k = 0.511$), the difference reached had statistical significance ($p = 0.002$). Previous studies regarding inter-observer agreement related to disc morphology showed highly variable results with kappa values between 0.32 and 0.79.¹⁸⁻²² In the study conducted by Lurie J et al. the data showed only fair agreement ($k = 0.32$),¹⁸ Brant-Zawadzki et al. found moderate inter-reader agreement (unweighted $k = 0.59$),¹⁹ Jarvik et al reported moderate to substantial inter-reader agreement with weighted k s of 0.50 to 0.75²⁰ and Solgaard et al and Weishaupt et al found substantial agreement for classifying disc morphology, with inter-observer k s of 0.79 and 0.68, respectively.^{21,22}

Different reasons for this variation between inter-observer variation have been suggested. Brant-Zawadzki et al. comparing two nomenclatures for lumbar herniations concluded that bulging disk was the main reason for moderate agreement.¹⁹ Similar results were seen in study by Van Rijn where it was concluded that more than 50% of inter-observer variation in MRI evaluation of patients with lumbosacral radicular pain is caused by disagreement on bulging disks.¹⁶ Bulging disks usually are assumed to be asymptomatic lesions because they are common in the general asymptomatic population. A patient with a bulging disk and radicular pain is likely to undergo conservative treatment and so this finding has very little clinical significance.

Analysing our data, we found that the neurosurgeon had labelled considerably fewer discs as herniated as compared to the radiologist i.e. 19 as compared to 42 by the radiologist, who had labeled 65 discs as bulging as compared to the radiologist's 51. Despite this difference, the agreement regarding nerve root compression is excellent (Percentage agreement for nerve root compression= 88.89 %; $k = 0.774$; $p = 0.737$) and is comparable to a study earlier conducted by

Van Rijn despite the fact that in his study the inter-observer agreement was calculated between two neuroradiologists. Van Rijn reported a Kappa value of 0.77 for inter-observer agreement for nerve root compression, which is almost similar to our study.¹⁶

A probable explanation of this apparent discrepancy may be that since it is the nerve compression that is significant from surgical point of view, the neurosurgeon considered those herniations as bulges or normal which were not resulting in nerve root compression since no surgical treatment was required in such cases. The difference may have academic importance, but from surgical perspective it is unlikely to alter the course of management.

This disagreement, however, further emphasises the point made in earlier studies that increased effort is required by lumbar MRI readers to use terms regarding disc morphology in accordance with published guidelines. In this respect the Combined Task Force of the North American Spine Society, the American Society of Spine Radiology, and the American Society of Neuroradiology have issued guidelines that provide standardisation of terms to characterise disc herniation, as well as other disc pathologies.²³ However, the degree of it being followed across the relevant specialties appears questionable.

Another cause of this disagreement may be the lack of supervised training of neurosurgeons regarding the interpretation of lumbar MRI. This weakness may be overcome by arranging supervised training sessions of neurosurgeons in radiology by neuroradiologists. This would improve their understanding of lumbar MRI and enable them to be able to interpret lumbar MRI accurately.

Conclusion

Excellent inter-observer agreement was seen between the neuroradiologist and the neurosurgeon in diagnosing nerve root compression due to lumbar disc disease. Agreement regarding disc bulge and herniation was fair. Although this is clinically insignificant, but we suggest arranging supervised training sessions of neurosurgeons in radiology by neuroradiologists for their better understanding and interpretation of lumbar MRI.

References

1. Carrino JA, Lurie JD, Tosteson AN, Tosteson TD, Carragee EJ, Kaiser J. Lumbar spine: reliability of MR imaging findings. *Radiology* 2009; 250: 161-70.
2. Jarvik JG, Deyo RA. Diagnostic evaluation of low back pain with emphasis on imaging. *Ann Intern Med* 2002; 137: 586-97.
3. Breslau J, Seidenwurm D. Socioeconomic aspects of spinal imaging: impact of radiological diagnosis on lumbar spine-related disability. *Top Magn Reson Imaging* 2000; 11: 218-23.
4. Brant-Zawadzki MN, Dennis SC, Gade GF, Weinstein MP. Low back pain. *Radiology* 2000; 217: 321-30.
5. Siddiqui AH, Rafique MZ, Ahmad MN, Usman MU. Role of magnetic

- resonance imaging in lumbar spondylosis. *J Coll Physicians Surg Pak* 2005; 15: 396-9.
6. Thornbury JR, Fryback DG, Turски PA, Javid MJ, McDonald JV, Beinlich BR, et al. Disk-caused nerve compression in patients with acute low-back pain: diagnosis with MR, CT myelography and plain CT. *Radiology* 1993; 186: 731-8.
 7. Goscinski I, Ulatowski S, Urbanik A. [Comparison of the clinical usefulness of magnetic resonance (MR), computer tomography (CT) and radiculography (R) in diagnostic lumbar discopathy]. *Przegl Lek* 2001; 58: 885-8.
 8. Janssen ME, Bertrand SL, Joe C, Levine MI. Lumbar herniated disk disease: comparison of MRI, myelography, and postmyelographic CT scan with surgical findings. *Orthopedics* 1994; 17: 121-7.
 9. Albeck MJ, Hilden J, Kjeer L, Praestholm J, Hensiksen O, Gjerris F. A controlled comparison of myelography, computed tomography and magnetic resonance imaging in clinically suspected lumbar disc herniation. *Spine* 1995; 20: 443-8.
 10. Tullberg T, Grane P, Rydberg J, Isacson J. Comparison of contrast-enhanced computed tomography and gadolinium enhanced magnetic resonance imaging one year after lumbar discectomy. *Spine* 1994; 19: 183-8.
 11. van Rijn JC, Klemetsö N, Reitsma JB, Bossuyt PM, Hulsmans FJ, Peul WC, et al. Observer variation in the evaluation of lumbar herniated discs and root compression: spiral CT compared with MRI. *Br J Radiol* 2006; 79: 372-7.
 12. Chawalparit O, Churojana A, Chiewvit P, Thanapipatsir S, Vamvanij V, Charnchaowanish P. The limited protocol MRI in diagnosis of lumbar disc herniation. *J Med Assoc Thai* 2006; 89: 182-9.
 13. Patel N. Surgical disorders of the thoracic and lumbar spine: a guide for neurologists. *J Neurol Neurosurg Psychiatry* 2002; 73 (Suppl. 1): i42-8.
 14. Milette PC. Classification, diagnostic imaging, and imaging characterization of a lumbar herniated disk. *Radiol Clin North Am* 2000; 38: 1267-92.
 15. Herzog RJ. The radiologic assessment for a lumbar disc herniation. *Spine*. 1996; 21: 19S-38S.
 16. van Rijn JC, Klemetsö N, Reitsma JB, Majoie CB, Hulsmans FJ, Peul WC, et al. Observer variation in MRI evaluation of patients suspected of lumbar disk herniation. *AJR Am J Roentgenol* 2005; 184: 299-303.
 17. Gwet K. Inter-Rater Reliability: Dependency on Trait Prevalence and Marginal Homogeneity. *Statistical Methods For Inter-Rater Reliability Assessment* 2; 2002.
 18. Lurie JD, Doman DM, Spratt KF, Tosteson AN, Weinstein JN. Magnetic resonance imaging interpretation in patients with symptomatic lumbar spine disc herniations: comparison of clinician and radiologist readings. *Spine (Phila Pa 1976)*. 2009; 34: 701-5.
 19. Brant-Zawadzki MN, Jensen MC, Obuchowski N, Ross JS, Modic MT. Interobserver and intraobserver variability in interpretation of lumbar disc abnormalities. A comparison of two nomenclatures. *Spine* 1995; 20: 1257-63.
 20. Jarvik JG, Haynor DR, Koepsell TD, Bronstein A, Ashley D, Deyo RA. Interreader reliability for a new classification of lumbar disk disease. *Acad Radiol* 1996; 3: 537-44.
 21. Solgaard Sorensen J, Kjaer P, Jensen ST, Anderson P. Low-field magnetic resonance imaging of the lumbar spine: reliability of qualitative evaluation of disc and muscle parameters. *Acta Radiol* 2006; 47: 947-53.
 22. Weishaupt D, Zanetti M, Hodler J, Boos N. MR imaging of the lumbar spine: prevalence of intervertebral disk extrusion and sequestration, nerve root compression, end plate abnormalities, and osteoarthritis of the facet joints in asymptomatic volunteers. *Radiology* 1998; 209: 661-6.
 23. Fardon DF, Milette PC; Combined task Forces of the North American Spine Society, American Society of Spine Radiology and American Society of Neuroradiology. Nomenclature and classification of lumbar disc pathology. Recommendations of the Combined task Forces of the North American Spine Society, American Society of Spine Radiology and American Society of Neuroradiology. *Spine* 2001; 26: E93-113.
-