



THE AGA KHAN UNIVERSITY

eCommons@AKU

---

Department of Anaesthesia

Medical College, Pakistan

---

7-2020

## In response to letter title "immediate hemodynamic and gaseous exchange effect of bi-level positive airway pressure after cardiac surgery: Our insight to Hamid et al.'s study"

Mohammad Hamid

*Aga Khan University*, mohammad.hamid@aku.edu

Mohammad Irfan Akhtar

*Aga Khan University*, mohammad.irfan@aku.edu

Saba Ahmed

*Aga Khan University*

Follow this and additional works at: [https://ecommons.aku.edu/pakistan\\_fhs\\_mc\\_anaesth](https://ecommons.aku.edu/pakistan_fhs_mc_anaesth)



Part of the [Anesthesiology Commons](#), and the [Cardiology Commons](#)

---

### Recommended Citation

Hamid, M., Akhtar, M., Ahmed, S. (2020). In response to letter title "immediate hemodynamic and gaseous exchange effect of bi-level positive airway pressure after cardiac surgery: Our insight to Hamid et al.'s study". *Annals of Cardiac Anaesthesia*, 23(3), 373-374.

Available at: [https://ecommons.aku.edu/pakistan\\_fhs\\_mc\\_anaesth/399](https://ecommons.aku.edu/pakistan_fhs_mc_anaesth/399)

## In Response to Letter Title “Immediate Hemodynamic and Gaseous Exchange Effect of Bi-Level Positive Airway Pressure after Cardiac Surgery: Our Insight to Hamid *et al.*’s Study”

The Editor,

Thank you for the opportunity to respond to the letter by Karim *et al.*,<sup>[1]</sup> in reference to our study.<sup>[2]</sup> The authors in this letter have expressed few concerns regarding our study and I would like to respond in the same sequence.

First critique is that we only used Bilevel Positive Airway Pressure (BiPAP) in nonhypercapnoeic patients. It is erroneous, as we also included hypercapnoeic patients. In our study eight patients had PaCO<sub>2</sub> 50 mm Hg or greater. The mean value of 44 mm Hg in pre-BiPAP period probably gives a false impression that these patients were excluded.

About echocardiographic evaluation of cardiac function before and after applying a BiPAP, I think it is difficult as you know that once you decide about BiPAP application it should be done quickly to prevent decompensation. I was unable to find any study where ECHO was used in such situation. It is a good idea but may be impractical in this scenario. Hoffman *et al.*<sup>[3]</sup> also used PA catheter readings for cardiac index and cardiac output rather than relying on echocardiography. Their study showed improved cardiac function after BiPAP and this effect was also present in our study. Another question regarding pulmonary HTN, we did not find pulmonary HTN in preoperative evaluation and second it would have made no difference as we compared the pre and post BiPAP hemodynamics.

Second, about comorbidities such as COPD and sleep apnea, we were unable to find COPD and sleep apnea complaints during preoperative evaluation. These questions were asked specifically and past medical record was checked for these conditions.

Regarding pain and sedative medications influencing parameters. During the short study period, no one was given sedatives or pain medications so I do not think it is an issue here. Generally, we avoid sedatives in all postoperative patients and even analgesics, which we use have minimal effect.

About the effect of BiPAP on reintubation rate, there are several studies mentioned that BiPAP application reduces reintubation rate in respiratory

failure patients.<sup>[4,5]</sup> I agree with the authors that the prophylactic use of BiPAP is controversial. They mentioned Ađirođlu G *et al.*<sup>[6]</sup> study where BiPAP was used prophylactically in nonrespiratory distress patients. It was used twice for 20 min with 3 h interval while we used different criteria for included patients and duration of application was different as well. We firmly believe that BiPAP application does help the patient in respiratory distress after cardiac surgery and reduces reintubation rate.

### Financial support and sponsorship

Nil.

### Conflicts of interest

There are no conflicts of interest.

**Mohammad Hamid, Mohammad I Akhtar,  
Saba Ahmed**

*Department of Anaesthesia, Aga Khan University,  
Karachi, Pakistan*

*Address for correspondence: Dr. Mohammad Hamid,  
Associate Professor, Department of Anaesthesia,  
Second Floor PW II, Aga Khan University, Stadium Road,  
Karachi, Sindh, Pakistan.  
E-mail: mohammad.hamid@aku.edu*

**Submitted:** 27-Mar-2020


**Accepted:** 09-May-2020

**Published:** 17-Jul-2020

### References

1. Karim HMR, Gonalves G, Esquinas AM. Immediate hemodynamic and gaseous exchange effect of Bi-level positive airway pressure after cardiac surgery: Our insight to Hamid *et al.*’s study. *Ann Card Anaesth* 2020;23:372.
2. Hamid M, Akhtar MI, Ahmed S. Immediate changes in hemodynamics and gas exchange after initiation of noninvasive ventilation in cardiac surgical patients. *Ann Card Anaesth* 2020;23:59-64.
3. Hoffmann B, Jepsen M, Hachenberg T, Huth C, Welte T. Cardiopulmonary effects of non-invasive positive pressure ventilation (NPPV)--A controlled, prospective study. *Thorac Cardiovasc Surg* 2003;51:142-6.
4. Liu YJ, Zhao J, Tang H. Non-invasive ventilation in acute respiratory failure: A meta-analysis. *Clin Med (Lond)* 2016;16:514-23.
5. Olper L, Corbetta D, Cabrini L, Landoni G, Zangrillo A. Effects of non-invasive ventilation on reintubation rate: A systematic review and meta-analysis of randomised studies of patients undergoing cardiothoracic surgery. *Crit Care Resusc* 2013;15:220-7.

6. Ağiroğlu G, Baysal A, Copuroğlu E, Gül Y, Karamustafaoglu Y, Dogukan M. Does early use of Bilevel Positive Airway Pressure (BiPAP) in cardiothoracic intensive care unit prevent reintubation? *Int J Clin Exp Med* 2014;7:3439-46. eCollection 2014.

Access this article online	
<b>Quick Response Code:</b> 	<b>Website:</b> <a href="http://www.annals.in">www.annals.in</a>
	<b>DOI:</b> 10.4103/aca.ACA_57_20

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

**How to cite this article:** Hamid M, Akhtar MI, Ahmed S. In response to letter title "Immediate hemodynamic and gaseous exchange effect of bi-level positive airway pressure after cardiac surgery: Our insight to Hamid *et al.*'s study". *Ann Card Anaesth* 2020;23:373-4.  
 © 2020 Annals of Cardiac Anaesthesia | Published by Wolters Kluwer - Medknow