



THE AGA KHAN UNIVERSITY

eCommons@AKU

Community Health Sciences

Department of Community Health Sciences

June 2002

Anorexic behaviour and attitudes among female medical and nursing students at a private university hospital

N Babar
Aga Khan University

M Alam
Aga Khan University

S S. Ali
Aga Khan University

A Ansari
Aga Khan University

M Atiq
Aga Khan University

See next page for additional authors

Follow this and additional works at: https://ecommons.aku.edu/pakistan_fhs_mc_chs_chs

Recommended Citation

Babar, N., Alam, M., Ali, S. S., Ansari, A., Atiq, M., Awais, A., Amin, F., Israr, S. M. (2002). Anorexic behaviour and attitudes among female medical and nursing students at a private university hospital. *Journal of Pakistan Medical Association*, 52(6), 272-276.
Available at: https://ecommons.aku.edu/pakistan_fhs_mc_chs_chs/399

Authors

N Babar, M Alam, S S. Ali, A Ansari, M Atiq, A Awais, F Amin, and S M. Israr

Anorexic Behaviour and Attitudes Among Female Medical and Nursing Students at a private University Hospital

N.Babar, M.Alam, S.S.Ali, A.Ansari, M.Atiq, A.Awais, F.Amin (Fourth Year Medical Students, The Aga Khan University Hospital, Karachi.)

S.M.Israr (Department of Community Health Sciences, The Aga Khan University Hospital, Karachi.)

Abstract

Objective: To Study the anorexic behaviour and attitude among female medical and nursing students in a tertiary care hospital.

Subjects and Methods: A cross-sectional survey to determine the proportion of anorexic behaviour among female medical and nursing students at The Aga Khan University Hospital Karachi was conducted. A multi-staged sampling technique was utilized in which our study population was first divided according to field of enrollment (medical or nursing school) and then stratified by class. Within each stratum, we used convenience sampling due to time and availability constraints. We utilized a modified Eating Attitudes Test (EAT-26) to collect data. The Eating Attitude Test is probably the most widely used standardized measure of symptoms characteristic of eating disorders, including anorexia nervosa.

Results: A total of 180 female students were interviewed. Ninety-four (52.2%) were medical students and 86 (47.8%) were nursing students. Among the 94 medical students, 26 (27.7%) were from first year, 14 (14.9%) from second year, 15 (16.0%) from third year, 20 (21.3%) from fourth year and 19 (20.2%) from the final year of medical college. Among the 86 nursing students 34 (39.5%) were from first year, 23 (26.7%) from second year, 27 (31.4%) from third year and 2 (2.4%) from fourth year. The proportion of anorexic behaviour among medical students was 8.0%, 7.1% and 20.0% in first, third and fourth years respectively. No individuals with anorexic behaviour were found in second and fifth years. The total proportion of anorexic behaviour among female health care students was 21.7%.

Conclusion: Results showed a 21.7% prevalence of anorexic behaviour, a figure much higher than that reported in similar studies conducted in Asia. We also found that the proportion was much higher among female nursing students as compared to female medical students. A previous visit to a psychiatrist for reasons other than eating disorders was found to be associated with anorexic behaviour (JPMA 52: 272; 2002).

Introduction

Anorexic behaviour causes significant physical and psychological stress in a large number of adolescents and young adults throughout the world. This clinical condition involves symptoms such as binge eating, purging and a constant preoccupation with food and body image. Individuals may engage in severe dieting behaviour, experience extreme guilt after eating and feel terrified of being overweight. They may actually feel as though food controls their lives.

Anorexic behaviour has been found in multiple studies to precede or actually be an indicator of anorexia nervosa, an eating disorder involving weight loss severe enough to cause significant

physiologic consequences. Subsequent malnutrition leads to protein and nutritional deficiencies and disruption of multiple organ systems, with mortality rates ranging from 5-10%¹. Early recognition of anorexia nervosa has favorable implications on the clinical outcome of the disease. For this reason identifying anorexic behaviour is a vital exercise if the burden of anorexia nervosa is to be reduced.

Research has disclosed a significant cultural effect on the pattern and distribution of the eating disorders. An important determinant appears to be the Western premium of thinness as an esthetic and moral value²⁻⁴. A high prevalence of eating disorders has been reported from many non-Western societies and several ethnic minorities in the United States,⁷ but their presenting features often differ somewhat from the DSM-IV criteria for anorexia nervosa and bulimia nervosa². With the advent of satellite television and growing Western trends in our society, we hypothesized that we might establish results comparable with Western data, indicating that anorexic behaviour poses a significant burden on our population as well.

Subjects and Methods

We conducted a cross-sectional survey to determine the prevalence of anorexic behaviour among female medical and nursing students at The Aga Khan University Hospital Karachi. Our study population included 509 female students from the Medical College and the School of Nursing. Out of 509 students, 185 were enrolled at the Medical College while the remainder belonged to the School of Nursing. A number of the students reside in Karachi and commute to and from the University every day, while the remainder, with families settled in other cities of Pakistan as well as abroad, live at the University Hostel.

We assumed the prevalence of our study variable, anorexic behavior, to be 12% as depicted by Martinez et al. who utilized the EAT-26 survey to identify anorexic behaviour among high school girls⁵. With a confidence interval of 95%, a sample size of 168 was calculated. Adding an extra 10% to counter the biases, refusals and wrong answers, we aimed to achieve a total sample size of 185 but only 180 survey forms (duly filled in) were obtained.

We utilized a multi-staged sampling technique in which the study population was first divided into two clusters according to their school (medical or nursing) and then stratified by class. Thus we collected our sample data from nine strata; five from the Medical College (first year through fifth year) and four from the School of Nursing (first year through fourth year). Within each stratum, we used convenience sampling due to time and availability constraints.

Data was collected using a modified version of the Eating Attitudes Test (EAT-26)⁵. EAT-26 has not been established as the sole mean of identifying eating disorders, but studies have shown that it can be used as a screening instrument in part of a two-stage process in which individuals who score above 20 are evaluated for eating disorders in a diagnostic interview. In this manner, early identification of an eating disorder can lead to earlier treatment thereby reducing morbidity and mortality.

The EAT-26 survey comprised of three sections (A, B, and C). Section A dealt with demographics, Section B contained 26 questions relating to eating attitudes and was used to calculate the score by addition of the values of the responses. Questions assessed various aspects of distorted eating behaviour, including bulimic tendencies, body image perception and degree of willful control over eating behavior. A score above 20 for Section B was considered diagnostic of anorexic behaviour. Section C consisted of specific questions regarding weight, height, psychiatric issues and menstrual irregularities. For the purpose of our study, we modified Section A so that the questions

relating to demographics pertained to our study population. We also added a Section D that dealt with factors we hypothesized were associated with anorexic behaviour in our target population. These included: living in the hostel, having friends who went on diets, stress related reduction in food intake, influence of media on the way students desired to look, pressure from parents or siblings to lose weight, the presence of family conflicts, having feelings of self-worthlessness or guilt and having been to a psychiatrist for reasons other than eating behaviour.

Results

A total of 180 female students were interviewed. Ninety-four (52.2%) were medical students and 86 (47.8%) nursing students.




Figure 1. Proportions of various strata (n = 180).

Figure 1 shows the number of medical and nursing students included in the study. Their mean age was 20.86 years with a standard deviation of 2.12 and a range from 18 to 32. The mean age of medical students was 21 ± 2.08 and that of nursing students was 20.6 ± 2.20 years.

The eating behaviour and attitudes of students were assessed by the use of the Eating Attitude Test-26. Of 180 females, 56 (31.1%) resided in the hostel and 118 (65.6%) were day-scholars. Among them 123 (68.3%) reported that their friends were dieting' at the time of the interview. Eighty-nine (49.4%) students thought that their food intake decreased during examinations or other periods of stress.

One hundred and nine (60.6%) agreed that media influenced the way they desired to look. The parents or siblings of 62 (34.4%) students had advised them to lose weight. Forty-two (23.3%) students had family conflicts and 57 (31.7%) had feelings of self-worthlessness or guilt. Thirteen

(7.2%) students had history of a psychiatrist consultation for reasons other than eating behaviour. Non-anorexic behaviour was assumed if the EAT-26 score was less than 20. There were 141 (78.33%) cases of non-anorexic behaviour. The mean age in this group was 20.95 with standard deviation of 2.23 and a range from 18 to 32. The mean EAT-26 score was 7.60 with a standard deviation of 5.00 and a range from zero to 19. The BMI was available in 119 students and the mean BMI was 20.83 with a standard deviation of 2.88 and a range from 16.09 to 34.89. The desired BMI's of only 110 students were available. The mean desired BMI was 20.01 with a standard deviation of 3.25 and a range from 17.04 to 29.38.

A case was labeled as having anorexic behaviour if EAT-26 score was 20 or more. Of the total 180 students sampled, 39 (21.67%) students had anorexic behaviour. The mean age in this group was 20.54 with standard deviation of 1.65 and range from 18 to 27. The mean EAT-26 score was 32.25 with a standard deviation of 8.53 and a range from 20 to 53. The BMI was available in 37 students. The mean BMI was 22.30 with a standard deviation of 4.17 and a range from 17.07 to 39.06. We were able to calculate the desired BMI of 31 students and the mean was 20.89 with a standard deviation of 2.53 and a range from 17.58 to 28.65.

On the basis of high scoring in the anorexic group, we identified the top six questions from the EAT-26 questionnaire, namely: am terrified about being overweight (84.6%), am aware of the calorie content of the foods I eat (87.2%), am preoccupied with the desire to be thinner (79.5%), think of burning calories when I exercise (89.7%), am preoccupied with the thought of having fat on my body (76.9%) and display self-control around food (76.9%) as shown in Figure 2.




Figure 2. Questions with maximum positive response in students with anorexic behavior (n = 180).

A Chi-square test was applied to all of the factors we hypothesized to be associated with anorexic behaviour. These factors included living in the hostel, having friends who went on diets, stress related reduction in food intake, influence of media on the way one desired to look, pressure from parents or siblings to lose weight, the presence of family conflicts, having feelings of self-worthlessness or guilt, and having been to a psychiatrist for reasons other than eating behavior. Of all these factors, a statistically significant value was found only for the last mentioned risk factor, that is, consulting a psychiatrist for reasons other than eating behavior ($X^2=8.142$, $df=1$). In order to determine the relevance of the relationship between current and desired BMI and anorexic status, a t-test of independent sample was carried out. It was found that the current BMI of those with anorexic behaviour was statistically different from those without anorexic behaviour ($p < 0.05$). However, this did not hold true for desired BMI.

Discussion

Anorexic behaviour, which precedes anorexia nervosa, has been studied by various centers in an effort to recognize anorexia nervosa in its preliminary stages. The premorbid condition comprises of the attitudes towards eating seen in anorexia nervosa as well as other psychiatric symptoms. These include markers such as low self-esteem, depression, identity conflicts and high anxiety levels.

Of the total 180 female students included in our study, 21.7% screened positive for anorexic behaviour with a cut-off value of 20 on the EAT-26 scale. A look at Western data shows that our result exceeds the figures they have obtained. Anstine et al. showed that 17.0% of college freshman women residing in dormitories at the University of Florida had anorexic behavior⁶. Chandarana et al. conducted a survey among high school students in London using a demographic questionnaire, EAT-26 and Basic Personality Inventory⁷. With a cut-off score of 20 on EAT-26, 7.5% of the sample size had anorexic - behaviour. Martinez et al. carried out a comprehensive study on anorexic behaviour of high school girls in which results showed anorexic behaviour to be present in 12.8% of the study population⁸.

Interestingly, Sjostedt et al showed that Indian women scored higher on EAT-26 as compared to Australian women⁹. In Iran, Nobakht et al. targeted school girls and found a lifetime prevalence of 0.9% for anorexia nervosa, 3.2% for bulimia nervosa and 6.6% for the partial syndrome⁵. Our results show a frequency of anorexic behaviour greater than all of the above-mentioned reports. This reiterates the fact that anorexic behaviour is not an entity, which could solely be attributed to the Western society; the lack of data from the developing world, however, may have left this false impression.

One interesting feature in our study was the comparison between medical and nursing students. We hypothesized that since both groups of students belonged to nearly same age group (mean age 20.8 ± 2.12), there would be no significant difference between their eating behaviour and attitude. However, we found a significant difference between the two groups with regards to the proportion of anorexic behavior ($\chi^2=23.44$, $df=1$). Any suitable explanation for this difference, however, could not be generated.

Lanzi et al pointed out an association between distorted attitudes towards weight, body and food with disturbances of body image among students of Pavia, Italy¹⁰. They also showed a correlation between abnormal eating behaviours and a low level of self-esteem, suicide attempts, and drug abuse. In our study, consulting the psychiatrist for reasons other than eating disorders was the only factor shown to be significantly associated with anorexic behaviour ($\chi^2=8.142$, $df=1$). This indicates that individuals with anorexic behaviour may have other psychiatric complaints, that need to be evaluated. Perhaps the presence of certain psychiatric illnesses predisposes one towards the development of anorexic behaviour. However, our questionnaire could not evaluate this relationship effectively.

The current BMI of students with anorexic behaviour was significantly different from those without anorexic behaviour ($p < 0.05$). However, this did not hold true for the desired BMI. The difference in current BMI could be accounted for by the dietary habits of students with anorexic behaviour. Since there was no significant difference in the desired BMI of the anorexic and the nonanorexic groups, we can infer that the two groups have the same perception of the ideal body image.

With reference to the EAT-26 questionnaire, we identified the six highest scoring questions among those with anorexic behaviour. Similar trends have been documented by Maloney et al., who conducted a survey among school going children between grades 3 through 6 in Ohio¹¹. He

showed that 45% of the children wanted to be thinner, 37% had already tried to lose weight, and 6.9% scored in the anorexia nervosa range.

Limitations

Our sample was drawn from one medical university due to the limited time available to conduct our study. Thus, our results may not be truly representative of the female healthcare student population. Certain subsets of students in our target population were unavailable due to electives or vacations, so their surveys were administered to students from other classes. For this reason, the number of students sampled from each strata of our population is not exactly proportionate to the study population.

Conclusion

Anorexic behaviour emerged as an important premorbid entity in our defined population. A 21.7% prevalence of anorexic behaviour is higher than that found in similar studies conducted elsewhere. We also found that the proportion was much higher among female nursing students as compared to female medical students. A previous visit to a psychiatrist for reasons other than eating disorders was found to be associated with anorexic behaviour. The target population for our study belonged to a selected stratum of the society (high socio-economic class), so the results cannot be extrapolated to the general population.

Recommendations

Measures need to be taken to identify this premorbid condition to avoid undesirable outcomes. Other psychiatric illnesses that are more relevant to our society i.e. major depression and anxiety disorders should be evaluated in context of anorexic behaviour and should there be a causal relationship then it must be duly addressed. Anorexic behaviour might just be indicative of underlying or future anorexia nervosa and a cohort study should be done to illuminate absolute and relative risk associated with such anorexic behaviour. Programs should be implemented to increase awareness regarding necessary daily caloric intake, impact of undue dieting and exercise and normalcy of body shape and weight. Media is one way to realize these educational goals.

Acknowledgements

We are grateful for the kind assistance extended to us by Mr. Hamza A. Rasheed and Dr. Ajmal Agha.

References

1. Gull WW. Anorexia Nervosa (apepsia hysterics, anorexia hysterica). Trans. Clin. Soc. London, 1874;7:22-28.
2. Steinhausen HC, Rauss-Mason C, Seidel R. Follow-up studies of anorexia nervosa: A review of four decades of outcome research. Psychol. Med., 1991; 21:447.
3. Herzog DB, Nussbaum KM, Marmor AK: Comorbidity and outcome in eating disorders. Psychiatr. Clin. North. Am., 1996; 19: 843-59.

4. Herzog DB, Keller MB, Lavori PW. Outcome in anorexia nervosa and bulimia nervosa: A review of the literature. *i. Nerv. Ment. Dis.*, 1988; 176:131.
5. Nobakht M, Dezhkam M: An epidemiological study of eating disorders in Iran. *Int J. Eat. Disord.*, 2000; 28:265-71.
6. Anstine D, Grinenko D. Rapid screening for disordered eating in college-aged females in the primary care setting. *J. Adolesc. Health*, 2000 ;26:338-42.
7. Chandarana P, Helmes E, Benson N. Eating attitudes as related to demographic and personality characteristics: a high school survey. *Can. J. Psychiatry*, 1988; 33:834-37.
8. Martinez MA, Menendez D, Sanchez TMJ, et al. *Aten Primaria Int. J. Eat. Disord.*, 2000; 25:313-19.
9. Sjostedt JP, Schumaker JF, Nathawat SS. Eating disorders among Indian and Australian university students. *J. Soc. Psychol.*, 1998;138:351-7.
10. Lanzi G, Rossi G, Balottin U, et al. Eating behaviour disorders: epidemiologic study of 434 adolescents. *Minerva Pediatr.*, 1997; 49: 249-60.
11. Maloney MJ, McGuire J, Daniels SR, et al. Dieting behaviour and eating attitudes in children *Pediatrics* 1990; 85; 714.