



THE AGA KHAN UNIVERSITY

eCommons@AKU

---

Department of Paediatrics and Child Health

Division of Woman and Child Health

---

10-21-2006

## Child health and survival in the eastern mediterranean region

Zulfiqar A. Bhutta

*Aga Khan University, zulfiqar.bhutta@aku.edu*

Asim Belgaumi

*King Faisal Specialist Hospital and Research Center, Riyadh, Saudi Arabia, asim.belgaumi@aku.edu*

Rab Mohammad Abdur

*Research Policy and Cooperation Unit, WHO Eastern Mediterranean Region, Cairo, Egypt*

Zein Karrar

*School of Medicine, Khartoum, Sudan*

Mohamed Khashaba

*Mansourah Children's Hospital, Mansourah, Egypt*

*See next page for additional authors*

Follow this and additional works at: <https://ecommons.aku.edu/>

[pakistan\\_fhs\\_mc\\_women\\_childhealth\\_paediatr](https://ecommons.aku.edu/pakistan_fhs_mc_women_childhealth_paediatr)



Part of the [Hematology Commons](#), [Oncology Commons](#), and the [Pediatrics Commons](#)

---

### Recommended Citation

Bhutta, Z. A., Belgaumi, A., Abdur, R. M., Karrar, Z., Khashaba, M., Mouane, N. (2006). Child health and survival in the eastern mediterranean region. *BMJ*, 333(7573), 839-842.

Available at: [https://ecommons.aku.edu/pakistan\\_fhs\\_mc\\_women\\_childhealth\\_paediatr/427](https://ecommons.aku.edu/pakistan_fhs_mc_women_childhealth_paediatr/427)

---

**Authors**

Zulfiqar A. Bhutta, Asim Belgaumi, Rab Mohammad Abdur, Zein Karrar, Mohamed Khashaba, and Nezha Mouane

## The way forward

Naturally, health professionals are expected to have a central role in promoting reform through actions to improve health. Such a role is fundamentally different from the roles demanded of health professionals in countries that have undergone political reform, such as South Africa and Eastern European countries<sup>19-20</sup> where the focus has been on monitoring of health effects of democratisation. Although priorities for health actions depend on the country and context, health professionals may consider three broad strategies:

**Mobilising health professionals**—Acknowledging our current disengagement and historical responsibilities towards the current situation and rallying to the cause of reform are prerequisites to engaging the wider public. This will determine our credibility

**Mobilising the public**—By engaging the public, civil institutions, opposition politicians, and even government agencies in discussions and projects, health professionals will be able to highlight the social and political determinants of health

**Health system actions**—We become more effective advocates for reform if we are successful at reforming our hospitals, clinics, and health ministries. Democratising structures, undoing hierarchies, and advancing transparency and accountability in our institutions will contribute to wider reforms.

The box gives some specific actions corresponding to these strategies. These are not meant to be prescriptive but to stimulate debate. Regional collaborations will prove helpful in advancing the cause of reform through drawing on rich perspectives and experiences from different settings. We need wide contributions and research from countries and regions to inform action leading to change.

We thank Dean Huda Zurayk for her pivotal role in creating a stimulating intellectual environment at the faculty of health sciences at the American University of Beirut and for her uncompromising engagement with ideas on health and social concerns.

Competing interests: None declared.

Contributors and sources: SJ is a cardiologist with a research interest in health and social change in the Arab world. AE-Z has a research interest in health and environmental policy, IN conducts research into occupational health in developing countries, and RG researches health as a social construction and population health in situations of conflict. The ideas in this article arose from discussions among all four of us during 2002-4 while we were visiting or working at the American University of Beirut.

- 1 United Nations Development Programme, Arab Fund for Economic and Social Development. *Arab human development report 2004*. New York: UN, 2005.
- 2 Ayodeji Akala F, El-Saharty S. Public health challenges in the Middle East and North Africa. *Lancet* 2006;367:961-4.
- 3 WHO Eastern Mediterranean Regional Office. *Country profiles*. [www.emro.who.int/emrinfo](http://www.emro.who.int/emrinfo) (accessed 5 Sep 2006).
- 4 Jabbour S. Health and development in the Arab world: Which way forward? *BMJ* 2003;326:1141-3.
- 5 Fouad FM, Jabbour S. New concepts in health contradict interests of medical "clergy". *An-Nahar Daily* 2004 Jun 13.
- 6 Giacaman G. In the throes of Oslo: Palestinian society, civil society, and the future. In: Giacaman G, Lonning D, eds. *After Oslo: new realities, old problems*. London: Pluto Press, 1998:1-51.
- 7 Ammar W. *Health system and reform in Lebanon*. Beirut: World Health Organization, 2003.
- 8 Riska E. Towards gender balance: but will women physicians have an impact on medicine? *Soc Sci Med* 2001;52:179-87.
- 9 Zuhur S. Women and empowerment in the Arab World. *Arab Stud Q* 2003;25(4):17-38.
- 10 Barghouthi M, Giacaman R. The emergence of an infrastructure of resistance. The case of health. In: Nassar JR, Heacock R, eds. *Intifada. Palestine at the crossroads*. New York: Praeger, 1990:73-87.
- 11 Giacaman R. A community of citizens: disability rehabilitation in the Palestinian transition to statehood. *Disabil Rehabil* 2001;23:639-44.
- 12 Reproductive Health Working Group. *What is the Reproductive Health Working Group?* [www.rhwg.org](http://www.rhwg.org) (accessed 25 Sep 2006).
- 13 Younis N, Khattab H, Zurayk H, El-Mouelhy M, Main MF, Farag AM. A community study of gynecological and related morbidities in rural Egypt. *Stud Fam Plann* 1993;24:175-86.
- 14 Choices and Challenges in Changing Childbirth Research Network. Routines in facility-based maternity care: evidence from the Arab World. *BJOG* 2005;112:1270-6.
- 15 Joseph S. Patriarchy and development. *Gender Development* 1996;4:14-9.
- 16 Hariri Foundation Directorate of Health and Social Services. *Harrimed*. [www.haririmed.org](http://www.haririmed.org) (accessed 7 Sep 2006).
- 17 Islamic Health Society. <http://www.hayaa.net> (accessed 23 Sep 2006).
- 18 Ministry of Health. *Healthy villages programme in Syria*. [www.moh.gov.sy/arabic/health-services/hvillage.htm](http://www.moh.gov.sy/arabic/health-services/hvillage.htm) (accessed 1 May 2006).
- 19 McKee M, Nolte E. Lessons from health during the transition from communism. *BMJ* 2004;329:1428-9.
- 20 Ncayiyana DJ. Is democracy good for people's health? A South African perspective. *BMJ* 2004;329:1425-6.

doi 10.1136/bmj.38980.664074.94

## Child health and survival in the Eastern Mediterranean region

Zulfiqar A Bhutta, Asim Belgaumi, Mohammad Abdur Rab, Zein Karrar, Mohamed Khashaba, Nezha Mouane

Most child deaths in the region are preventable and occur in just a few of the 22 countries in the region. The interventions are not expensive, but governments need to implement them

Over 10 million children aged under 5 years die every year, almost 90% of them in a few countries in sub-Saharan Africa and South Asia.<sup>1-6</sup> Landmark series on child and neonatal survival suggested that this high mortality persists despite low cost solutions being known and that almost 60-70% of these deaths could be prevented by making these interventions widely available.<sup>2-4</sup> To evaluate whether countries have acted on this information, we did an in-depth assessment of child health in the World Health Organization Eastern Mediterranean region. The region comprises 22 predominantly Islamic countries, and health indicators vary widely.

### Review methods

We used data from WHO and Unicef to determine child deaths and cause specific mortality.<sup>7-8</sup> We obtained information from the official websites of ministries of health and nationally accredited institutions or research organisations, describing national research priorities for child health, funding for research, and policy. In addition, we reviewed research funding data from the WHO (Eastern Mediterranean

Department of Paediatrics and Child Health, Aga Khan University, Karachi, Pakistan  
Zulfiqar A Bhutta  
Husein Lalji Dewraj  
professor

continued over

BMJ 2006;333:839-42




Further details of the review are on [bmj.com](http://bmj.com)

Department of Paediatrics, King Faisal Specialist Hospital and Research Center, Riyadh, Saudi Arabia  
 Asim Belgaumi  
*head, section of leukaemia and lymphoma*  
 Research Policy and Cooperation Unit, WHO Eastern Mediterranean Region, Cairo, Egypt  
 Mohammad Abdur Rab  
*regional adviser*

Department of Paediatrics, School of Medicine, Khartoum, Sudan  
 Zein Karrar  
*professor*  
 Mansourah  
 Children's Hospital, Mansourah, Egypt  
 Mohamed Khashaba  
*head of department*

l'Unité de Nutrition, Service de Pédiatrie III, Hôpital d'enfants, Rabat, Morocco  
 Nezha Mouane  
*professor*

Correspondence to:  
 Z A Bhutta  
 zulfiqar.bhutta@aku.edu



Trends in child mortality for countries with highest burden

Region) research cooperation and policy unit on subjects directly or indirectly related to newborn and child health as well as information available on activities within the region from other United Nations agencies (Unicef, UN Population Fund, UN Development Programme), World Bank, and major bilateral donors. We also evaluated the evidence base for established interventions as well as relatively new strategies for addressing newborn and child health in health system settings. Finally, we assessed the relation between government effectiveness and child health. Full details of the methods are on [bmj.com](http://bmj.com).

### Child mortality in the region

The Eastern Mediterranean region accounts for almost 15% of the total global burden of newborn and child mortality, most of which is concentrated in a few countries. Of the current 1.4 million deaths among children under 5 in the region every year, over 1.2 million (91%) occur in just seven countries (Pakistan, Afghanistan, Egypt, Sudan, Somalia, Iraq, and Yemen) with mortality exceeding 50/1000 live births. Despite falls in mortality from 1970 to 1990, rates in these countries have recently increased (Afghanistan, Iraq, and Somalia) or stagnated (figure). Accurate data for child health indicators in the Gaza Strip and Darfur are not available, and we do not yet know the full effect of the current humanitarian crisis in Lebanon on child health.

### How many lives can we save and how?

Table 1 indicates the distribution of child deaths from specific causes and the proportions that are preventable in the seven high burden countries of the region. Table 2 details various intervention strategies and their effect on child mortality. We assumed a target of 90% coverage except for the expanded programme of immunisation and tetanus toxoid, for which we assumed 95% coverage. Although these targets are lower than the assumed 99% coverage in the child survival series,<sup>2</sup> our analysis indicates that these interventions can still potentially save 52% of all newborn and child deaths in these countries. If we could add newer interventions, including the vaccines for rotavirus diarrhoea and invasive pneumococcal

disease, as well as other strategies for secondary care of preterm infants,<sup>9</sup> an additional 10-15 % lives can be saved.

So why are these interventions not happening? The explanations relate to issues such as lack of awareness at the policy level of the importance of child survival, the specific packages of interventions selected by programmes, and efforts to scale up delivery strategies in health systems. We found a close correlation between our index of government effectiveness—that is, ability to implement policy—and child mortality (see fig B on [bmj.com](http://bmj.com)). Although specific data on national spending for newborn and child health in the high burden countries were not available, overall funding levels for health services varied considerably. Figure B on [bmj.com](http://bmj.com) indicates the relation of the health to defence expenditure ratio and infant mortality in Eastern Mediterranean countries with the highest numbers of child deaths

Scaling up delivery is recognised as essential to meeting the millennium development goals<sup>10</sup> and must be supported with financial and other resources. Current coverage for many of these interventions is extremely variable, with big contrasts between rural and urban populations for some basic newborn and child survival interventions such as skilled care at birth, measles vaccination, and nutrition support programmes tackling chronic under-nutrition problems such as stunting (see fig D on [bmj.com](http://bmj.com)).

The state of maternal care and strategies for newborn care in these countries clearly contribute to the persistently high rates of newborn deaths. A large proportion of all births take place at home with unskilled attendants, and few community based strategies or outreach programmes for maternal and newborn care are in place. Despite the success of composite community based initiatives,<sup>11</sup> only Pakistan and Iran have been able to set up a robust programme of interventions through community based health workers.<sup>12 13</sup> Although much of the focus has been on community delivery of specific interventions and commodities, the effect of such programmes can be

**Table 1** Total and preventable deaths in children aged under 5 years in Eastern Mediterranean countries with mortality above 50/1000 live births

Disease or condition	No of deaths in 2004 (000s)	No (%) preventable (000s)
<b>Non-neonatal:</b>		
Diarrhoea	238	180 (75.6)
Pneumonia	291	178 (61.2)
Measles	56	42 (74.6)
Malaria	37	32 (84.3)
HIV and AIDS	4	3 (68.1)
Other	186	0
<b>Neonatal:</b>		
Asphyxia	117	58 (49.1)
Prematurity	127	63 (49.6)
Sepsis/pneumonia	167	100 (59.9)
Tetanus	69	44 (64.1)
Diarrhoea	21	3 (15.9)
Congenital disorders	52	2 (4.2)
Other	33	1 (2.1)
<b>Total</b>	<b>1401</b>	<b>706 (50.4)</b>

Numbers do not always calculate exactly because of rounding.

**Table 2** Estimate of deaths averted (in thousands) among children aged under 5 years by interventions in Eastern Mediterranean countries with mortality above 50/1000 live births

Intervention	No (%) of deaths prevented
<b>Preventive</b>	
Family and community care:	
Periconceptual folic acid	4 (0.3)
Promotion of breast feeding	151 (10.8)
Appropriate complementary feeding strategies including dietary diversification	54 (3.9)
Water, sanitation, and hygiene strategies	31 (2.2)
Outreach services:	
Effective antenatal care	17 (1.2)
Tetanus toxoid administration	36 (2.6)
Intermittent preventive treatment of malaria in pregnancy	2 (0.2)
Insecticide treated materials for malaria	24 (1.7)
Vitamin A supplementation	11 (0.8)
Clinic based care:	
Treatment of asymptomatic bacteriuria in pregnancy	8 (0.6)
Skilled maternal and immediate newborn care	53 (3.8)
Antenatal steroids in high risk cases (preterm births)	23 (1.6)
Extra care of low birthweight infants	19 (1.4)
Haemophilus influenzae B vaccine (as part of standard vaccination schedule)	29 (2.1)
Measles vaccine	30 (2.1)
<b>Treatment</b>	
Family and community care:	
Oral rehydration therapy	65 (4.7)
Appropriate antibiotics for pneumonia	42 (3.0)
Outreach services:	
Zinc for diarrhoea	4 (0.3)
Appropriate antimalarials including artemisin combination	4 (0.3)
Clinic based care:	
Antibiotics for preterm premature rupture of membranes	5 (0.3)
Emergency obstetric care	19 (1.3)
Emergency newborn care	35 (2.5)
Appropriate treatment of neonatal pneumonia	31 (2.2)
Antibiotics for dysentery	6 (0.4)
Vitamin A for measles	1 (0.1)
Antiretroviral treatment	3 (0.2)

greatly increased by strategies aimed at empowerment of women, domiciliary practices, community engagement, and governance of services.

Over the past few years, the evidence on the feasibility and potential impact of community based interventions and innovative delivery strategies has greatly improved.<sup>13-15</sup> Many of the interventions to prevent child deaths that our analysis identified could be delivered to whole populations through community based approaches and outreach programmes. However, efforts for implementing and scaling up these interventions are slow in most countries. More importantly, where interventions exist, monitoring and evaluation frameworks are poor and effectiveness evaluation and research extremely limited.

### Importance of research into effectiveness

Evaluation of the effectiveness of child survival interventions is critical to generating the momentum to introduce and increase their coverage within health

### Summary points

Child deaths in the Eastern Mediterranean region account for 15% of the total global burden for newborn and child mortality

Most of the deaths occur in just seven countries, several of which are in a state of conflict

Available evidence based interventions, if fully implemented, can prevent half of all child deaths in these countries

Barriers to implementation of these interventions are not so much lack of knowledge as political will and appropriate allocation of resources

systems.<sup>16</sup> Evaluation does not always require randomised controlled trials. Alternative study methods, such as well designed observational studies or phased introduction of interventions through a step-wedge design can also provide strong evidence to inform policy making for child health.<sup>17</sup> Little research has been done on the effectiveness of essential child health interventions and delivery packages,<sup>18</sup> particularly community based interventions and outreach programmes. Such evidence and experience in real health systems is critical in developing policy, particularly for newer interventions such as use of zinc for diarrhoea. This was highlighted by the recent evaluation of supervised and unsupervised treatment of children with malaria in Myanmar, which showed that an inexpensive dihydroartemisinin-piperazine combination was as effective as artesunate-mefloquine.<sup>19</sup>

We also evaluated the evidence base and gaps for many of the interventions when packaged together according to current delivery strategies. As table B on bmj.com shows, much of the evidence has been derived from relatively small scale efficacy studies, and few interventions have been evaluated in whole health systems. Very few research projects are evaluating strategies for scaling up child survival interventions to whole health systems (total expenditure under \$4m (£2m, €3m)). We could identify only three projects during 2000-4 in the region that had evaluated the effectiveness of country-wide child survival interventions or packages.

### Role of social determinants and political will

Although we have largely focused on health interventions, this should not be misconstrued as downplaying the importance of factors such as education, social development, economic and sex equality, and feudal power structures.<sup>20</sup> Interventions related to some of these determinants would normally be implemented by sectors other than health but are critical to supporting public health interventions.

In addition, several elements in health sector planning such as devolution and democratisation of decision making, governance, and reform are

critical to progress in the region. Our analysis shows that relatively low levels of spending on health (in comparison with military expenditure) and ineffective health sector governance are important correlates of child survival in the poorest countries of the region. This suggests that much can be done, even with limited resources. Indeed, Morocco and Egypt have made remarkable gains in child survival over the past decade with relatively limited resources (box).

## Conclusions

Our review of child survival has several limitations. Our estimates of the effect of interventions are largely derived from small efficacy studies and have wide confidence intervals. However, we do not think that this detracts from the main message that much can be done with existing knowledge and resources. Because of the focus on survival, we have not discussed the emerging issues of mental health and child development, urbanisation, environmental change, and lifestyle issues such as obesity. The high rates of consanguinity and inherited disorders in the region will also emerge as public health issues once child mortality drops. We believe, however, that our data point to an unacceptable persistent burden of child mortality from common disorders in some countries in the region. In most instances, these deaths are largely preventable by well established interventions.

### Morocco's success story

Morocco has a population of 30 million and mortality for children under 5 years is 43/1000 live births. Mortality has fallen sharply since Morocco became independent, averaging 6% a year. Despite numerous challenges and a per capita annual income of \$1540, the government has been able to introduce several high quality health programmes:

**Immunisation**—A national programme was started in 1981 and reinforced in 1987 comprising systematic vaccinations in health posts, annual national vaccination days, and mobile teams going four times a year to rural areas. Current vaccine coverage exceeds 95%

**Diarrhoeal diseases**—A control programme was initiated in 1979 and extended in 1990

**Acute respiratory infections**—Programme started in 1997

**Nutrition programme** includes promotion of exclusive breast feeding and complementary feeding; monitoring of child growth; prevention of vitamin D, vitamin A, and iron deficiencies; and detection and treatment of severe malnutrition. Vitamin A is also distributed at time of vaccination.

Since 1998 many of these vertical programmes have been brought together in the integrated management of sick children strategy. The decline in mortality has been faster for children aged 1-4 years than for infants. A recent evaluation concluded that vaccine preventable diseases contributed to 23% of the total fall in mortality, diarrhoea and malnutrition to 33%, and other infectious diseases to 29%. However, mortality for accidents, acute respiratory infection in infancy, and some neonatal conditions (birth trauma and prematurity) has not changed much. These findings have led to the recent inclusion of the neonatal period in the national strategy for management of sick children.

These existing interventions, as well as the promising new targeted strategies, must be delivered to all those who need them most. This will require concerted efforts by public health policy makers, development agencies, and civic societies to garner resources for child health. Not only must these interventions be based on robust evidence but their implementation in health systems must also be part of a learning process.

We thank Arjumand Rizvi for help with the analysis of the effect of interventions and Mohammad Afzal for additional information on research funding.

Competing interests: None declared.

Contributors and sources: This paper is part of a larger analysis of child health and survival in the Eastern Mediterranean region and the global evidence base of interventions. ZAB conceived of the idea for the article and analysis, collected the primary information, and wrote the first draft. All co-authors contributed to the analytical and paper writing process as well as contributing specific information as required. ZAB is guarantor.

- Black RE, Morris SS, Bryce J. Where and why are 10 million children dying every year? *Lancet* 2003;361:2226-34.
- Jones G, Steketee RW, Black RE, Bhutta ZA, Morris SS, Bellagio Child Survival Study Group. How many child deaths can we prevent this year? *Lancet* 2003;362:65-71.
- Lawn JE, Cousens S, Zupan J. Lancet Neonatal Survival Steering Team. 4 million neonatal deaths: when? where? why? *Lancet* 2005;365:891-900.
- Darmstadt GL, Bhutta ZA, Cousens S, Adam T, Walker N, de Bernis L, Lancet Neonatal Survival Steering Team. Evidence-based, cost-effective interventions: how many newborn babies can we save? *Lancet* 2005;365:977-88.
- World Health Organization. *World health report 2005. Make every mother and child count*. www.who.int/whr/2005/en/index.html (accessed 26 Jul 2006).
- Task Force on Child Health and Maternal Health. *Who's got the power? Transforming health systems for women and children*. New York: UN Millennium Project, 2005. www.millenniumproject.org/reports/health.htm (accessed 26 Jul 2006).
- World Health Organization. *Statistics*. www.who.int/research/en/ (accessed 27 Jul 2006).
- Unicef. *Monitoring the status of women and children*. www.childinfo.org/ (accessed 26 Jul 2006).
- Bhutta ZA, Khan I, Salat S, Raza F, Ara H. Reducing length of stay in hospital for very low birth weight infants by involving mothers in a stepdown unit: an experience from Karachi (Pakistan). *BMJ* 2004;329:1151-5.
- Travis P, Bennett S, Haines A, Pang T, Bhutta Z, Hyder AA, et al. Overcoming health-systems constraints to achieve the millennium development goals. *Lancet* 2004;364:900-6.
- WHO Regional Office for the Eastern Mediterranean. *Community based initiatives: Success stories from the Eastern Mediterranean Region*. 2006. www.emro.who.int/cbi/pdf/CBI\_SuccessStories.pdf (accessed 20 Sep 2006).
- Shadpour K. Primary health care networks in the Islamic Republic of Iran. *East Mediterranean Health J* 2000;6:822-5.
- Haines A, Sanders D, Rowe AK, Lawn J, Jan S, Walker D, et al. Achieving child survival goals: potential contribution of community health workers. *Lancet* (in press).
- Bhutta ZA, Darmstadt GL, Hasan BS, Haws RA. Community-based interventions for improving perinatal and neonatal health outcomes in developing countries: a review of the evidence. *Pediatrics* 2005;115(suppl):519-617.
- Bang AT, Baitule SB, Reddy HM, Deshmukh MD, Bang RA. Low birth weight and preterm neonates: can they be managed at home by mother and a trained village health worker? *J Perinatol* 2005;25(suppl 1):S72-81.
- Bryce J, Victora CG, MCE-IMCI Technical Advisors. Ten methodological lessons from the multi-country evaluation of integrated management of childhood illness. *Health Policy Plan* 2005;20(suppl 1):i94-105.
- Victora CG, Habicht JP, Bryce J. Evidence-based public health: moving beyond randomized trials. *Am J Public Health* 2004;94:400-5.
- Bhutta ZA, Lassi S. Child health: how can health research make a difference? *Global Forum Update on Research for Health* 2005;107-11. www.globalforumhealth.org/filesupld/global\_update1/GlobalUpdate4.pdf (accessed 26 Jul 2006).
- Smithuis F, Kyaw MK, Phe O, Aye KZ, Htet L, Barends M, et al. Efficacy and effectiveness of dihydroartemisinin-piperacillin versus artesunate-mefloquine in falciparum malaria: an open-label randomized comparison. *Lancet* 2006;367:2075-85.
- World Health Organization. *Social determinants of health and health equity in the EMR/MENA region*, 2006. www.who.int/social\_determinants/resources/emro\_workshop\_rpt.pdf (accessed 15 Sep 2006).

doi 10.1136/bmj.38979.379641.68