



9-2019

Cross-sectional study of knowledge of stroke Management amongst doctors of a tertiary Care Hospital; Peshawar, Pakistan

Muhammad Fozan Khan
Rehman Medical Institute Peshawar Pakistan

Farzana Behtani
Rehman Medical Institute, Peshawar, Pakistan

Omer Nasim
Rehman Medical Institute, Peshawar, Pakistan

Muhammad Shahid Iqbal
Rehman Medical Institute, Peshawar, Pakistan

Sohail Khan
Rehman Medical Institute, Peshawar, Pakistan

See next page for additional authors

Follow this and additional works at: <https://ecommons.aku.edu/pjns>



Part of the [Neurology Commons](#)

Recommended Citation

Fozan Khan, Muhammad; Behtani, Farzana; Nasim, Omer; Shahid Iqbal, Muhammad; Khan, Sohail; and Khizar Hayat, Muhammad (2019) "Cross-sectional study of knowledge of stroke Management amongst doctors of a tertiary Care Hospital; Peshawar, Pakistan," *Pakistan Journal of Neurological Sciences (PJNS)*: Vol. 14 : Iss. 3 , Article 9.

Available at: <https://ecommons.aku.edu/pjns/vol14/iss3/9>

Cross-sectional study of knowledge of stroke Management amongst doctors of a tertiary Care Hospital; Peshawar, Pakistan

Authors

Muhammad Fozan Khan, Farzana Behtani, Omer Nasim, Muhammad Shahid Iqbal, Sohail Khan, and Muhammad Khizar Hayat

CROSS-SECTIONAL STUDY OF KNOWLEDGE OF STROKE MANAGEMENT AMONGST DOCTORS OF A TERTIARY CARE HOSPITAL; PESHAWAR, PAKISTAN

Muhammad Fozan Khan¹, Farzana Behttani¹, Omer Nasim², Muhammad Shahid Iqbal¹, Sohail Khan¹, Muhammad Khizar Hayat³

1: Consultant Stroke Unit, Rehman Medical Institute Peshawar Pakistan, 2: Lecturer in Anatomy, Rehman Medical College Peshawar, 3: House Officer General Medicine, Rehman Medical Institute Peshawar

Correspondence to: Muhammad Fozan Khan, Incharge Stroke Unit Rehman Medical Institute, Peshawar, Pakistan Email: fozan5036@hotmail.com

Date of submission: February 19, 2019 **Date of revision:** May 22, 2019 **Date of acceptance:** June 15, 2019

ABSTRACT:

Background: Stroke is one of the leading causes of mortality and neurological disability worldwide. 'Time is brain' and prompt management of stroke is vital to improve the outcomes. Knowledge of stroke management is important for clinicians working in various specialties and this study focused on evaluation of this vital aspect.

Objective: To document the knowledge of stroke management amongst doctors, working at different stages of career in different specialties at a tertiary care hospital.

Materials & Methods: A questionnaire was developed using current stroke guidelines of the Royal College of Physicians London and American Heart and Stroke Association. The questionnaire was distributed among 73 doctors working in various specialties and grades. Data analysis was done using SPSS version 21.0.

Results: House officers (n=33), Medical officers (n= 7), Post graduate trainees (n=20), Registrars (n=3), Consultants (n=9) participated. Diffusion weighted (DWI) MRI was recognised as the best diagnostic tool for acute ischaemic stroke (58.33%). BP management knowledge in haemorrhagic stroke was poor (60.56%). Less than half (40.85%) chose tissue plasminogen activator (t-PA) as the preferred initial treatment. Majority were unaware of percutaneous endoscopic gastrostomy (PEG) for unsafe swallowing (45.07%). Few recognised (19.72%) that the upper limit in blood pressure management for acute ischaemic stroke was 220/120 without other end organ damage. Only (55.56%) thought that IPCs (intermittent pneumatic compressions) were the safest method for DVT prophylaxis.

Conclusion: The survey shows that the knowledge of doctors was inadequate. Improving medical training of stroke management and awareness of stroke as an emergency ('Brain attack') needs to be improved for optimal outcomes.

Keywords: stroke; cerebrovascular accident; knowledge; thrombolysis; Pakistan.

INTRODUCTION: Stroke is amongst the most common causes of death and the first leading cause of disability in developed and developing countries.¹ To determine the true incidence of stroke, there are no large-scale epidemiological studies in Pakistan. In previous retrospective studies of patients of stroke in other medical units in two major hospitals over an 8-year period, incidence was 796/12,454 (6.4%).² Stroke is a medical emergency and the estimated annual incidence in Pakistan is 250/100,000, translating to 350,000 new cases every year.³ Around 40% of people aged 18 years or above have hypertension, dyslipidemia or a history of active

smoking.⁴ Burden of major vascular risk factors i.e. hypertension, diabetes mellitus, smoking, dyslipidemia and obesity is enormous in Pakistan. The study participants were healthy individuals who came for routine checkups and only 40% of the hypertensive patients had optimal blood pressure levels.⁴ There was significant family history of ischaemic heart disease in 42%, obesity in 24%, hypertension in 19%, and diabetes mellitus in 15%.⁴ Contrary to decline in the incidence of the disease in the Western population, the burden of stroke in South Asian countries (India, Pakistan, Bangladesh, and Sri Lanka) has been rising.⁵

These are challenging times in the health care system in Pakistan. Limited resources are allocated for prevention of vascular diseases. There are compelling epidemiological and scientific reasons that the incidence of such diseases will increase in the coming decades. Considering the high mortality and disability associated with stroke, good knowledge of identifying stroke risk factors, symptoms and management are important for doctors working in most specialties. This study documents the knowledge of stroke management amongst doctors, working at different stages of career in different specialties at a tertiary care hospital.

Material and Methods:

A cross-sectional survey design was used. Study was conducted at Rehman Medical Institute (RMI) Peshawar from June-July 2019. The data collection tool had three distinct parts. The first being an introduction to study objective, informed consent and assurance of anonymity of the data. The second section had demographic details, including the specialty and working grade of the doctor. The last part consisted of 15 multiple choice questions (MCQs) which were prepared in the single best choice format. A questionnaire was developed using current stroke guidelines (2016) of the Royal College of Physicians London and American Heart and Stroke Association.^{6, 7}

The pre-tested self-administered questionnaire was distributed among 73 doctors. Data was analysed using SPSS Version 21.0.

Results:

Respondents included House officers (n=33), Medical officers (n= 7), Post graduate trainees (n=20), Registrars (n=3), Consultants (n=9). Most correctly identified CT brain as the initial diagnostic modality (95.83%), DWI MRI as the best diagnostic tool for acute ischemic stroke (58.33%). Majority were unaware of BP management in acute haemorrhagic stroke, (60.56%). Most were familiar with the risk of deep vein thrombosis in immobilised stroke patients (94.44%). On the other hand, less than half (40.85%) chose tissue plasminogen activator (t-PA) as the preferred initial choice of intervention in eligible patients within 4.5 hours of presentation with acute ischaemic stroke. Majority were unaware of percutaneous endoscopic gastrostomy (PEG) for unsafe swallowing greater than 4 weeks, correct answer was given by less than half of participants (45.07%). Very few recognised (19.72%) that the upper limit in blood pressure management for ischemic stroke was 220/120 within the first 48 to 72 hours without other end organ damage. Also, only (55.56%) thought that IPCs (intermittent pneumatic compressions) were the safest method for DVT prophylaxis after stroke. Knowledge that lowering and maintaining the blood pressure lowers the risk of stroke recurrence (80.28%) and that haemorrhagic stroke is an absolute contraindication to thrombolytic therapy (79.17%) was known to majority of the respondents. The percentage of missing responses ranged from 0.2 % - 1.0% which was not significant. The complete questionnaire with the highlighted correct answer is displayed in Table. 3.

The detailed results are shown in Table 1.0.

#	Item-wise Response Distribution on Stroke Knowledge Test	A	B	C	D
		n (%)	n (%)	n (%)	n (%)
1	The first emergency investigation of choice in a patient suspected of having an acute stroke is	95.8	4.2	-	-
2	Urgent Computed tomography (CT) of brain in patients with stroke helps in	1.4	95.8	2.8	-
3	The most accurate diagnostic test for detecting ischaemic stroke is	1.4	58.3	29.2	11.1
4*	A 55-year-old man is in Emergency Dept. with dysphasia and right sided weakness (power 1/5). He has been diagnosed as having acute left anterior circulation ischaemic stroke for last 2.5 hours. His CT brain is unremarkable and CT angiogram has not shown any large vessel occlusion. The emergency treatment should be	48.6	40.3	1.4	8.3
5*	After managing airway and breathing, which of the following should be the next step in management of a patient with acute haemorrhagic stroke	59.7	1.4	31.9	5.6
6	Evidence shows that patients with acute stroke have better outcomes if managed	1.4	2.8	94.4	1.4
7	Intravenous thrombolysis in ischaemic stroke can be performed	76.4	1.4	20.8	1.4
8*	The risk of recurrence of stroke can be significantly reduced by	-	13.9	5.6	79.2
9	Which of the following is an absolute contraindication to thrombolytic therapy	79.2	13.9	2.8	4.2
10	The development of which serious condition would you be concerned most in patients immobilised due to acute stroke	-	4.2	1.4	94.4
11	The most important contraindication to anticoagulation is	2.8	80.6	2.8	13.9
12*	If swallowing is unsafe after 4 weeks of stroke, the next recommended step of management to consider is	38.9	1.4	13.9	44.4
13	The safest method of Deep Vein Thrombosis (DVT) prophylaxis after acute stroke is	13.9	11.1	19.4	55.6
14*	Target systolic Blood Pressure in case of acute haemorrhagic stroke for initial one week is	-	34.7	8.3	55.6
15	Within 48 to 72 hours after acute ischaemic stroke (without other end organ damage), acceptable BP not requiring urgent lowering of BP is up to	19.4	13.9	47.2	18.1

Table. 1.0 Question-wise response distribution on stroke knowledge test
Correct answer bold | *1-3 participants did not respond to these questions.

QUESTIONS		PARTICIPANT STATUS									
		HOUSE OFFICERS		MEDICAL OFFICERS		TRAINEE MEDICAL OFFICERS		REGISTRARS		CONSULTANTS	
		Frequency (n)	Percentage (%)	Frequency (n)	Percentage (%)	Frequency (n)	Percentage (%)	Frequency (n)	Percentage (%)	Frequency (n)	Percentage (%)
Question 1	Incorrect	1	3	0	0	2	10	0	0	0	0
	Correct	32	97	8	100	18	90	2	100	9	100
Question 2	Incorrect	2	6	0	0	1	5	0	0	0	0
	Correct	31	94	8	100	17	95	2	100	9	100
Question 3	Incorrect	22	67	3	38	4	20	1	50	0	0
	Correct	11	33	5	63	16	80	1	50	9	100
Question 4	Incorrect	21	64	6	75	10	50	2	100	3	33
	Correct	12	36	2	25	10	50	0	0	5	56
Question 5	Incorrect	16	48	2	25	7	35	2	100	2	22
	Correct	17	52	6	75	13	65	0	0	7	78
Question 6	Incorrect	3	9	0	0	1	5	0	0	0	0
	Correct	30	91	8	100	19	95	2	100	9	100
Question 7	Incorrect	12	36	3	38	2	10	0	0	0	0
	Correct	21	64	5	63	18	90	2	100	9	100
Question 8	Incorrect	7	21	3	38	4	20	0	0	1	11
	Correct	26	79	5	63	16	80	2	100	8	89
Question 9	Incorrect	9	27	1	13	4	20	0	0	0	0
	Correct	24	73	7	88	16	80	2	100	9	100
Question 10	Incorrect	3	9	0	0	0	0	0	0	1	11
	Correct	30	91	8	100	20	100	2	100	8	89
Question 11	Incorrect	6	18	1	13	5	25	0	0	1	11
	Correct	26	79	7	88	15	75	2	100	8	89
Question 12	Incorrect	23	70	2	25	14	70	0	0	1	11
	Correct	10	30	6	75	6	30	2	100	8	89
Question 13	Incorrect	15	45	1	13	13	65	2	100	1	11
	Correct	18	55	7	88	7	35	0	0	8	89
Question 14	Incorrect	19	58	3	38	6	30	0	0	4	44
	Correct	14	42	5	63	14	70	2	100	5	56
Question 15	Incorrect	27	82	5	63	16	80	2	100	8	89
	Correct	6	18	3	38	4	20	0	0	1	11

Table 2: Question-wise response distribution on stroke knowledge test based on participants' career status (missing responses taken as incorrect, 1-2%)

Table 3: Complete Questionnaire with highlighted answers.

Question	Stem A	Stem B	Stem C	Stem D
1. The first emergency investigation of choice in a patient suspected of having an acute stroke is?	Computed tomography (CT) scan	Magnetic Resonance Imaging (MRI)	X-ray head	Lumbar puncture
2. Urgent Computed tomography (CT) of brain in patients with stroke helps in?	Identification of post stroke depression	Distinguishing between haemorrhagic and ischaemic stroke	Localizing the aneurysm	Preventing further haemorrhage
3. The most accurate diagnostic test for detecting ischaemic stroke is?	Non-Diffusion Weighted Magnetic Resonance Imaging (MRI)	Diffusion Weighted (DWI) Magnetic Resonance Imaging (MRI)	Computed Tomography (CT) with contrast	CT Intracranial Angiogram
4. A 55-year-old man is in Emergency Dept. with dysphasia and right sided weakness (power 1/5). He has been diagnosed as having acute left anterior circulation ischaemic stroke for last 2.5 hours. His CT brain is unremarkable and CT angiogram has not shown any large vessel occlusion. The emergency treatment should be?	Aspirin	Tissue Plasminogen Activator (tPA)/Alteplase	Warfarin	Clopidogrel

5. After managing airway and breathing, which of the following should be the next step in management of a patient with acute haemorrhagic stroke?	Lower blood pressure to optimal targets	Give statins	Administer intravenous fluids	Administer subcutaneous Heparin
6. Evidence shows that patients with acute stroke have better outcomes if managed?	In General Medicine ward	Coronary Care Unit	Dedicated Stroke Unit	Vascular surgery ward
7. Intravenous thrombolysis in ischemic stroke can be performed?	Within 4.5 hours after onset of symptoms	After 4.5 hours of onset of symptoms	Within 12 hours of onset of symptoms	After 12 hours
8. The risk of recurrence of stroke can be significantly reduced by?	Good sleep	Regular exercise	Subcutaneous Heparin	Optimal control of blood pressure
9. Which of the following is an absolute contraindication to thrombolytic therapy?	Intracerebral haemorrhage	Pregnancy	Age >70 years	Major surgery within the last 1 year
10. The development of which serious condition would you be concerned most in patients immobilised due to acute stroke?	Depression	Hemiplegic Shoulder pain	Urinary tract infection	Deep vein thrombosis
11. The most important contraindication to anticoagulation is?	Epilepsy	Acute subarachnoid haemorrhage	Deep vein thrombosis	Pregnancy
12. If swallowing is unsafe after 4 weeks of stroke, the next recommended step of management to consider is?	Continue Nasogastric feeding	Oral sips while accepting risk of aspiration	Total Parenteral Nutrition	Percutaneous Endoscopic Gastrostomy
13. The safest method of Deep Vein Thrombosis (DVT) prophylaxis after acute stroke is?	Enoxaparin	Warfarin	Rivaroxaban	Intermittent Pneumatic Compression (IPC)
14. Target systolic Blood Pressure in case of acute hemorrhagic stroke for initial one week is?	200mmHg	<130 mmHg	120 mmHg	140-150 mmHg
15. Within 48 to 72 hours after acute ischaemic stroke (without other end organ damage), acceptable BP not requiring urgent lowering of BP is up to?	220/120 mmHg	190/100 mmHg	140/90 mmHg	120/80 mmHg

Discussion:

In our study only a few recognised (19.72%) that the upper limit in blood pressure management for ischaemic stroke was 220/120 within the first 48 to 72 hours, in that study less than half adequately treated hypertension.⁸ The practical implications can lead to inadequate or even detrimental outcomes.

In Pakistan, 50% of transient ischemic attacks (TIAs) and more than 35% strokes are most often misdiagnosed.⁹ Multiple surveys around the world have revealed that the generic knowledge of stroke in the general population is generally inadequate in both the developed¹⁰⁻¹² and developing countries.¹³⁻¹⁶ In a study carried out in the Gulf countries, many patients were not aware of the term 'stroke'.¹³ Also, in Pakistan respondents from high schools in a single district showed that they were unaware of the risk factors for stroke and the symptoms and did not know what to do in a case of acute stroke.¹⁶

Less than half (40.84%) of the respondent's chose tPA as the correct initial intervention in eligible candidates within time window. It also shows that awareness and knowledge of use of thrombolytic therapy is insufficient amongst the doctors. There is urgent need of promoting stroke care education amongst the doctors and public so that patients can present early to the tertiary care stroke centres within the 4.5 hours, 'golden timeframe' for

intravenous thrombolysis. Evidence clearly shows that strokes are preventable and treatable. And thousands of people are dying due to inadequate care and all these lives could be saved and could avoid living with permanent disability with service improvements.¹⁸

The public and private tertiary care hospitals should ensure establishment of fully resourced and dedicated stroke units. Majority of the respondents in our study had a consensus that a dedicated stroke unit is required for better outcome in patients and such units could curtail the rate of death and dependency rates by as great as 20% - which in terms of numbers mean saving thousands of lives and reducing lifelong disabilities.¹⁸

As an estimate, Pakistan's population might have 4.8% stroke annually. 17 It is equivalent to approximately 8.7 million stroke patients countrywide. Amongst these patients, 63% develop complications of stroke, and 7-20% unfortunately die. 18 Stroke is a huge burden on the healthcare system, families and the economy, hence the necessity of timely diagnosis and early intervention can prevent death and disability.

All medical professionals should be educated in a structured manner, e.g. lectures, interactive workshops and seminars. All hospitals must follow a standard, evidence-based stroke management protocol. Stroke management should be an important part of curriculum in the medical colleges. There is a need to perform a

nationwide survey on stroke that will reveal its accurate epidemiological factors. There is also an urgent need to establish dedicated stroke units at tertiary care hospitals.

Future studies need to be conducted, involving higher number of physicians to yield more detailed analysis of the knowledge and in turn it could help to highlight the short comings, to deduce the solutions and improvement of the quality of stroke care.

The survey shows that the knowledge of doctors was inadequate. Improving medical training of stroke management and awareness of stroke as an emergency ('Brain attack') needs to be improved for optimal outcomes. This study provides a basis for further research and analyses the trends in stroke knowledge amongst doctors working at a large tertiary care hospital. Training sessions, curriculum improvements and public awareness campaigns are likely to significantly improve stroke care in Pakistan.

References:

1. Feigin VL. Stroke epidemiology in the developing world. *Lancet* 2015;365:2160-61.
2. Vohra EA, Ahmed WU, Ali M. Aetiology and prognostic factors of patients admitted for stroke. *J Pak Med Assoc.* 2000;50:234-36.
3. Khealani BA, Hameed B, Mapari UU. Stroke in Pakistan. *J Pak Med Assoc.* 2008;58(7):400-3.
4. Iqbal SP, Dodani S, Qureshi R. Risk factors and behaviors for coronary artery disease (CAD) among ambulatory Pakistanis. *J Pak Med Assoc.* 2004;54:261- 66.
5. Bulatao RA, Stephens PW. Global estimates and projections of mortality by cause. Washington, DC: Population, Health and Nutrition Department; World Bank, preworking paper 1992;1007.
6. William J. Powers, Alejandro A. Rabinstein, Teri Ackerson, Opeolu M. Adeoye, Nicholas C. Bambakidis, Kyra Becker, et al. Guidelines for the Early Management of Patients With Acute Ischemic Stroke: A Guideline for Healthcare Professionals From the American Heart Association/American Stroke Association and on behalf of the American Heart Association Stroke Council. *Stroke.* 2018;49:e46–e99
7. College of Physicians National Clinical Guideline for Stroke (2019). *Strokeaudit.org*. Retrieved 8 July 2019, from [https://www.strokeaudit.org/SupportFiles/Documents/Gudelines/2016-National-Clinical-Guideline-for-Stroke-5t-\(1\).aspx](https://www.strokeaudit.org/SupportFiles/Documents/Gudelines/2016-National-Clinical-Guideline-for-Stroke-5t-(1).aspx)
8. Wasay M, Khealani B, Yousuf A, Azam I, Rathi SL, Malik A, Haq A: Knowledge gaps in strokecare: results of a survey of family physicians in Pakistan. *J Stroke Cerebrovasc Dis.* 2011;20:282–86.
9. Khubaib MU, Rathore FA, Waqas A, Jan MM, Sohail S. Knowledge regarding basic facts of Stroke among Final Year MBBS Students and House Officers: A cross-sectional survey of 708 respondents from Pakistan. *Cureus.*2016;8(3):e539.
10. Spark JI, Blest N, Sandison S, Puckridge PJ, Saleem HA, Russell DA: Stroke and transient ischaemic attack awareness. *Med J Aust.* 2011;195:16–19.
11. Nedeltchev K, Fischer U, Arnold M, Kappeler L, Mattle HP: Low awareness of transient ischemic attacks and risk factors of stroke in a Swiss urban community. *J Neurol.* 2007;254:179–84.
12. Johnston SC, Fayad PB, Gorelick PB, Hanley DF, Shwayder P, van Husen D, Weiskopf T: Prevalence and knowledge of transient ischemic attack among US adults. *Neurology.* 2003, 60:1429-34.

13. Kamran S, Bener AB, Deleu D, Khoja W, Jumma M, Al Shubali A, et. al. The level of awareness of stroke risk factors and symptoms in the Gulf Cooperation Council Countries: Gulf Cooperation of Stroke awareness study. *Neuroepidemiology*. 2007;29:235-42.
14. Góngora-Rivera F, Gutiérrez-Jiménez E, Zenteno MA; GEPEVC Investigators: Knowledge of ischemic stroke among a Mexico City population. *J Stroke Cerebrovasc Dis*. 2009, 18:208-13.
15. Falavigna A1, Teles AR, Vedana VM, Kleber FD, Mosena G, Velho MC, Mazzocchin T, Silva RC, Lucena LF, Santin JT, Roth F: Awareness of stroke risk factors and warning signs in southern Brazil. *Arq Neuropsiquiatr*. 2009, 67:1076-81.
16. Farooq MU, Bhatt A, Safdar A, Kassab MY, Majid A: Stroke symptoms and risk factor awareness in high school children in Pakistan. *Int J Stroke*. 2012;7:e15.
17. Jafar TH: Blood pressure, diabetes, and increased dietary salt associated with stroke--results from a community-based study in Pakistan. *J Hum Hypertens*. 2006;20:83-85.
18. Malik A, Khatri I, Wasay M: Stroke manifesto- What must be done for stroke care in Pakistan. *Pak J Neurol Sci*. 2015;10:1-3.

Conflict of interest: Author declares no conflict of interest.

Funding disclosure: Nil

Author's contribution:

MFK designed and led the project

FB, ON, MFK did data collection/correction, manuscript writing & editing of the manuscript

ON, MSI, SK did data collection, statistical analysis & manuscript writing

ON, FB did data correction, computer generation of the data bank

MKH, MFK, MSI, SK did the editing, review & final approval of the manuscript