



THE AGA KHAN UNIVERSITY

eCommons@AKU

Department of Medicine

Department of Medicine

January 2010

Future prospects for the utility of interferon assays in tuberculosis diagnosis

Taimur Saleem

Madiha Syed

Bushra Jamil

Aga Khan University, bushra.jamil@aku.edu

Follow this and additional works at: https://ecommons.aku.edu/pakistan_fhs_mc_med_med



Part of the [Internal Medicine Commons](#)

Recommended Citation

Saleem, T., Syed, M., Jamil, B. (2010). Future prospects for the utility of interferon assays in tuberculosis diagnosis. *Journal of Pakistan Medical Association*, 704.

Available at: https://ecommons.aku.edu/pakistan_fhs_mc_med_med/467

Students' Corner

Letter to the Editor

Future prospects for the utility of interferon assays in tuberculosis diagnosis

Madam, tuberculosis remains an important global health problem. For Pakistan, the control of the disease should be a top priority as Pakistan is the 7th leading country with tuberculous disease.¹ Tuberculosis remains a diagnostic dilemma for physicians in endemic areas. Its ability to involve virtually any organ of the body along with its high prevalence in the region warrants its assignment to the list of differential diagnoses for a multitude of presentations. The pursuit of a definitive diagnosis for tuberculosis is imperative because of the consideration of side effects of anti-tuberculous therapy in the recipient. On the other hand, it is important to treat this condition because of its potential for curability; failure to treat can have devastating effects on individuals with this disease. Moreover, it is important to correctly recognize individuals with the disease from amongst the pool of

the most important parameters. Many studies have reported that the sensitivity and specificity of interferon assays is higher as compared to tuberculin skin test as far as latent tuberculosis is concerned.²⁻⁴ Limited data has also shown an association between interferon gamma response and progression of latent tuberculosis to active tuberculosis.³ Mycobacterial antigen six kDa early secreted antigenic target (ESAT6) and culture filtrate protein 10 (CFP10) can be used to detect *M. tuberculosis* associated interferon gamma responses. In a study conducted in Pakistan recently, it was seen that ESAT6-induced interferon gamma and CXCL9 responses discriminated the severity of tuberculous infections.⁵

However, the cost of this assay and the juxtaposed need for appropriate laboratory facilities and personnel are factors that ostensibly preclude its wider utility in developing countries.² Based on available literature, interferon gamma assays may not be conclusively recommended as a replacement for tuberculin skin testing because of lack of studies in endemic areas. Future studies

infected individuals; especially where prior BCG vaccination can confound the results of commonly employed diagnostic assays.

The traditional tuberculin skin test has remained a widely employed tool for diagnosing tuberculosis with various "risk-stratified" cut-offs being followed in various regions of the world for different sub-groups of the population.² However, the tuberculin skin test is fraught with shortcomings such as high dependency on the reader and the need for patients to return to get the test read after an appropriate time period;² both of these factors are important considerations in developing countries.

Interferon gamma assay is an alternative assay that is now rapidly emerging as a modern diagnostic tool for tuberculosis.² For any screening test, sensitivity is one of

done in endemic countries will, therefore, be instrumental in delineating the superiority of interferon gamma assays over the tuberculin skin test.

Taimur Saleem,¹ Madiha Syed,² Bushra Jamil³

Class of 2009, Medical College,^{1,2} Department of Medicine,³
Aga Khan University, Stadium Road, Karachi, Pakistan.

References

1. Ansari A, Talat N, Jamil B, Hasan Z, Razzaki T, Dawood G, et al. Cytokine gene polymorphisms across tuberculosis clinical spectrum in Pakistani patients. *PLoS One* 2009; 4: e4778
2. Pai M. Alternatives to the tuberculin skin test: interferon-gamma assays in the diagnosis of mycobacterium tuberculosis infection. *Indian J Med Microbiol* 2005; 23: 151-8.
3. Pai M, Riley LW, Colford JM Jr. Interferon-gamma assays in the immunodiagnosis of tuberculosis: a systematic review. *Lancet Infect Dis* 2004; 4: 761-76.
4. Luarti Ruiz F, Audran R, Harari A, Spertini F. [Specific interferon-gamma assays: a modern tool for tuberculosis diagnosis]. *Rev Med Suisse* 2006; 2: 1042, 1044-7.
5. Hasan Z, Jamil B, Ashraf M, Islam M, Yusuf MS, Khan JA, et al. ESAT6-induced IFN γ and CXCL9 can differentiate severity of tuberculosis. *PLoS One* 2009; 4: e5158.