



THE AGA KHAN UNIVERSITY

eCommons@AKU

Department of Paediatrics and Child Health

Division of Woman and Child Health

8-5-2020

The spectrum of primary immunodeficiencies at a tertiary care hospital in Pakistan

Sonia Qureshi

Aga Khan University, sonia.qureshi@aku.edu

Fatima Mir

Aga Khan University, fatima.mir@aku.edu

Samina Junejo

Indus Hospital, Korangi Road, Karachi, Pakistan

Khalid Saleem

Children's Hospital and The Institute of Child Health, Multan, Pakistan

Samreen Zaidi

National Institute of Blood Disease & Bone Marrow Transplantation, Pakistan

See next page for additional authors

Follow this and additional works at: [https://ecommons.aku.edu/](https://ecommons.aku.edu/pakistan_fhs_mc_women_childhealth_paediatr)

[pakistan_fhs_mc_women_childhealth_paediatr](https://ecommons.aku.edu/pakistan_fhs_mc_women_childhealth_paediatr)



Part of the [Allergy and Immunology Commons](#), [Maternal and Child Health Commons](#), and the [Pediatrics Commons](#)

Recommended Citation

Qureshi, S., Mir, F., Junejo, S., Saleem, K., Zaidi, S., Naveed, A. B., Ahmad, K., Qamar, F. N. (2020). The spectrum of primary immunodeficiencies at a tertiary care hospital in Pakistan. *World Allergy Organization Journal*, 13(7), 100133.

Available at: https://ecommons.aku.edu/pakistan_fhs_mc_women_childhealth_paediatr/946

Authors

Sonia Qureshi, Fatima Mir, Samina Junejo, Khalid Saleem, Samreen Zaidi, Abdullah B. Naveed, Khalil Ahmad, and Farah Naz Qamar



The spectrum of primary immunodeficiencies at a tertiary care hospital in Pakistan

Sonia Qureshi^a, Fatima Mir^a, Samina Junejo^b, Khalid Saleem^c, Samreen Zaidi^d, Abdullah B. Naveed^e, Khalil Ahmad^a and Farah Naz Qamar^{a*}

ABSTRACT

Background: Primary Immunodeficiency Disorders (PIDs) are well-known disorders in the West, but the recognition and diagnosis of these disorders is challenging in developing countries. We present the spectrum of PIDs seen at a tertiary care center in Pakistan, identified using clinical case definitions and molecular methods.

Methods: A retrospective chart review of children suspected to have PID was conducted at the Aga Khan University Hospital (AKUH) Karachi, Pakistan from 2010 to 2016. Data on demographics, clinical features, family history of consanguinity, sibling death, details of laboratory workup done for PID and molecular tests targeted panel next generation sequencing (NGS) or whole exome sequencing (WES) performed at the Geha laboratory at Boston Children's Hospital, USA was collected. The study was exempted from the Ethical Review Committee of AKUH.

Results: A total of 43 children visited the hospital with suspected PID during the study period. Genetic testing was performed in 31/43 (72.1%) children. A confirmed diagnosis of PID was established in 20/43 (46.5%) children. A pathogenic gene variant was identified in 17 (85%) of the 20 confirmed cases (Table 1). Twelve (60%) of the confirmed cases of PID were male. The most common presenting symptom was recurrent diarrhea 11/20 (55%). The mean (\pm S.D) age of the cases at the time of diagnosis was 4.2 (\pm 4.1) years. Chronic granulomatous disease (CGD) was the most common 6/20 (30%) disorder, followed by severe combined immunodeficiency (SCID) 3/20 (15%), leukocyte adhesion deficiency (LAD) 3/20 (15%), agammaglobulinemia/hypogammaglobulinemia 3/20 (15%), and Hermansky-Pudlak Syndrome (HPS) 2/20 (10%). Wiskott-Aldrich Syndrome, Immunodeficiency Centromeric Instability and Facial Anomalies Syndrome (ICF 2), Trichohepatoenteric syndrome (TRES), and C3 deficiency were each diagnosed once {1/20 (4.3%) each} (Table 1). Of these 20 confirmed cases, almost all 19/20 (95%) had a family history of consanguinity. Sibling death was reported in 5/20 (25%) of these cases. Five out of the 20 (25%) children died over the 7-year period for various reasons.

Conclusion: PIDs are not uncommon in Pakistan; their diagnosis may be missed or delayed due to the overlapping of clinical features of PID with other diseases and a lack of diagnostic facilities. There is a need to build capacity for early recognition and diagnosis of PIDs to decrease morbidity and mortality.

Keywords: Children, Primary immunodeficiency disorders, Chronic granulomatous disease, Consanguineous marriages

^aDepartment of Pediatrics and Child Health, Aga Khan University, Stadium Road, Karachi, 74800, Pakistan

*Corresponding author. Department of Pediatric & Child Health, Aga Khan University, Stadium Road, P. O Box 3500, Karachi, 74800, Pakistan. E-mail: farah.qamar@aku.edu

Full list of author information is available at the end of the article

<http://doi.org/10.1016/j.waojou.2020.100133>

Received 4 November 2019; Received in revised form 29 April 2020; Accepted 18 May 2020

Online publication date xxx

1939-4551/© 2020 The Authors. Published by Elsevier Inc. on behalf of World Allergy Organization. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

INTRODUCTION

Primary immunodeficiency disorders (PIDs) are a heterogeneous group of genetic disorders characterized by an impaired ability of the immune system to produce a normal immune response. This is due to inherited defects in either cellular or humoral immunity, which results in a spectrum of issues such as recurrent infections, allergies, autoimmunity, and malignancies.^{1,2} In neonates, PIDs often present with severe infections leading to death; whereas in adolescents these infections are less severe albeit recurrent. Diagnosing PIDs is challenging because of the variability in clinical presentation and limited availability of diagnostic tests, particularly in low middle-income countries (LMIC). When diagnostic tests are available, their cost often becomes a limiting factor. Advances in molecular diagnostic techniques and the identification of known gene defects have helped to facilitate the diagnosis of patients with PIDs.³ The true global prevalence and distribution of PIDs remain unclear. The prevalence statistics available from nationwide registries are mostly derived from limited areas of the world. The data obtained from these registries often underestimate the true prevalence, because not all cases are reported to these registries, and due to ambiguity in what constitutes a PID case, some cases are missed. These issues are compounded in developing countries because of the lack of physician training in identification of these disorders and the limited access to diagnostics in these countries. Recent studies have shown that PIDs are more common than previously thought, and that around 1% of the population may have an underlying PID.⁴ The burden of PID varies by region, being highest in the United States of America (USA), followed by Europe, Latin America, Middle East, Asia, and finally Africa.⁴ This frequency may be biased by the availability of resources for diagnosis of these disorders.

Most PIDs are autosomal recessive, which makes it safe to assume that the incidence of PIDs is greater in regions having higher rates of consanguinity. However, limited studies have been carried out in such regions and hence the burden and type of PIDs in such areas is relatively unknown.

Consanguineous marriages are common in developing Asian countries ranging between 20 and 70% of all marriages, and Pakistan is no different, with about 70% of the marriages being consanguineous.⁵⁻⁷ The data on PIDs in Pakistan is currently confined to a few case reports, case series, and editorials; there are no detailed reports on the spectrum of PIDs seen in the country.⁸⁻¹¹

Diagnosing and treating PIDs is a challenge in Pakistan. Patients often die before the disorder is recognized by a physician, leading to a delay in diagnosis or death of the child. Families are often referred to tertiary care centers following the death of a sibling or for admission for life-threatening infections. Even in large tertiary care centers with well-equipped laboratories, there are limited diagnostic facilities available for confirming PIDs. Even when these are available, the diagnostic tests are expensive and beyond the reach of most families as the health care system of Pakistan relies on out-of-pocket payments for health expenditures. Once the diagnosis of PID is confirmed, supportive therapies like antibiotic and antifungal prophylaxis, intravenous (I/V), or subcutaneous (S/C) immunoglobulin therapy, depending on the underlying disorder, are offered. Intravenous immunoglobulin (IVIG) is the standard therapy for most humoral deficiencies, but it is expensive and beyond the reach of most patients in low middle income countries (LMICs). S/C immunoglobulins provide ease of administration, but their availability is limited in LMICs. Bone marrow transplant is available in certain settings but has its limitations in countries like Pakistan where no donor registries are available, and the cost of the transplant coupled with the risk of infections in the post-transplant period make this option challenging to pursue. Gene therapy is another promising albeit costly treatment option undergoing experimentation and is now considered an option for the treatment of multiple non-life-threatening disorders (ie, immunological disorders and systemic protein deficiency). This study aims to report the spectrum of PID cases observed at a tertiary care center in Pakistan, with a focus on the molecular diagnosis in this patient population over the last 7 years.

MATERIALS AND METHODS

The Aga Khan University Hospital (AKUH) is a 700 bed, not-for-profit tertiary care hospital in Karachi, Pakistan. The hospital receives referrals from all over the country. A retrospective chart review of children suspected to have PID was conducted at the AKUH, Pakistan from 2010 to 2016. Data on demographics, clinical features, age at onset of symptoms, age at presentation, history of recurrent infections, skin allergies, family history of consanguinity, sibling death, details related to laboratory workup done for PID (complete blood count, serum immunoglobulin levels, nitro-tetrazolium blue test (NBT), dihydro-rhodamine(DHR) and flow cytometric analysis) were collected. Molecular tests targeted panel next generation sequencing (NGS) or whole exome sequencing (WES) were performed at Geha laboratory at Boston Children's Hospital, USA. Targeted panel NGS was performed using the PID v2 panel and Ion Torrent™ S5 sequencer (ThermoFisher), with an average coverage of 328x. Variant annotation was performed with VarSeq™ software (Golden Helix). Whole exome sequencing (WES) was performed using a previously described pipeline with an average on-target coverage of 80x.¹² The cost of shipping and molecular testing was covered by a philanthropic grant from the Perkins Fund. Data were analyzed using IBM SPSS Statistics v. 20.0. Mean \pm S.D is reported for quantitative variable like age at diagnosis, delay in diagnosis (days), etc. Frequency and percentages are reported for categorical variables such as gender, type of PID, clinical features, history of sibling death, consanguinity in family, mortality, etc.

RESULTS

A total of 43 children visited the hospital with a suspected PID during the study period. Among them, genetic testing was performed in 31/43 (72.1%) children. A confirmed diagnosis was established in 20/43 (46.5%) children. A pathogenic gene variant was identified in 17 (85%) of the 20 confirmed cases (Table 1). Of the remaining 23 patients, samples for 9 patients were not sent for molecular testing, 7 had no known genetic mutation for PID (Table 2), no pathogenic genetic variant was identified in 2 patients, 2 died before

a final diagnosis could be established, 2 were lost to follow-up, and 1 had normal sequencing results (Fig. 1). Twelve (60%) of the 20 confirmed children with PID were male. The mean (\pm S.D) age at the time of diagnosis was 4.2 (\pm 4.1) years. The mean (\pm S.D) delay in diagnosis was 1333 \pm 1420 days. The most common presenting symptom was recurrent diarrhea 11/20 (55%) followed by recurrent pneumonia 8/20 (40%), oral thrush 7/20 (35%), skin rashes and abscesses 6/20 (30%), osteomyelitis 1/20 (5%), and meningitis 1/20 (5%). Chronic granulomatous disease (CGD) was the most common 6/20 (30%) disorder, followed by severe combined immunodeficiency (SCID) 3/20 (15%), leukocyte adhesion deficiency (LAD) 3/20 (15%), agammaglobulinemia/hypogammaglobulinemia 3/20 (15%), and Hermansky-Pudlak Syndrome (HPS) 2/20 (10%). Wiskott-Aldrich Syndrome, Immunodeficiency Centromeric Instability and Facial Anomalies Syndrome (ICF 2), Trichohepatoenteric syndrome (TRES), and C3 deficiency were each diagnosed once {1/20 (4.3%) each} (Table 1). Of these 20 confirmed cases, almost all 19/20 (95%) had a family history of consanguinity. Sibling death was reported in 5/20 (25%) of these cases. Five out of 20 (25%) children died due to various reasons. A pathogen (*Pneumocystis jiroveci*) was identified in only 1 of the 5 deceased patients. Among the deceased children, 2 developed acute respiratory distress syndrome (ARDS), 2 died of sepsis, and 1 had plastic bronchitis.

DISCUSSION

In this case series, 85% of the children were able to get a molecular diagnosis through the support of Boston Children's Hospital. CGD was the commonest disorder observed. This is consistent with data reported from other Asian countries in which phagocytic defects were the most common type of PIDs ranging from 29% to 60%.¹³⁻¹⁵ Almost all cases in this study had a family history of consanguinity. Death of siblings was seen in 25% of the cases. Out of the children who died during the study period, 80% of them were diagnosed after they had died.

Due to a lack of facilities, diagnosing PIDs continues to be a challenge in developing countries. Partnering with international organizations is

S.No	Age/ Sex	Clinical Features	Consan quinity	Sibling Death (Age/ Cause of death)	Special Investigations (Normal range)	Mutated Gene	Final Diag nosis	Age at onset of symptom	Age at diag nosis	Out come
Phagocyte Defects (n = 9)										
1	11 mo/M	Recurrent abscesses and osteomyelitis	Yes	Yes (18 mo old brother & 16 days old brother with no apparent cause)	NBT: Negative DHR: Positive Immunoglobulins IgG:15.2 (6.5-16) IgA: 3.68 (0.4-3.5) IgM:1.15 (0.5-3.0) IgE: not done Flow cytometry Not done HIV-serology: not done	Gene NCF1 Loss of coverage Exon 2 could not be amplified, suggesting a deletion that includes this exon	AR CGD	Birth	6 yr	Alive
2	3.3 yr/M	Recurrent abscesses, lymphadenopathy	No	Yes (9 mo old brother due to pneumonia)	NBT: Negative DHR: Not done Immunoglobulins IgG:23.1 (5.7-17.1) IgA: 3.58 (0.47-2.5) IgM:2.38 (0.64-2.7) IgE: not done Flow Cytometry CD3: 5203 (1500-2900) CD4: 2328 (1000-2100) CD8: 1999 (700-1100) CD19: 2407(500-1200) CD 56: 556 (300-600) HIV-serology: Negative	Gene CYBB Zygosity Homozygous Genomic coordinate c.1222G > A Amino acid Change p.Gly408Arg	X-linked CGD	9 mo	8 yr 9 mo	Alive
3	3 mo/M	Recurrent diarrhea, perianal abscess, skin rash	Yes	No	NBT: Negative DHR: Not done Immunoglobulins IgA: 0.2 (0.09-1.07) Others: not done Flow Cytometry CD3: 1652(1800-3300) CD4: 1224 (900-2300) CD8: 237 (700-1500) CD19: 839(700-1700) CD 56: not done HIV status: not checked	Gene NCF1 Zygosity Homozygous Genomic coordinate c.294_295insA Amino acid change p.Gly99fs	AR CGD	2 mo	2 yr 9 mo	Alive
4	1.3 yr/F	Recurrent diarrhea, cow milk protein allergy	Yes	Yes (1.5 years old sister due to septic shock & had similar complaints, no workup done for PID)	NBT: Absent activity DHR: Not done Immunoglobulins IgG:13.4 (6.5-16.0) IgA:2.49 (0.4-3.5) IgM:1.14 (0.5-3.0) IgE: 1336 (0.4-351.6) Flow cytometry CD3: 5030 (1900-2900) CD4: 3617 (1000-1800) CD8: 1388 (700-1200) CD19: 5073(600-1400) CD56:1279 (300-600) HIV status: not checked	Gene NCF2 Zygosity Homozygous Genomic coordinate Chr1: 183532647 Loss of coverage: 1098_1099delTC = p.Gln367fs (Frameshift deletion)	AR CGD	Birth	2 yr	Alive

5	3 yr/M	Recurrent pneumonias, oral thrush, perianal abscess lymphadenopathy	Yes	No	NBT: Normal DHR: Not done Immunoglobulins IgG: 7.62 (.05-14.0) IgA:1.56((0.18-1.7)) IgM: 1.20(0.63-2.9) IgE: >2000(0.4-351.6) Flow cytometry CD3: 6124(1500-2900) CD4: 3355(1000-2100) CD8: 2539(700-1100) CD19:1830(500-1200) CD56:not done HIV serology: Negative	Gene NCF1 Zygoty Homozygous Genomic coordinate Chr7: 74191664 Amino acid change p.Arg42Trp	AR CGD	2 yr	4 yr 5 mo	Alive
6.	7 yr/M	Recurrent pneumonia	Yes	No	NBT: Not done DHR: Positive Immunoglobulins Not done Flow cytometry Not done HIV status: not checked	Not done	CGD	9 mo	7 yr	Alive
7.	3.5 mo/F	Delayed shedding of umbilical cord, recurrent oral thrush, skin rash,	Yes	No	Immunoglobulins Not done Flow cytometry Not done HIV status: not checked	Gene FERMT3 Zygoty Homozygous Loss of coverage (Kindlin-3) c.126_126delC = p.Ile42fs (Frameshift deletion)	LAD3	2 mo	1yr 4 mo	Alive
8.	7 mo/F	Delayed shedding of umbilical cord, recurrent diarrhea, skin rash	Yes	No	Immunoglobulins IgG:4.02 (6.5-16) IgM:1.36 (0.5-3.0) IgA: 0.36 (0.4-3.5) IgE: 31.0 (1.4-52.3) Flow Cytometry CD3: 3987 (2800-3500) CD4: 3122 (1700-2400) CD8: 784 (800-1100) CD 56: 444 j(300-800) CD19: 2168 (1000-1700) HIV status: not checked	Gene ITGB2 Zygoty: Homozygous Genomic coordinate (GRCh37/hg19): c.134G > A Amino acid change p.Trp45Ter (Nonsense)	LAD1	12 days	5 mo	Alive
9.	4 mo/M	Delayed shedding of umbilical cord and umbilical abscess	Yes	No	Flow cytometry for LAD markers: CD11C deficiency	Not done	LAD2	Birth	3 mo	Alive
Combined/Cellular defects (n = 7)										
10.	4 mo/M	Recurrent diarrhea, skin rash	Yes	No	Immunoglobulins IgG: <0.7 (6.5-16.0) IgA: <0.15 (0.4-3.5) IgM: 0.0 (0.5-3.0) IgE: 440 (1.4-52.3) Flow Cytometry	Gene RAG2 Zygoty Homozygous Genomic coordinate	SCID Omenn syndrome	1.5 mo	5 mo	Alive

(continued)

S.No	Age/ Sex	Clinical Features	Consan guinity	Sibling Death (Age/ Cause of death)	Special Investigations (Normal range)	Mutated Gene	Final Diag nosis	Age at onset of symptom	Age at diag nosis	Out come
11.	5 mo/F	Recurrent pneumonia, diarrhea, otitis media, skin rash	Yes	No	CD3: 150 (1800-3300) CD4: 150 (900-2300) CD8: 1 (700-1500) CD56:108 (300-800) CD19: 0 (700-1700) HIV status: not checked	c.1247G > T Amino acid change p.Trp416Leu	SCID	3 mo	6mo	Died due to ^b ARDS and ^b PCP pneumonia (1 month before establishing diagnosis)
12.	1 yr/F	Recurrent diarrhea, oral thrush	Yes	No	Immunoglobulins IgG:0.66 (1.9-7.9) IgA: <0.15 (0.01-0.59) IgM: not done IgE: 18 (1.4-52.3) Flow Cytometry CD3: 428 (2800-3500) CD4: 283 (1700-2400) CD8: 122 (800-1100) CD56: not done CD19: 25(1000-1700) HIV serology: Negative	Gene JAK3 Amino acid change p.P402L SIFT (0). Polyphen (0.85)	SCID	8 mo	1 yr 2 mo	Died due to ^b ARDS (2 months before establishing diagnosis)
13	2.5 yr/M	Oral thrush, skin rash, Food allergy, anemia, thrombocytopenia	Yes	No	Immunoglobulins Ig G:19.4 (not available) Ig A: 2.99(not available) Ig M: 0.76(not available) Ig E: >5000(not available) Flow Cytometry CD3: 1126(not available) CD4: 285(not available) CD8: 675(not available) CD19: 75(not available) CD56: 299(not available) HIV status: not checked	Gene WAS Zygosity Hemizygous Genomic coordinate ChrX: 48543941 (Deletion of C) Arg94fs (Frameshift deletion)	WAS	9 mo	3 yr 1 mo	Died due to sepsis (5 months before establishing diagnosis)
14.	9 mo/F	Recurrent diarrhea, pneumonia, oral and perianal candidiasis	Yes	No	Immunoglobulins IgG: 4.6 (6.5-16) IgA: <0.33 (0.4-3.5) IgM: 0.21 (0.5-3.0) Ig E: 1.23 (1.4-52.3) Flow cytometry	Gene ZBTB24 Zygosity Homozygous Genomic coordinate	ICF 2	4 mo	1 yr 10 mo	Died due to Plastic bronchitis. (2.5 months after diagnosis)

					CD3: 4286 (1900-2900) CD4: 2942 (1000-1800) CD8: 824 (700-1200) CD56: 331 (300-600) CD19: 944 (600-1400) HIV status: not checked	Chr6: 109802832 c.396_397delTA = p.His132fs (Frameshift deletion)				
15	6 yr/M	Recurrent diarrhea, pneumonia, occulocutaneous albinism, blond hairs, easy bruisibility, oral thrush	Yes	No	Immunoglobulins Ig G: 9.7 (6.5-16) Ig A: 4.0 (0.4-3.5) IgM: 0.28 (0.5-3.0) IgE: 28 (0.5-393) Flow cytometry Not done HIV status: not checked	No pathogenic variant found <i>(Diagnosed on the basis of clinical findings and family history of similar illness in his first cousin)</i>	Hermansky-Pudlack syndrome	Birth	7 yr	Alive
16.	16 y/F	Recurrent diarrhea, pneumonia, occulocutaneous albinism, blond hairs, easy bruisibility, oral thrush, epistaxis, otitis media	Yes	Yes (2.5 years old brother due to pneumonia & had similar features and complaints)	Immunoglobulins Ig G:17.8 (6.5-16) Ig A: 4.48 (0.4-3.5) IgM: 0.45 (0.5-3.0) IgE: 12.0 (1.9-170) Flow cytometry CD3:1330 (720-2320) CD4: 592 (430-1010) CD8: 687 (170-1050) CD19:175 (100-430) CD56: 48 (100-430) HIV serology: Negative	Not done <i>(Diagnosed on the basis of clinical findings and family history of similar illness in her first cousin)</i>	Hermansky-Pudlack syndrome	1 yr	17 yr	Alive
Humoral/Antibody defects (n = 3)										
17.	3 yr/M	Recurrent diarrhea Recurrent abscesses	Yes	No	Immunoglobulins IgG: <0.7 (5.7-17.1) IgM: <0.08 (0.64-2.7) IgA: <0.15 (0.47-2.5) IgE: 51 (0.4-351.6) Flow cytometry CD3: 1444 (1500-2900) CD4: 389 (1000-2100) CD8: 1064 (700-1100) CD19: 2 (500-1200) CD56: 165 (300-600) HIV serology: Negative	Gene BTK Loss of coverage frameshift deletion resulting in exon18 = c.1883delC = p.T628fs	X-linked Agammaglobulinemia	2yr	4yr 9 mo	Died due to sepsis (4 months before establishing diagnosis)
18.	8 yr/F	Recurrent diarrhea and pneumonia	Yes	No	Immunoglobulins IgG:1.11(7.3-15.1) IgM: 0.14(0.55-2.1) IgA: <0.15 (0.7-3.25) IgE: <0.1 (0.5-393) Flow cytometry CD3: 3472(1300-2200) CD4: 1513 (600-1100) CD8: 1787(500-1000) CD19: 49(300-500) CD56:176(300-500) HIV status: not checked	Gene TCF3 Zygosity Homozygous Loss of coverage Total loss of coverage of exons 5-11.	Agammaglobulinemia	1 yr	8 yr 7 mo	Alive

(continued)

S.No	Age/ Sex	Clinical Features	Consan quinity	Sibling Death (Age/ Cause of death)	^a Special Investigations (Normal range)	Mutated Gene	Final Diag nosis	Age at onset of symptom	Age at diag nosis	Out come
19.	1.9 yr/M	Recurrent pneumonia	Yes	Yes (5 years old brother due to measles)	Immunoglobulins: IgG: 9.64 (6.5-16.0) IgM: 0.6 (0.5-3.0) IgA: 0.69 (0.4-3.5) IgE: 3.37 (0.4-351.6) Flow cytometry: CD3:1653 (1500-2900) CD4:1242 (1000-2100) CD8: 427 (700-1100) CD19: 12 (500-1200) CD56: 132 (300-600) HIV status: not checked	Gene TTC37 Zygoty Homozygous Genomic coordinates (GRCh37/ hg19): c.1864A > C Amino acid change p.Thr622Pro Polyphen: 0.89 SIFT: 0.41	Trichohe patoenteric syndrome (THES)	Birth	1 yr 10 mo	Alive
Complement Deficiency (n = 1)										
20.	4.5 yr/M	Recurrent meningitis	Yes	No	Complement Level C3: 0.05 g/L(0.8-1.5)	Gene C3 Zygoty Homozygous Genomic coordinates (GRCh37/ hg19): Chr19: 6711208 C > A (5' Splicesite, Exon 12)	C3 deficiency	1 yr	5 yr	Alive

Table 1. (Continued) Clinical and Laboratory characteristics of patients with a confirmed Primary immunodeficiency disorder. a. CD count unit: cells/mm³, Immunoglobulin unit: G/L, IgE unit: IU/ml, NBT: Nitrotetrazolium blue test, DHR: Dihydrorhodamine. b. ARDS: Acute Respiratory Distress Syndrome, PCP: Pneumocystis carini c. Sample sent to Armed Forces of Institute of Pathology (AFIP), Rawalpindi to check the CD markers for Leukocyte Adhesion Defect (LAD) on flow cytometry

S.No	Age/Sex	Clinical Features	Special Investigations (Normal range)	Suspected diagnosis	Gene Sequence Result	Age at onset of symptom	Age at diagnosis	Outcome
1	10 mo/M	Recurrent diarrhea	Immunoglobulins IgG:<0.7 (1.9-7.9) IgA:<0.15 (0.01-0.59) IgM:<0.08 (0.09-2.1) IgE:1.0 (1.4-52.3) Flow cytometry CD3:166(1800-3300) CD4:132(900-2300) CD8: 27 (700-1500) CD19: 22(700-1700) CD56: 530(300-800) HIV serology: negative	T ⁻ B ⁺ NK ⁻ SCID	No Known candidate gene for PID identified	Birth	10 mo.	Died due to sepsis
2	2.5 yr/M	Work up was done because of family history of SCID.	Immunoglobulins Not done Flow cytometry CD3:503(1500-2900) CD4: 451(1000-2100) CD8: 68(700-1100) CD19:12(500-1200) CD56: not done HIV status: not checked	T ⁻ B ⁻ NK ⁻ SCID	No Known candidate gene for PID identified	10th day of life	1 yr 8 mo.	Died due to sepsis
3	6 mo/M	Fever, leg gangrene, seizures	Immunoglobulins IgG:10.7(6.5-16) IgA: 0.31(0.4-3.5) IgM:0.2(0.5-3) IgE:12.4(1.4-52.3) Flow Cytometry CD3:1140(1800-3300) CD4:735(900-2300) CD8: 449(700-1500) CD56: 49 (300-800) CD19: 273(700-1700) HIV serology: negative	SCID	No Known candidate gene for PID identified	1.5 mo.	4 mo.	Alive No recurrence of infections in follow up. Less likely to be a PID
4	13 yr/M	Recurrent diarrhea and joint swelling	Immunoglobulins IgG: 1.11(5.7-17.1) IgA: 0.41(0.47-2.5) IgM: 0.05(0.64-2.7) IgE: 2.85(1.9-170) Flow Cytometry CD3:1718(1100-2200) CD4: 651(600-1600)	No	No Known candidate gene for PID identified	6 yr.	13 yr.	Alive

(continued)

S.No	Age/Sex	Clinical Features	^a Special Investigations (Normal range)	Suspected diagnosis	Gene Sequence Result	Age at onset of symptom	Age at diagnosis	Outcome
			CD8: 850(500-1200) CD19: 454(200-600) CD56: 83(300-600) HIV status: not checked					
5	8 mo/F	Fever and seizures	Immunoglobulins IgG:13.54(6.5-16) IgA: 0.19 (0.4-3.5) IgM: 2.8(0.5-3) IgE: not done Flow Cytometry CD3:1624(2800-3500) CD4: 592(1700-2400) CD8: 831(800-1100). CD19: 425(1000-1700) CD56: 913(300-800) HIV status: not checked	No	No Known candidate gene for PID identified	4.5 mo.	6 mo.	Alive
6	11 mo/M	Recurrent diarrhea and pneumonia	Immunoglobulins IgG:4.96((6.5-16)) IgM:0.09(0.5-3) IgA: 0.16(0.4-3.5) IgE:1.0(1.4-52.3) Flow Cytometry CD3: 244(1800-3300) CD4: 105(900-2300) CD8: 146(700-1500) CD19:1.0(700-1700) CD56: 89(300-800) HIV status: Negative	SCID	No Known candidate gene for PID identified	4 mo.	11 mo.	Alive No recurrence of infections in follow up. Less likely to be a PID
7	5.5 mo/M	Recurrent pneumonia, skin rash and hyperglycemia	Immunoglobulins IgG:0.26(6.5-16) IgA: 0.09(0.4-3.5) IgM:0.12(0.5-3) IgE:5.96 (1.4-52.3) Flow cytometry CD3: 2244(1800-3300) CD4: 879 (900-2300) CD8: 1167(700-1500) CD56:196 (300-800) CD19: 21 (700-1700) HIV status: not checked	Early onset diabetes and Agamma globulinemia STAT3 GOF? LRBA?	No Known candidate gene for PID identified	2 mo.	5 mo.	Alive Recurrent otitis media, eczematous skin rash and failure to thrive

Table 2. Clinical and laboratory characteristics of suspected primary immunodeficiency patients with No known gene mutation for PID. a. CD count unit: cells/mm³, Immunoglobulin unit: G/L, IgE unit: IU/ml




Fig. 1 Study flow showing distribution of study subjects

crucial in aiding and establishing the diagnosis of suspected PIDs in these countries.⁹

Patients with PIDs are often diagnosed based on a clinical history of recurrent infections due to atypical or less virulent pathogens.¹⁶ However, they can also present with non-infectious manifestations, such as autoimmune disease, or albinism.¹⁷ Verma et al published a case series of 27 PID cases in whom recurrent pneumonia and recurrent diarrhea were the most common presentations. In our case series, recurrent diarrhea was the most common clinical presentation followed by recurrent pneumonia, oral thrush, skin rash, and abscesses.¹⁸

Live vaccines (eg, BCG, OPV) should not be administered to children with PIDs due to the risk of developing mycobacterial disease with the bacillus Calmette–Guérin (BCG) vaccine or Vaccine Derived Poliovirus (VDP) with the oral polio virus vaccine (OPV).¹⁹ Shahmohammadi S et al reviewed 17 cases of disseminated BCG infection in Iran from 2005 to 2010 and observed that 10 (59%) out of the 17 cases had impaired immunity due to an incompetent immune system.²⁰ In Pakistan, these vaccines (BCG and OPV) are a part of the national immunization programme; hence, they are administered routinely, much earlier before a diagnosis of immunodeficiency can be suspected or established in children. In our case series, almost all cases were vaccinated with the BCG and OPV at birth; so far, none of

them have developed any sign of disease due to the vaccination (VDP or Tuberculosis). Although this report is based on a limited number of cases, it does raise the need for a newborn screening program in the long term, so that children suspected of having a PID do not receive a live vaccine at birth.

Early recognition and prompt diagnosis of PIDs help in preventing significant disease related morbidity and mortality. A clinical history and physical examination together with supportive laboratory investigations can provide clues about an underlying PID. Nevertheless, molecular testing plays an essential role in not only confirming the diagnosis but also in identifying the exact nature of the defect. It also assists in patient care as it allows for prognostic counseling.²¹

Varying degree of support for diagnosing PIDs is available from different institutions and philanthropic foundations such as the (International Union of Immunological Sciences (IUIS), European Society for Immunodeficiencies (ESID), Immune Deficiency Foundation (IDF), and Jeffrey Modell Foundation JMF). In most LMICs, diagnostic and therapeutic services are sparse and/or inaccessible. Armed Forces Institute of Pathology (AFIP) in Rawalpindi, Pakistan is a center where limited immune diagnostics are available but molecular techniques are still lacking. Furthermore, this is the only center where some form of diagnostics is available for PIDs. For a country with a population

of 200 million and a birth rate of 29.8 births per 1,000 people, a single facility is not enough to meet the demands.

To our knowledge, this is the first report of PID cases from Pakistan confirmed with molecular diagnosis. The cases reported here only reflect cases referred to a large tertiary care center and represent only the tip of the iceberg.

As a next step, we plan to set up a national registry to bring together individual efforts so that medical care for affected individuals can be improved. Having a national registry will facilitate exchange of experience in the diagnosis and management of PIDs with international registries and will help push policies for support of children with an identified PID. We plan to work towards achieving philanthropic support for IVIG/SCIG or bone marrow transplant (BMT) for the treatment of diagnosed patients. Hopefully, these collective efforts will help with early identification of PIDs, aid in tracking and establishing epidemiological data of PIDs in Pakistan, and help decrease the mortality of patients in developing countries around the world.

CONCLUSION

PIDs are not uncommon in Pakistan; their diagnosis may be missed or delayed due to the overlapping of clinical features of PIDs with other diseases and a lack of diagnostic facilities. There is a need to increase testing capacity for early recognition, diagnosis and management of PIDs to decrease morbidity and mortality. There is also a need to establish a national registry for estimating the true burden of PIDs in the country to aid in policy recommendations.

Abbreviations

PIDs: Primary Immunodeficiency Disorders; NGS: Next-Generation Sequencing; WES: Whole Exome Sequencing; NBT: Nitrotetrazolium blue test; DHR: Dihydrorhodamine; CGD: Chronic Granulomatous Disease; SCID: Severe Combined Immunodeficiency Disorder; LAD: Leukocyte Adhesion Deficiency; HPS: Hermansky-Pudlak Syndrome; ICF-2: Immunodeficiency Centromeric Instability and Facial Anomalies Syndrome; TRES: Trichohepatoenteric syndrome; LMIC: Low Middle Income Countries; USA: United States of America; I/V: Intravenous; S/C: Subcutaneous; ARDS: Acute Respiratory Distress Syndrome; BCG: Bacille Calmette-Guerin; OPV: Oral Polio Vaccine; VDP: Vaccine

Derived Poliovirus; BMT: Bone Marrow Transplant; AFIP: Armed Forces Institute of Pathology

Funding disclosure

There was no funding for this study.

Ethical consideration

The study was exempted by Ethical Review Committee (ERC) of Aga Khan University Hospital Karachi (4652-Ped-ERC-17).

Author's contribution

Sonia Qureshi and Farah Naz Qamar contributed to the study conception and design. Material preparation, data collection and analysis were performed by Sonia Qureshi. Sonia Qureshi, Farah Naz Qamar, Fatima Mir, Samina Junejo, Khalid Saleem, Samreen Zaidi and Khalil Ahmad were involved in patient care and management of PID cases. The first draft of the manuscript was written by Sonia Qureshi and all authors commented on the previous manuscript. Abdullah. B. Naveed contributed in the revised version of the manuscript. All authors read and approved the final manuscript.

Availability of data and materials

The data set is available from Farah Naz Qamar.

Consent for publication

All authors agreed to the publication of this work.

Declaration of Competing Interest

All the authors declare that they have no conflict of interest, whether financial or otherwise.

Acknowledgement

We are grateful to The Jeffrey Modell Foundation and Prof. Geha's lab and his team members from the Immunology Division, Boston Children's Hospital, for their continuous support in diagnosing these patients and for performing the molecular testing. We would also like to acknowledge Col. Hamid Nawaz Tipu and his team at the Department of Immunology, Armed Forces Institute of Pathology, and Rawalpindi, Pakistan for providing laboratory support for our patients.

Author details

^aDepartment of Pediatrics and Child Health, Aga Khan University, Stadium Road, Karachi, 74800, Pakistan.

^bDepartment of Pediatrics, The Indus Hospital, Korangi Road, Karachi, Pakistan. ^cChildren's Hospital and The Institute of Child Health, Multan, Pakistan. ^dNational Institute of Blood Disease & Bone Marrow Transplantation, P.E.C.H.S, Karachi, Pakistan. ^eMedical College, Aga Khan University, Stadium Road, Karachi, 74800, Pakistan.

REFERENCES

1. Renzi S, Langenberg-Verregaert KPS, Waespe N, et al. Primary immunodeficiencies and their associated risk of malignancies in children: an overview. *Eur J Pediatr.* 2020;1-9.
2. Raje N, Dinakar C. Overview of immunodeficiency disorders. *Immunol Allergy Clin.* 2015;35(4):599-623.
3. Bousfiha A, Jeddane L, Picard C, et al. The 2017 IUIS phenotypic classification for primary immunodeficiencies. *J Clin Immunol.* 2018;38(1):129-143.
4. Modell V, Orange JS, Quinn J, Modell F. Global report on primary immunodeficiencies: 2018 update from the Jeffrey Modell Centers Network on disease classification, regional trends, treatment modalities, and physician reported outcomes. *Immunol Res.* 2018;66(3):367-380.
5. Al-Herz W. Primary immunodeficiency disorders in Kuwait: first report from Kuwait national primary immunodeficiency registry (2004-2006). *J Clin Immunol.* 2008;28(2):186-193.
6. Aghamohammadi A, Mohammadinejad P, Abolhassani H, et al. Primary immunodeficiency disorders in Iran: update and new insights from the third report of the national registry. *J Clin Immunol.* 2014;34(4):478-490.
7. Al-Saud B, Al-Mousa H, Al Gazlan S, et al. Primary immunodeficiency diseases in Saudi arabia: a tertiary care hospital experience over a period of three years (2010-2013). *J Clin Immunol.* 2015;35(7):651-660.
8. El-Sayed ZA, Abramova I, Aldave JC, et al. X-linked agammaglobulinemia (XLA):Phenotype, diagnosis, and therapeutic challenges around the world. *World Allergy Organ J.* 2019;12(3):100018.
9. Wallace JG, Tipu HN, Stafstrom K, et al. Rethinking newborn screening for severe combined immunodeficiency: lessons from an international partnership for patients with primary immunodeficiencies in Pakistan. *Clin Immunol.* 2019;202:29-32.
10. Qamar F, Junejo S, Qureshi S, et al. A novel mutation in the JH4 domain of JAK3 causing severe combined immunodeficiency complicated by vertebral osteomyelitis. *Clin Immunol.* 2017;183:198-200.
11. Qureshi S, Sheikh MDA, Qamar FN. Autosomal Recessive Agammaglobulinemia - first case with a novel TCF3 mutation from Pakistan. *Clin Immunol.* 2019;198:100-101.
12. Seleman M, Hoyos-Bachiloglu R, Geha RS, Chou J. Uses of next-generation sequencing technologies for the diagnosis of primary immunodeficiencies. *Front Immunol.* 2017;8:847.
13. de Silva NR, Gunawardena S, Rathnayake D, Wickramasingha GD. Spectrum of primary immunodeficiency disorders in Sri Lanka. *Allergy Asthma Clin Immunol.* 2013;9(1):50.
14. Shah I. Hyper IgM syndrome with tuberculous osteomyelitis and scrofuloderma. *Indian Pediatr.* 2005;42(9):952-953.
15. Modell V, Knaus M, Modell F, Roifman C, Orange J, Notarangelo LD. Global overview of primary immunodeficiencies: a report from Jeffrey Modell Centers worldwide focused on diagnosis, treatment, and discovery. *Immunol Res.* 2014;60(1):132-144.
16. Lederman HM. The clinical presentation of primary immunodeficiency diseases. *Clin Focus Prim Immunol Def.* 2000;2(1).
17. Azizi G, Ahmadi M, Abolhassani H, et al. Autoimmunity in primary antibody deficiencies. *Int Arch Allergy Immunol.* 2016;171(3-4):180-193.
18. Verma S, Sharma PK, Sivanandan S, et al. Spectrum of primary immune deficiency at a tertiary care hospital. *Indian J Pediatr.* 2008;75(2):143-148.
19. Sobh A, Bonilla FA. Vaccination in primary immunodeficiency disorders. *J Allergy Clin Immunol Pract.* 2016;4(6):1066-1075.
20. Shahmohammadi S, Saffar MJ, Rezai MS. BCG-osis after BCG vaccination in immunocompromised children: case series and review. *J Pediatr Rev.* 2014;2(1):62-74.
21. Ameratunga R, Woon ST, Neas K, Love DR. The clinical utility of molecular diagnostic testing for primary immune deficiency disorders: a case based review. *Allergy Asthma Clin Immunol.* 2010;6(1):12.